

The Delta Smelt Working Group and the Delta Smelt Risk Assessment Matrix

The delta smelt risk assessment matrix (DSRAM) consists of month by month criteria which, when exceeded will trigger a meeting of the Delta Smelt Working Group (Working Group). The Working Group consists of experts in delta smelt biology from the U.S. Fish and Wildlife Service (Service), U.S. Bureau of Reclamation (Reclamation), U.S. Environmental Protection Agency (EPA), and California Departments of Water Resources (DWR) and Fish and Game (DFG). The purpose of the DSRAM is to take actions to protect delta smelt in a proactive manner prior to salvage events at the Federal and State Delta export facilities. Reclamation and/or DWR are responsible for monitoring the DSRAM criteria and reporting back to the Service and the Working Group. The DSRAM has been modified from the delta smelt decision tree, which was peer-reviewed and presented in the Interagency Ecological Program (IEP) Newsletter. It is the intent of the Service that the DSRAM be sent out for independent peer review. The DSRAM is an adaptive management tool which may be further modified by the Working Group and the interagency Water Management Operations Team (WOMT) as new information becomes available, without undergoing formal reconsultation. An informative link to an existing website will be developed that compiles monitoring data from IEP and DFG to enable members of the Working Group to easily track the progress of the triggering criteria. Data will be updated at least weekly to determine the need for a meeting.

Should a triggering criterion be met or exceeded, Reclamation and/or DWR will inform the members of the Working Group and the Working Group will determine the need to meet. Any member of the Working Group may initiate a meeting of the Working Group at any time. A meeting of the Working Group may consist of an in-person meeting, a conference call, or a discussion by email. If needed, the Working Group will meet prior to the weekly meetings of the DAT and the WOMT and information will be shared with these groups. The Working Group will be available to present management briefings as needed.

If a meeting of the Working Group proves necessary, the group will review the available monitoring and survey data and decide whether to recommend a change in water project operations (referred to as "fish actions"). These potential fish actions are listed in the DSRAM by the months wherein each of these tools generally becomes available. Generally, if the Working Group recommends a fish action, it will be shared with the Data Assessment Team (DAT) during its weekly conference call and forwarded to the WOMT for discussion and potential implementation; however, the Working Group may make recommendations to WOMT at any time. Recommendations will include a discussion of the level of concern for delta smelt and will include a list of the participants in the Working Group discussions. All dissenting opinions and/or discussion points will also be forwarded to the WOMT. The Working Group will meet at least weekly throughout the period in which the triggering criteria are met or exceeded, to determine the need to provide further recommendations to the WOMT.

Notes and findings of Working Group meetings will be submitted to the Service and members of the WOMT for their records. Meeting notes will also be available to the public on the Sacramento Fish and Wildlife Office's web page. The WOMT will respond to the Working Group's recommendations and the actions taken by the WOMT will be summarized by Reclamation and/or DWR annually and submitted to all WOMT agencies.

If an action is taken, the Working Group will follow up on the action to attempt to ascertain its effectiveness. An assessment of effectiveness will be attached to the notes from the Working Group's discussion concerning the action.

Life Stage	Adults	Adults	Adults	Adults and larvae	Adults and larvae	Larvae and juveniles	Larvae and juveniles	Juveniles
Previous Year's Fall Midwater Trawl Recovery Index (1)	Index below 74	Index below 74	Index below 74	Index below 74	Index below 74	Index below 74	Index below 74	Index below 74
Risk of Entrainment (2)				X2 upstream of Chipps Island and temps are $\geq 12^{\circ}$	X2 upstream of Chipps Island and temps are between 12° and 18°C	X2 upstream of Chipps Island and mean delta-wide temps $<18^{\circ}\text{C}$ and south delta temps below 25°C	X2 upstream of Chipps Island and temps are below 25°C	X2 upstream of Chipps Island and temps are below 25°C
Duration of Spawning period (number of days temperatures are between 12 and 18°C) (3)					39 days or less by April 15	50 days or less by May 1		
Spawning Stage as determined by spring Kodiak trawl and/or salvage (4)			Presence of Adults at spawning stage ≥ 4	Adult spawning stage ≥ 4	Adult spawning stage ≥ 4			
smelt distribution (5)	See footnote #5	See footnote #5	See footnote #5	See footnote #5 or negative 20mm centroid or low juvenile abundance	Negative 20mm centroid or low juvenile abundance	Negative 20mm centroid or low juvenile abundance	Negative 20mm centroid or low juvenile abundance	Negative 20mm centroid or low juvenile abundance
Salvage Trigger (6)	Adult concern level calculation	Adult concern level calculation	Adult concern level calculation	Adult concern level calculation		If salvage is above zero	If salvage is above zero	

Tools for Change (7)	December	January	February	March	April	May	June	July
Export reduction at one or both facilities	X	X	X	X	X	X	X	X
Change in barrier operations						X	X	X
Change in San Joaquin River flows				X	X	X	X	X
Change position of cross channel gates						X	X	

Delta Smelt Risk Assessment Matrix Footnotes (note: the references for the DSRAM are also included in the literature cited section of the biological opinion)

- 1 The Recovery index is calculated from a subset of the September and October Fall Midwater Trawl sampling (<http://www.delta.dfg.ca.gov/>). The number in the matrix, 74, is the median value for the 1980-2002 Recovery Index (Figure DS1)

The temperature range of 12 to 18 degrees Celsius is the range in which most successful delta smelt spawning occurs. This has been analyzed by using observed cohorts entering the 20-mm Survey length frequency graphs (1996-2002). Cohorts were defined by having a noticeable peak or signal and occurring over three or more surveys during the rearing season. Back calculations were made using the first survey of that cohort with fish less than 15 mm fork length. Temperature data from IEP's HEC-DSS Time Series Data web site was compiled using three stations representing the south Delta (Mossdale), confluence (Antioch), and north Delta (Rio Vista) and averaged together. Spawning dates for each cohort were back-calculated by applying an average daily growth rate (wild fish) of 0.45 mm/day (Bennett, DFG pers. comm.) and egg incubation period of 8-14 days (Baskerville-Bridges, Lindberg pers. comm.) (Mager et al. 2004) from the median value of the analyzed cohort. Each spawning event was then plotted against temperature over time (Figure DS2.1). While spawning does occur outside of the 12-18 degree range, larval survival is most likely reduced when temperatures are either below (DFG pers. comm.) or above this range (Baskerville-Bridges & DFG pers. comm.).

Critical thermal maxima for delta smelt was reached at 25.4 degrees Celsius in the laboratory (Swanson et al., 2000); and at temperatures above 25.6 degrees Celsius smelt are no longer found in the delta (DFG, pers. comm.).

Websites for the temperature data: <http://iep.water.ca.gov/cgi-bin/dss/dss1.pl?station=RSAN007>

<http://iep.water.ca.gov/cgi-bin/dss/dss1.pl?station=RSAN087>

<http://iep.water.ca.gov/cgi-bin/dss/dss1.pl?station=RSAC101>

Mager RC, Doroshov SI, Van Eenennaam JP, and Brown RL. 2004. Early Life Stages of Delta Smelt. American Fisheries Society Symposium 39:169-180.

Swanson C, Reid T, Young PS, and Cech JJ. 2000. Comparative environmental tolerances of threatened delta smelt (*Hypomesus transpacificus*) and introduced Wakasagi (*H. nipponensis*) in an altered California estuary. *Oecologia* 123:384-390.

- 3 Figure DS3: The working hypothesis for delta smelt is that spawning only occurs when temperatures are suitable during the winter and spring. In years with few days having suitable spawning temperatures, the spawning "window" is limited, so the species produces fewer cohorts of young smelt. When there are fewer

cohorts the risk that mortality sources such as entrainment may substantially reduce population size increases. The figures below were used to help define years when there were relatively few days with suitable temperatures. For April 15 and May 1, the figures show the cumulative spawning days for each year during 1984-2002. The cumulative spawning days for each year were calculated based on the number of days that the mean water temperature for three Delta stations (Antioch; Mossdale and Rio Vista) was in the 12 - 18 C range starting on February 1. The results are plotted in terms of the ranks to identify the lower quartile. In other words, years in the lower quartile represent examples of years with relatively few spawning days.

- 4 The adult spawning stage is determined by the Spring Kodiak Trawl and/or fish collected at the salvage facilities (<http://www.delta.dfg.ca.gov/>). A stage greater than or equal to 4 indicates female delta smelt are ripe and ready to spawn or have already spawned (Mager 1996).

Mager RC. 1996. Gametogenesis, Reproduction and Artificial Propagation of Delta Smelt, *Hypomesus transpacificus*. [Dissertation] Davis: University of California, Davis. 115 pages. Published.

- 5 The spring kodiak trawl will be used to generally evaluate the distribution of adult delta smelt. However, since the spring kodiak trawl is not intended to be a survey for abundance or distribution, no definitive trigger for concern can be determined at this time.

Juveniles (March-July) – distribution of juvenile delta smelt where the centroid is located upstream (negative) or downstream (positive) of the Sacramento-San Joaquin River confluence (Sacramento RKI 81; Figure DS5.1). The 20-mm Survey centroid is calculated by multiplying the observed delta smelt station CPUE (fish/10,000 m³) by a distance parameter in km from Sacramento RKI 81. The summed result (summed over a survey) is divided by the survey CPUE which gives the survey centroid position (Figure DS5.2).

Low juvenile abundance will also be a trigger. When juvenile abundance is low, concern is high. Low abundance is indicated when the total cumulative catch in the 20-mm Survey is less than or equal to the 1995-2003 median value of cumulative 20-mm Survey catch for the same surveys (Table DS5).


- 6 Adult salvage trigger: the adult delta smelt salvage trigger period is December through March and the trigger is calculated as the ratio of adult delta smelt salvage to the fall MWT index. This ratio will increase as fish are salvaged during the winter months. If the ratio exceeds the median ratio observed during December-March 1980-2002, then the trigger has been met (see Figure DS6 for more explanation of the calculation)

Juvenile salvage trigger: During May and June, if delta smelt salvage at the SWP/CVP facilities is greater than zero, then the working group will meet. This

is because May and June are the peak months of delta smelt salvage and salvage densities cannot be predicted. Therefore, during these two months, the delta smelt working group expects to meet regularly to look at relevant information such as salvage, delta temperatures, delta hydrology and delta smelt distribution and decide whether to recommend proactive measures to protect these fish.

- 7 The tools for change are actions that the working group can recommend to the WOMET to help protect delta smelt. Exports may be reduced at one or both of the south delta export facilities and a proposed duration of the reduction would be recommended by the working group. Export reductions and changes in San Joaquin River flows may be covered by B(2) or EWA assets. Details of past fish actions can be found at the Calfed Ops website:
<http://www.woco.water.ca.gov/calfedops/index.html>; >Operations [year]

Figure DS1



Points are labeled with the year representing the recovery index. The winter salvage for this analysis starts on December 1 of the recovery index year and continues through March 31 of the following year.

Figure DS2.1. Successful delta smelt spawning periods (shaded blue area) and cohorts (black bars) plotted against water temperature (1996-2002). Spawning periods and cohorts were back calculated using 20-mm Survey catch data. Start of spawning season uses an egg incubation period of 14 d and a growth rate of 0.45 mm/day and end of spawning season 8 d with a growth rate of 0.45 mm/day. Black bars represent the range of 8-14 d egg incubation with a growth rate of 0.45 mm/day from laboratory results. Temperature data ($^{\circ}\text{C}$) was compiled from IEP's HEC-DSS Time Series Data using mean daily temperatures from the confluence (Antioch), south Delta (Mossdale), north Delta (Rio Vista) and averaged together.




Figure DS2.1 cont.




Figure DS2.1 cont.




Figure DS3.





Figure DS5.1. A 20-mm Survey delta smelt bubble plot map with calculated centroid position from the confluence of Sacramento-San Joaquin Rivers with one standard deviation.




Figure DS5.2. Historic juvenile centroid position (20-mm Survey) with one standard deviation.




Table DS5. Median values of cumulative catch from the 20-mm Survey. When cumulative catch per survey during a season is at or below the calculated value, concern is high.

	survey 1	survey 2	survey 3	survey 4	survey 5	survey 6	survey 7	survey 8
Median Value	12	40	144	188	346	500	924	1019

Figure DS6


The objective is to quantify a level of concern for adult delta smelt during the winter that is based upon the number of fish salvaged and the overall abundance of delta smelt. Our trigger reflects that when abundance is low and salvage is high concern is high, and conversely when abundance is high and salvage is low that concern is low.

Below is a Quantile plot of the ratio of winter salvage to the MWT recovery index ($\ln(\text{winter salvage}/\text{recovery index})$). Winter salvage is defined as the total salvage from December through March. In the figure below, the size of the bubbles is proportional to the log of the fall midwater trawl to demonstrate that concern may be high in years of

high or low fall abundance. The resulting quartiles of the ratio are as follows: 25% = 2.950; 50% = 3.575; 75% = 5.029.

Using this approach to calculate winter concern levels, all years above the 1999 point in the graph would have been years of concern. In other words, these are the years in which we may have recommended some protection. Comparing it to the protection afforded adult delta smelt in the winter by the 1995 Biological Opinion: "red light" was, or would have been, reached in fewer winters (1980, 1981, 1982, 1984 and 1999).

The median was selected as the measure of concern and will be calculated by:
 concern level = $\text{anti ln}(3.575) * \text{Recovery index}$



The goal for the DSRAM is to avoid the upper quartile of the above graph, which the Working Group thinks will avoid salvage events that are high relative to fall abundance. Actions may be taken prior to major salvage events.