References

- Anderson CM, Zhu SH. The California Smokers' Helpline and low-SES smokers. Paper presented at the National Conference on Tobacco or Health, San Francisco, CA, November 19–21, 2002.
- Anderson CM, Zhu SH. *The California Smokers' Helpline: A Case Study.* Sacramento, CA: California Department of Health Services, 2000.
- Bailey L. The alliance for tobacco cessation: lessons learned about expanding coverage for evidencebased tobacco dependence treatment. Paper presented at the Ninth Annual Meeting of the Society for Research on Nicotine and Tobacco, New Orleans, LA, February 19–22, 2003.
- Borland R, Balmford J, Hunt D. The effectiveness of personally tailored computer-generated advice letters for smoking cessation. *Addiction* 2004;99(3):369–377.
- Borland R, Segan CJ, Livingston PM, Owen N. The effectiveness of callback counselling for smoking cessation: a randomized trial. *Addiction* 2001;96(6):881–889.
- Burns DM. Smoking cessation: recent indicators of what's working at a population level. In: National Cancer Institute. *Population Based Smoking Cessation: Proceedings of a Conference on What Works to Influence Cessation in the General Population*. Smoking and Tobacco Control Monograph No. 12. Bethesda, MD: U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute, NIH Pub. No. 00-4892, November 2000:99–128.
- Centers for Disease Control and Prevention, Office on Smoking and Health. Unpublished quitline data.
- Centers for Disease Control and Prevention. Annual smoking-attributable mortality, years of potential life lost, and economic costs—United States, 1995–1999. *Morbidity and Mortality Weekly Report* 2002a;51(14):300–303.
- Centers for Disease Control and Prevention. Cigarette smoking among adults—United States, 2000. *Morbidity and Mortality Weekly Report* 2002b;51(29):642–645.
- Centers for Disease Control and Prevention. *Best Practices for Comprehensive Tobacco Control Programs*—August 1999. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1999.

- Centers for Disease Control and Prevention. Physician and other health-care professional counseling of smokers to quit—United States, 1991. *Morbidity and Mortality Weekly Report* 1993a;42(44);854–857.
- Centers for Disease Control and Prevention. Use of smokeless tobacco among adults—United States, 1991. *Morbidity and Mortality Weekly Report* 1993b;42(14):263–266.
- Cummins SE, Zhu SH, Anderson CM. Telephone quitlines for high-risk or hard-to-reach populations. Paper presented at the National Conference on Tobacco or Health, San Francisco, CA, November 19–21, 2002.
- Curry S, McBride C, Grothaus LC, et al. A randomized trial of self-help materials, personalized feedback, and telephone counseling with nonvolunteer smokers. *Journal of Consulting and Clinical Psychology* 1995;63(6):1005–1014.
- Earle R. *The Art of Cause Marketing: How to Use Advertising to Change Personal Behavior and Public Policy.* Lincolnwood, IL: NTC Business Books, 2000.
- Farrelly MC, Pechacek TF, Chaloupka FJ. The impact of tobacco control program expenditures on aggregate cigarette sales: 1981–2000. *Journal of Health Economics* 2003;22(5):843–859.
- Fiore MC, Croyle RT, Curry SJ, et al. Preventing 3 million premature deaths and helping 5 million smokers quit: a national action plan for tobacco cessation. *American Journal of Public Health* 2004;94(2):205–210.
- Fiore MC, Bailey WC, Cohen SJ, et al. *Treating Tobacco Use and Dependence*. Clinical Practice Guideline. Rockville, MD: U.S. Department of Health and Human Services, Public Health Services, June 2000.
- Fishbein HA, Unger JB, Johnson CA, et al. Interaction of population-based approaches for tobacco control. In: National Cancer Institute. *Population Based Smoking Cessation: Proceedings of a Conference on What Works to Influence Cessation in the General Population*. Smoking and Tobacco Control Monograph No. 12. Bethesda, MD: U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute, NIH Pub. No. 00-4892, November 2000:223–233.
- Flay BR. Efficacy and effectiveness trials (and other phases of research) in the development of health promotion programs. *Preventive Medicine* 1986;15(5):451–474.
- Floyd RL, Rimer BK, Giovino GA, et al. A review of smoking in pregnancy: effects on pregnancy outcomes and cessation efforts. *Annual Review of Public Health* 1993;14:379–411.
- Greenwald P, Cullen JW. The new emphasis in cancer control. *Journal of the National Cancer Institute* 1985;74(3):543–551.
- Hatsukami DK, Severson HH. Oral spit tobacco: addiction, prevention and treatment. *Nicotine & Tobacco Research* 1999;1(1):21–44.

- Hollis J, Polen M, Whitlock E, et al. Efficacy of a brief tobacco prevention and cessation program for teens seen in routine medical care (Teen REACH). Paper presented at the National Cancer Institute Youth Tobacco Research Meeting, Palm Harbor, FL, June 2002.
- Hopkins DP, Briss PA, Ricard CJ, et al. Reviews of evidence regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke. *American Journal of Preventive Medicine* 2001;20(Suppl 2):16–66.
- Hughes JR, Keely JP, Niaura RS, et al. Measures of abstinence in clinical trials: issues and recommendations. *Nicotine & Tobacco Research* 2003;5(1):13–25.
- Lando HA, Hellerstedt WL, Pirie PL, et al. Brief supportive telephone outreach as a recruitment and intervention strategy for smoking cessation. *American Journal of Public Health* 1992;82(1):41–46.
- Lichtenstein E, Glasgow RE. Smoking cessation: what have we learned over the past decade? *Journal* of Consulting and Clinical Psychology 1992;60(4):518–527.
- Lichtenstein E, Zhu SH, McAfee T. Smoking cessation quitlines: fusing research and practice. Symposium presented at the 24th Annual Meeting of the Society for Behavioral Medicine, Salt Lake City, UT, March 19–22, 2003.
- Lichtenstein E, Glasgow RE, Lando HA, et al. Telephone counseling for smoking cessation: rationales and meta-analytic review of evidence. *Health Education Research* 1996;11(2):243–257.
- McAfee T. Increasing the population impact of quitlines. Paper presented at the North American Quitline Conference, Phoenix, AZ, May 8–10, 2002.
- McAfee T, Sofian NS, Wilson J, et al. The role of tobacco intervention in population-based health care: a case study. *American Journal of Preventive Medicine* 1998;14(Suppl 3):46–52.
- McAlister AL, Rabius V, Geiger A, et al. Telephone assistance for smoking cessation: one year cost effectiveness estimations. *Tobacco Control* 2004;13(1):85–86.
- McDonald P, Colwell B, Backinger CL, et al. Better practices for youth tobacco cessation: evidence of review panel. *American Journal of Health Behavior* 2003;27(Suppl 2):S144–S158.
- Melvin CL, Dolan-Mullen P, Windsor RA, et al. Recommended cessation counselling for pregnant women who smoke: a review of the evidence. *Tobacco Control* 2000;9(Suppl III):iii80–iii84.
- Mermelstein R. Youth smoking cessation. Paper presented at Tobacco Control Research: Investing in Science for the Public's Health. Washington, DC, June 17–19, 2003.
- Niaura R, Abrams DB. Smoking cessation: progress, priorities, and prospectus. *Journal of Consulting and Clinical Psychology* 2002;70(3):494–509.
- New Jersey Comprehensive Tobacco Control Program. *2001 Annual Report.* New Jersey Department of Health and Senior Services, 2001. Available at http://www.state.nj.us/health/as/ctcp/annualreport.htm. Accessed March 24, 2004.

- Orleans CT, Melvin CL, Marx JF, et al. National action plan to reduce smoking during pregnancy: the National Partnership to Help Pregnant Smokers Quit. *Nicotine & Tobacco Research* 2004; 6(Suppl 2):S269–S277.
- Orleans CT, Schoenbach VJ, Wagner EH, et al. Self-help quit smoking interventions: effects of selfhelp materials, social support instructions, and telephone counseling. *Journal of Consulting and Clinical Psychology* 1991;59(3):439–448.
- Ossip-Klein DJ. The concept of a consortium for quitlines. Paper presented at the North American Quitline Conference, Phoenix, AZ, May 8–10, 2002.
- Ossip-Klein DJ, McIntosh S. Quitlines in North America: evidence base and applications. *American Journal of Medical Science* 2003;326(4):201–205.
- Ossip-Klein DJ, Giovino GA, Megahed N, et al. Effects of a smoker's hotline: results of a 10-county self-help trial. *Journal of Consulting and Clinical Psychology* 1991;59(2):325–332.
- Owen L. Impact of a telephone helpline for smokers who called during a mass media campaign. *Tobacco Control* 2000;9(2):148–154.
- Pacific Center on Health and Tobacco. Linking a Network: Integrating Quitlines with Health Care Systems, 2003. Available at http://www.paccenter.org. Accessed March 24, 2004.
- Padgett DI, Zhu SH, Sun J, Strychacz C. Characteristics of smokeless tobacco users calling a statewide tobacco cessation quitline. Paper presented at the National Conference on Tobacco or Health, San Francisco, CA, November 19–21, 2002.
- Peto R, Darby S, Deo H, et al. Smoking, smoking cessation, and lung cancer in the UK since 1950: combination of national statistics with two case-control studies. *British Medical Journal* 2000;321(7257):323–329.
- Powers P. Impact of media on the Arizona Smokers' Helpline: three-year comparison. 11th World Conference on Tobacco or Health, Chicago, August 6–11, 2000. Available at http://www.tepp.org/presentations. Accessed March 24, 2004.
- Powers P, Wentzel TM, Ranger-Moore J. Impact of pharmacotherapy usage and social support on quit rates for Hispanic clients of the Arizona Smokers' Helpline. Paper presented at the National Conference on Tobacco or Health, New Orleans, LA, November 27–29, 2001. Available at http://128.196.174.132/bigfiles/hispanics-NCTOH2001.ppt. Accessed March 24, 2004.
- Powers P, Wentzel TM, Ranger-Moore J, et al. Attracting motivated quitters to a smokers' helpline using television advertising. Poster presented at the 21st Annual Meeting of the Society for Behavioral Medicine, Nashville, TN, April 5–8, 2000a. Available at http://www.tepp.org/presentations/chuck2000/index.htm. Accessed March 24, 2004.
- Powers P, Ranger-Moore J, Wentzel TM, et al. Impact of television advertising on helpline clients' stages of change. Poster presented at the Sixth Annual Meeting of the Society for Research on Nicotine and Tobacco, Arlington, VA, February 18–20, 2000b. Available at http://www.tepp.org/presentations/Impact_2000/index.htm. Accessed March 24, 2004.

- Powers P, Wentzel TM, Ranger-Moore J, et al. Effects of television advertising on tobacco use cessation helpline calls. Poster presented at the 127th Annual Meeting of the American Public Health Association, Chicago, November 7–11, 1999.
- Rosen E. The Anatomy of Buzz New York: Doubleday, 2000.
- Schroeder SA. Thirty seconds to save a life: what busy clinicians can do to help their patients quit smoking. Presented at a meeting of BlueCross/BlueShield Association executives, San Francisco, CA, November 12, 2003. Available at http://smokingcessationleadership.ucsf.edu. Accessed April 10, 2004.
- Severson HH, Andrews JA, Lichtenstein E, et al. A self-help cessation program for smokeless tobacco users: comparison of two interventions. *Nicotine & Tobacco Research* 2000;2(4):363–370.
- Stead LF, Lancaster T, Perera R. Telephone counselling for smoking cessation (Cochrane Review). In: *The Cochrane Library 2004*, Issue 1. Chichester, UK: John Wiley & Sons, Ltd, 2004.
- Stevens C. Designing an effective counteradvertising campaign—California. *Cancer* 1998;83 (Suppl 12):2736–2741.
- Stevens VJ, Glasgow RE, Hollis JF, Mount K. Implementation and effectiveness of a brief smoking-cessation intervention for hospital patients. *Medical Care* 2000;38(5):451–459.
- Swan GE, McAfee T, Curry SJ, et al. Effectiveness of bupropion sustained release for smoking cessation in a health care setting: a randomized trial. *Archives of Internal Medicine* 2003;163(19):2337–2344.
- Taylor DH, Hasselblad V, Henley SJ, et al. Benefits of smoking cessation for longevity. *American Journal of Public Health* 2002;92(6):990–996.
- U.S. Department of Health and Human Services. *Healthy People 2010.* 2nd ed. With Understanding and Improving Health and Objectives for Improving Health. 2 vols. Washington, DC: U.S. Department of Health and Human Services, 2000a.
- U.S. Department of Health and Human Services. *Reducing Tobacco Use: A Report of the Surgeon General.* Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Office on Smoking and Health, 2000b.
- U.S. Department of Health and Human Services. *Tobacco Use Among U.S. Racial/Ethnic Minority Groups—African Americans, American Indians and Alaska Natives, Asian Americans and Pacific Islanders, and Hispanics: A Report of the Surgeon General.* Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1998.
- Velicer WF, Prochaska JO, Rossi JS, Snow MG. Assessing outcome in smoking cessation studies. *Psychological Bulletin* 1992;111(1):23–41.
- Waa A, Glasgow H, McCulloch M. Subsidisation of nicotine replacement therapies through the New Zealand Quitline. Paper presented at the Second European Quitline Conference, Barcelona, Spain, September 6–8, 2000.

References

- Wakefield M, Borland R. Saved by the bell: the role of telephone helpline services in the context of mass-media anti-smoking campaigns. *Tobacco Control* 2000;9(2):117–119.
- Weinreich NK. Hands-On Social Marketing. Thousand Oaks, CA: Sage Publications, 1999.
- Zhu SH. A quitline for teen smokers. Paper presented at the Youth Tobacco Research Meeting, Washington, DC, June 17–19, 2003.
- Zhu SH. A survey of quitlines in North America. Paper presented at the North America Quitline Conference, Phoenix, AZ, May 8–10, 2002a.
- Zhu SH. Special populations: adolescents, pregnant women, SLT users. Paper presented at the North American Quitline Conference, Phoenix, AZ, May 8–10, 2002b.
- Zhu SH. Telephone quitlines for smoking cessation. In: National Cancer Institute. Population Based Smoking Cessation: Proceedings of a Conference on What Works to Influence Cessation in the General Population. Smoking and Tobacco Control Monograph No. 12. Bethesda, MD: U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute, NIH Pub. No. 00–4892, November 2000:189–198.
- Zhu SH, Anderson CM, Johnson CE, et al. A centralized telephone service for tobacco cessation: the California experience. *Tobacco Control* 2000;9(Suppl II):ii48–ii55.
- Zhu SH, Anderson CM. Bridging the clinical and public health approaches to smoking cessation: California Smokers' Helpline. In: Jamner MS, Stokols D, eds. *Promoting Human Wellness: New Frontiers for Research, Practice, and Policy.* Berkeley, CA: University of California Press, 2000:378–394.
- Zhu SH, Anderson CM, Tedeschi GJ, et al. Evidence of real-world effectiveness of a telephone quitline for smokers. *New England Journal of Medicine* 2002;347:1087–1093.
- Zhu SH, Stretch V, Balabanis M, et al. Telephone counseling for smoking cessation: effects of singlesession and multiple-session interventions. *Journal of Consulting and Clinical Psychology* 1996;64(1):202–211.
- Zhu SH, Rosbrook B, Anderson CM, et al. The demographics of help-seeking for smoking cessation in California and the role of the California Smokers' Helpline. *Tobacco Control* 1995;4(Suppl I): S9–S15.