

TABLE 2**Laboratory confirmed *Shigella* isolates reported to the CDC by Serotype in 2004**

Rank	Serotype	Reported	Percent
1	<i>S. sonnei</i>	6433	68.9
2	<i>S. flexneri</i> unspecified	814	8.7
3	<i>S. flexneri</i> 2 unspecified	185	2.0
4	<i>S. flexneri</i> 3 unspecified	112	1.2
5	<i>S. boydii</i> unspecified	111	1.2
6	<i>S. flexneri</i> 1 unspecified	98	1.1
7	<i>S. flexneri</i> 2a	89	1.0
8	<i>S. flexneri</i> 4 unspecified	69	0.7
9	<i>S. flexneri</i> 3a	53	0.6
10	<i>S. flexneri</i> 4a	53	0.6
11	<i>S. flexneri</i> 6	42	0.5
12	<i>S. flexneri</i> 1b	27	0.3
13	<i>S. flexneri</i> variant y	26	0.3
14	<i>S. dysenteriae</i> unspecified	22	0.2
15	<i>S. boydii</i> 2	20	0.2
16	<i>S. boydii</i> 4	15	0.2
17	<i>S. flexneri</i> 3b	13	0.1
18	<i>S. boydii</i> 1	10	0.1
19	<i>S. flexneri</i> 2b	10	0.1
20	<i>S. flexneri</i> variant x	5	0.1
21	<i>S. dysenteriae</i> 1	4	0.0
22	<i>S. dysenteriae</i> 2	4	0.0
23	<i>S. dysenteriae</i> 3	4	0.0
24	<i>S. boydii</i> 10	3	0.0
25	<i>S. boydii</i> 14	3	0.0
26	<i>S. boydii</i> 5	3	0.0
27	<i>S. flexneri</i> 5 unspecified	3	0.0
28	<i>S. boydii</i> 8	2	0.0
29	<i>S. dysenteriae</i> 12	2	0.0
30	<i>S. flexneri</i> 5a	2	0.0
31	<i>S. boydii</i> 12	1	0.0
32	<i>S. boydii</i> 20	1	0.0
33	<i>S. dysenteriae</i> 4	1	0.0
34	<i>S. flexneri</i> 1a	1	0.0
35	<i>S. flexneri</i> 88-893	1	0.0
	Sub Total	8242	88.2
	Unknown	1101	11.8
	Sub Total	1101	11.8
	Total	9343	100.0