LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|-----------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | United States | | | |
| | All causes | 2,443,387 | 100.0 | 847. |
| | Diseases of heart (I00-I09,I11,I13,I20-I51) | 696,947 | 28.5 | 241. |
| | Malignant neoplasms (C00-C97) | 557,271 | 22.8 | 193 |
| | Cerebrovascular diseases (I60-I69) | 162,672 | 6.7 | 56 |
| | Chronic lower respiratory diseases (J40-J47) | 124,816 | 5.1 | 43 |
| | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 106,742 | 4.4 | 37 |
| | Diabetes mellitus (E10-E14) | 73,249 | 3.0 | 25 |
| | Influenza and pneumonia (J10-J18) | 65,681 | 2.7 | 22 |
| | Alzheimer's disease (G30) | 58,866 | 2.4 | 20 |
| | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 40,974 | 1.7 | 14 |
| 0 | Septicemia (A40-A41) | 33,865 | 1.4 | 11 |
| 1 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 31,655 | 1.3 | 11 |
| 2 | Chronic liver disease and cirrhosis (K70,K73-K74) | 27,257 | 1.1 | 9 |
| 3 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 20,261 | 0.8 | 7 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 17,638 | 0.7 | 6 |
| 5 | Pneumonitis due to solids and liquids (J69) | 17,593 | 0.7 | 6 |
| | All other causes (Residual) | 407,900 | 16.7 | 141 |
| | Alabama | | | |
| | All causes | 46,069 | 100.0 | 1,026 |
| | Diseases of heart (I00-I09,I11,I13,I20-I51) | 13,197 | 28.6 | 294 |
| | Malignant neoplasms (C00-C97) | 9,698 | 21.1 | 216 |
| | Cerebrovascular diseases (I60-I69) | 3,201 | 6.9 | 71 |
| | Chronic lower respiratory diseases (J40-J47) | 2,328 | 5.1 | 51 |
| | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,228 | 4.8 | 49 |
| | Diabetes mellitus (E10-E14) | 1,486 | 3.2 | 33 |
| | Influenza and pneumonia (J10-J18) | 1,218 | 2.6 | 27 |
| | Alzheimer's disease (G30) | 1,189 | 2.6 | 26 |
| | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,032 | 2.2 | 23 |
|) | Septicemia (A40-A41) | 765 | 1.7 | 17 |
| | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 514 | 1.1 | 11 |
| 2 | Chronic liver disease and cirrhosis (K70,K73-K74) | 425 | 0.9 | Ç |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 416 | 0.9 | Ç |
| 1 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 343 | 0.7 | 7 |
| 5 | Pneumonitis due to solids and liquids (J69) | 323 | 0.7 | 7 |
| | All other causes (Residual) | 7,706 | 16.7 | 171 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|-------------------------------------------------------------------------------|--------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Alaska | | | |
| | All causes | 3,030 | 100.0 | 470.7 |
| 1 | Malignant neoplasms (C00-C97) | 715 | 23.6 | 111.1 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 567 | 18.7 | 88.1 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 346 | 11.4 | 53.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 158 | 5.2 | 24.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 142 | 4.7 | 22.1 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 132 | 4.4 | 20.5 |
| 7 | Diabetes mellitus (E10-E14) | 86 | 2.8 | 13.4 |
| 8 | Alzheimer's disease (G30) | 61 | 2.0 | 9.5 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 55 | 1.8 | 8.5 |
| 10 | Influenza and pneumonia (J10-J18) | 51 | 1.7 | 7.9 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 40 | 1.3 | 6.2 |
| 12 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q95 | 29 | 1.0 | 4.5 |
| 13 | Atherosclerosis (I70) | 26 | 0.9 | 4.0 |
| 14 | Septicemia (A40-A41) | 25 | 0.8 | 3.9 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 21 | 0.7 | 3.3 |
| | All other causes (Residual) | 576 | 19.0 | 89.5 |
| | Arizona | | | |
| | All causes | 42,816 | 100.0 | 784.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,852 | 25.3 | 198.9 |
| 2 | Malignant neoplasms (C00-C97) | 9,359 | 21.9 | 171.5 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,577 | 6.0 | 47.2 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,575 | 6.0 | 47.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 2,535 | 5.9 | 46.5 |
| 6 | Alzheimer's disease (G30) | 1,433 | 3.3 | 26.3 |
| 7 | Influenza and pneumonia (J10-J18) | 1,319 | 3.1 | 24.2 |
| 8 | Diabetes mellitus (E10-E14) | 1,231 | 2.9 | 22.6 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 886 | 2.1 | 16.2 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 671 | 1.6 | 12.3 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 622 | 1.5 | 11.4 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 504 | 1.2 | 9.2 |
| 13 | Septicemia (A40-A41) | 461 | 1.1 | 8.4 |
| 14 | Parkinson's disease (G20-G21) | 398 | 0.9 | 7.3 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 303 | 0.7 | 5.6 |
| | All other causes (Residual) | 7,090 | 16.6 | 129.9 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|---------|------------------|---------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Arkansas | | | |
| | All causes | 28,513 | 100.0 | 1,052.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8,330 | 29.2 | 307.4 |
| 2 | Malignant neoplasms (C00-C97) | 6,282 | 22.0 | 231.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,232 | 7.8 | 82.4 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,441 | 5.1 | 53.2 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,311 | 4.6 | 48.4 |
| 6 | Diabetes mellitus (E10-E14) | 793 | 2.8 | 29.3 |
| 7 | Influenza and pneumonia (J10-J18) | 776 | 2.7 | 28.6 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 601 | 2.1 | 22.2 |
| 9 | Alzheimer's disease (G30) | 551 | 1.9 | 20.3 |
| 10 | Septicemia (A40-A41) | 502 | 1.8 | 18.5 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 377 | 1.3 | 13.9 |
| 12 | Pneumonitis due to solids and liquids (J69) | 224 | 0.8 | 8.3 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 222 | 0.8 | 8.2 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 194 | 0.7 | 7.2 |
| 15 | Aortic aneurysm and dissection (I71) | 179 | 0.6 | 6.6 |
| | All other causes (Residual) | 4,498 | 15.8 | 166.0 |
| | California | | | |
| | All causes | 234,565 | 100.0 | 668.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 68,797 | 29.3 | 195.9 |
| 2 | Malignant neoplasms (C00-C97) | 54,143 | 23.1 | 154.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 17,626 | 7.5 | 50.2 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 12,684 | 5.4 | 36.1 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 10,107 | 4.3 | 28.8 |
| 6 | Influenza and pneumonia (J10-J18) | 8,128 | 3.5 | 23.1 |
| 7 | Diabetes mellitus (E10-E14) | 6,807 | 2.9 | 19.4 |
| 8 | Alzheimer's disease (G30) | 5,421 | 2.3 | 15.4 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 3,747 | 1.6 | 10.7 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 3,228 | 1.4 | 9.2 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 2,485 | 1.1 | 7.1 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 2,337 | 1.0 | 6.7 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,164 | 0.9 | 6.2 |
| 14 | Atherosclerosis (I70) | 1,588 | 0.7 | 4.5 |
| 15 | Parkinson's disease (G20-G21) | 1,582 | 0.7 | 4.5 |
| | All other causes (Residual) | 33,721 | 14.4 | 96.0 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Colorado | | | |
| | All causes | 29,210 | 100.0 | 648.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,425 | 22.0 | 142.6 |
| 2 | Malignant neoplasms (C00-C97) | 6,384 | 21.9 | 141.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,915 | 6.6 | 42.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,848 | 6.3 | 41.0 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,812 | 6.2 | 40.2 |
| 6 | Alzheimer's disease (G30) | 954 | 3.3 | 21.2 |
| 7 | Influenza and pneumonia (J10-J18) | 752 | 2.6 | 16.7 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 727 | 2.5 | 16.1 |
| 9 | Diabetes mellitus (E10-E14) | 659 | 2.3 | 14.6 |
| 10 | Atherosclerosis (I70) | 503 | 1.7 | 11.2 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 417 | 1.4 | 9.3 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 415 | 1.4 | 9.2 |
| 13 | Septicemia (A40-A41) | 279 | 1.0 | 6.2 |
| 14 | Parkinson's disease (G20-G21) | 251 | 0.9 | 5.6 |
| 15 | Pneumonitis due to solids and liquids (J69) | 250 | 0.9 | 5.5 |
| | All other causes (Residual) | 5,619 | 19.2 | 124.7 |
| | Connecticut | | | |
| | All causes | 30,122 | 100.0 | 870.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8,815 | 29.3 | 254.7 |
| 2 | Malignant neoplasms (C00-C97) | 7,163 | 23.8 | 207.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,861 | 6.2 | 53.8 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,453 | 4.8 | 42.0 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,182 | 3.9 | 34.2 |
| 6 | Influenza and pneumonia (J10-J18) | 888 | 2.9 | 25.7 |
| 7 | Diabetes mellitus (E10-E14) | 675 | 2.2 | 19.5 |
| 8 | Alzheimer's disease (G30) | 570 | 1.9 | 16.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 554 | 1.8 | 16.0 |
| 10 | Septicemia (A40-A41) | 527 | 1.7 | 15.2 |
| 11 | Pneumonitis due to solids and liquids (J69) | 374 | 1.2 | 10.8 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 318 | 1.1 | 9.2 |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 260 | 0.9 | 7.5 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 256 | 0.8 | 7.4 |
| 15 | Parkinson's disease (G20-G21) | 200 | 0.7 | 5.8 |
| | All other causes (Residual) | 5,026 | 16.7 | 145.2 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|--------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Delaware | | | |
| | All causes | 6,861 | 100.0 | 849. |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,918 | 28.0 | 237. |
| 2 | Malignant neoplasms (C00-C97) | 1,621 | 23.6 | 200. |
| 3 | Cerebrovascular diseases (I60-I69) | 405 | 5.9 | 50. |
| ļ | Chronic lower respiratory diseases (J40-J47) | 350 | 5.1 | 43. |
| , | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 292 | 4.3 | 36. |
| 3 | Diabetes mellitus (E10-E14) | 215 | 3.1 | 26. |
| , | Influenza and pneumonia (J10-J18) | 168 | 2.4 | 20 |
| 3 | Septicemia (A40-A41) | 133 | 1.9 | 16 |
|) | Alzheimer's disease (G30) | 128 | 1.9 | 15 |
| 0 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 117 | 1.7 | 14 |
| 1 | Chronic liver disease and cirrhosis (K70,K73-K74) | 88 | 1.3 | 10 |
| 2 | Pneumonitis due to solids and liquids (J69) | 74 | 1.1 | 9 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 74 | 1.1 | 9 |
| 4 | Human immunodeficiency virus (HIV) disease (B20-B24) | 70 | 1.0 | 8 |
| 5 | Certain conditions originating in the perinatal period (P00-P96) | 62 | 0.9 | 7 |
| | All other causes (Residual) | 1,146 | 16.7 | 141 |
| | District of Columbia | | | |
| | All causes | 5,851 | 100.0 | 1,024. |
| | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,666 | 28.5 | 291 |
|) | Malignant neoplasms (C00-C97) | 1,298 | 22.2 | 227 |
| , | Cerebrovascular diseases (I60-I69) | 279 | 4.8 | 48 |
| ļ | Human immunodeficiency virus (HIV) disease (B20-B24) | 233 | 4.0 | 40 |
| ; | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 229 | 3.9 | 40 |
| i | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 200 | 3.4 | 35 |
| • | Diabetes mellitus (E10-E14) | 191 | 3.3 | 33 |
| } | Chronic lower respiratory diseases (J40-J47) | 133 | 2.3 | 23 |
|) | Septicemia (A40-A41) | 130 | 2.2 | 22 |
| 0 | Alzheimer's disease (G30) | 107 | 1.8 | 18 |
| 1 | Chronic liver disease and cirrhosis (K70,K73-K74) | 88 | 1.5 | 15 |
| 2 | Influenza and pneumonia (J10-J18) | 81 | 1.4 | 14 |
| 3 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 70 | 1.2 | 12 |
| 4 | Certain conditions originating in the perinatal period (P00-P96) | 44 | 0.8 | 7 |
| 5 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 39 | 0.7 | 6 |
| | All other causes (Residual) | 1,063 | 18.2 | 186 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|---------|------------------|--------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Florida | | | |
| | All causes | 167,814 | 100.0 | 1,004. |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 49,235 | 29.3 | 294. |
| 2 | Malignant neoplasms (C00-C97) | 39,140 | 23.3 | 234. |
| } | Cerebrovascular diseases (I60-I69) | 10,269 | 6.1 | 61. |
| ļ | Chronic lower respiratory diseases (J40-J47) | 9,062 | 5.4 | 54. |
| , | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 7,396 | 4.4 | 44 |
| ; | Diabetes mellitus (E10-E14) | 4,583 | 2.7 | 27. |
| • | Alzheimer's disease (G30) | 4,052 | 2.4 | 24 |
| } | Influenza and pneumonia (J10-J18) | 3,290 | 2.0 | 19 |
|) | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,338 | 1.4 | 14 |
| 0 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,201 | 1.3 | 13 |
| 1 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,151 | 1.3 | 12 |
| 2 | Septicemia (A40-A41) | 1,858 | 1.1 | 11 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,719 | 1.0 | 10 |
| 4 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1,448 | 0.9 | 8 |
| 5 | Parkinson's disease (G20-G21) | 1,284 | 0.8 | 7 |
| | All other causes (Residual) | 27,788 | 16.6 | 166 |
| | Georgia | | | |
| | All causes | 65,449 | 100.0 | 764 |
| | Diseases of heart (I00-I09,I11,I13,I20-I51) | 17,529 | 26.8 | 204 |
| | Malignant neoplasms (C00-C97) | 13,975 | 21.4 | 163 |
| } | Cerebrovascular diseases (I60-I69) | 4,261 | 6.5 | 49 |
| | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,333 | 5.1 | 38 |
| | Chronic lower respiratory diseases (J40-J47) | 3,163 | 4.8 | 36 |
| | Influenza and pneumonia (J10-J18) | 1,791 | 2.7 | 20 |
| • | Diabetes mellitus (E10-E14) | 1,576 | 2.4 | 18 |
| | Alzheimer's disease (G30) | 1,525 | 2.3 | 17 |
| | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,335 | 2.0 | 15 |
| 0 | Septicemia (A40-A41) | 1,293 | 2.0 | 15 |
| 1 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 909 | 1.4 | 10 |
| 2 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 712 | 1.1 | 8 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 708 | 1.1 | 8 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 696 | 1.1 | 8 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 672 | 1.0 | 7 |
| | All other causes (Residual) | 11,971 | 18.3 | 139 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|-------------------------------------------------------------------------------|--------|------------------|------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Hawaii | | | |
| | All causes | 8,801 | 100.0 | 707. |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,512 | 28.5 | 201. |
| 2 | Malignant neoplasms (C00-C97) | 1,945 | 22.1 | 156. |
| } | Cerebrovascular diseases (I60-I69) | 812 | 9.2 | 65. |
| ļ | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 393 | 4.5 | 31. |
| , | Chronic lower respiratory diseases (J40-J47) | 265 | 3.0 | 21 |
| ; | Influenza and pneumonia (J10-J18) | 246 | 2.8 | 19. |
| , | Diabetes mellitus (E10-E14) | 204 | 2.3 | 16 |
| } | Septicemia (A40-A41) | 145 | 1.6 | 11. |
|) | Alzheimer's disease (G30) | 141 | 1.6 | 11. |
| 0 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 136 | 1.5 | 10 |
| 1 | Pneumonitis due to solids and liquids (J69) | 129 | 1.5 | 10 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 120 | 1.4 | 9 |
| 3 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 89 | 1.0 | 7 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 79 | 0.9 | 6 |
| 5 | Parkinson's disease (G20-G21) | 74 | 0.8 | 5 |
| | All other causes (Residual) | 1,511 | 17.2 | 121. |
| | Idaho | | | |
| | All causes | 9,923 | 100.0 | 739. |
| | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,532 | 25.5 | 188 |
| <u>)</u> | Malignant neoplasms (C00-C97) | 2,138 | 21.5 | 159 |
| } | Cerebrovascular diseases (I60-I69) | 736 | 7.4 | 54 |
| Ļ | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 611 | 6.2 | 45 |
| | Chronic lower respiratory diseases (J40-J47) | 595 | 6.0 | 44 |
| | Diabetes mellitus (E10-E14) | 322 | 3.2 | 24 |
| • | Alzheimer's disease (G30) | 318 | 3.2 | 23 |
| | Influenza and pneumonia (J10-J18) | 265 | 2.7 | 19 |
| | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 202 | 2.0 | 15 |
| 0 | Chronic liver disease and cirrhosis (K70,K73-K74) | 105 | 1.1 | 7 |
| 1 | Parkinson's disease (G20-G21) | 103 | 1.0 | 7 |
| 2 | Pneumonitis due to solids and liquids (J69) | 89 | 0.9 | 6 |
| 3 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 86 | 0.9 | 6 |
| 4 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 83 | 0.8 | 6 |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99 | 64 | 0.6 | 4 |
| | All other causes (Residual) | 1,674 | 16.9 | 124 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|---------|------------------|------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Illinois | | | |
| | All causes | 106,667 | 100.0 | 846. |
| | Diseases of heart (I00-I09,I11,I13,I20-I51) | 30,821 | 28.9 | 244. |
| | Malignant neoplasms (C00-C97) | 24,737 | 23.2 | 196 |
| | Cerebrovascular diseases (I60-I69) | 7,183 | 6.7 | 57 |
| | Chronic lower respiratory diseases (J40-J47) | 4,827 | 4.5 | 38 |
| | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,222 | 4.0 | 33 |
| | Diabetes mellitus (E10-E14) | 3,011 | 2.8 | 23 |
| | Influenza and pneumonia (J10-J18) | 2,940 | 2.8 | 23 |
| | Alzheimer's disease (G30) | 2,398 | 2.2 | 19 |
| | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,328 | 2.2 | 18 |
| 0 | Septicemia (A40-A41) | 2,057 | 1.9 | 16 |
| 1 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,145 | 1.1 | 9 |
| 2 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,068 | 1.0 | 8 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,016 | 1.0 | 8 |
| 4 | Pneumonitis due to solids and liquids (J69) | 908 | 0.9 | 7 |
| 5 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 856 | 0.8 | 6 |
| | All other causes (Residual) | 17,150 | 16.1 | 136 |
| | Indiana | | | |
| | All causes | 55,396 | 100.0 | 899 |
| | Diseases of heart (I00-I09,I11,I13,I20-I51) | 15,321 | 27.7 | 248 |
| | Malignant neoplasms (C00-C97) | 12,865 | 23.2 | 208 |
| | Cerebrovascular diseases (I60-I69) | 3,717 | 6.7 | 60 |
| | Chronic lower respiratory diseases (J40-J47) | 3,138 | 5.7 | 50 |
| | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,148 | 3.9 | 34 |
| | Diabetes mellitus (E10-E14) | 1,688 | 3.0 | 27 |
| | Alzheimer's disease (G30) | 1,475 | 2.7 | 23 |
| | Influenza and pneumonia (J10-J18) | 1,360 | 2.5 | 22 |
| | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,222 | 2.2 | 19 |
|) | Septicemia (A40-A41) | 831 | 1.5 | 13 |
| l | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 743 | 1.3 | 12 |
| 2 | Atherosclerosis (I70) | 526 | 0.9 | ; |
| 3 | Chronic liver disease and cirrhosis (K70,K73-K74) | 514 | 0.9 | ; |
| 4 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 424 | 0.8 | (|
| 5 | Pneumonitis due to solids and liquids (J69) | 414 | 0.7 | 6 |
| | All other causes (Residual) | 9,010 | 16.3 | 146 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | lowa | | | |
| | All causes | 27,978 | 100.0 | 952.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8,181 | 29.2 | 278.6 |
| 2 | Malignant neoplasms (C00-C97) | 6,473 | 23.1 | 220.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,226 | 8.0 | 75.8 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,580 | 5.6 | 53.8 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,093 | 3.9 | 37.2 |
| 6 | Influenza and pneumonia (J10-J18) | 942 | 3.4 | 32.1 |
| 7 | Alzheimer's disease (G30) | 899 | 3.2 | 30.6 |
| 8 | Diabetes mellitus (E10-E14) | 734 | 2.6 | 25.0 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 314 | 1.1 | 10.7 |
| 10 | Atherosclerosis (I70) | 261 | 0.9 | 8.9 |
| 10 | Pneumonitis due to solids and liquids (J69) | 261 | 0.9 | 8.9 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 261 | 0.9 | 8.9 |
| 13 | Parkinson's disease (G20-G21) | 250 | 0.9 | 8.5 |
| 14 | Septicemia (A40-A41) | 231 | 8.0 | 7.9 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 220 | 0.8 | 7.5 |
| | All other causes (Residual) | 4,052 | 14.5 | 138.0 |
| | Kansas | | | |
| | All causes | 25,021 | 100.0 | 921.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,680 | 26.7 | 246.0 |
| 2 | Malignant neoplasms (C00-C97) | 5,362 | 21.4 | 197.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,845 | 7.4 | 67.9 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,367 | 5.5 | 50.3 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,139 | 4.6 | 41.9 |
| 6 | Diabetes mellitus (E10-E14) | 765 | 3.1 | 28.2 |
| 7 | Alzheimer's disease (G30) | 755 | 3.0 | 27.8 |
| 8 | Influenza and pneumonia (J10-J18) | 698 | 2.8 | 25.7 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 517 | 2.1 | 19.0 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 345 | 1.4 | 12.7 |
| 11 | Atherosclerosis (I70) | 343 | 1.4 | 12.6 |
| 12 | Septicemia (A40-A41) | 297 | 1.2 | 10.9 |
| 13 | Parkinson's disease (G20-G21) | 218 | 0.9 | 8.0 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 195 | 0.8 | 7.2 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 187 | 0.7 | 6.9 |
| | All other causes (Residual) | 4,308 | 17.2 | 158.6 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Kentucky | | | |
| | All causes | 40,697 | 100.0 | 994.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,696 | 28.7 | 285.8 |
| 2 | Malignant neoplasms (C00-C97) | 9,438 | 23.2 | 230.6 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,554 | 6.3 | 62.4 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,401 | 5.9 | 58.7 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,090 | 5.1 | 51.1 |
| 6 | Diabetes mellitus (E10-E14) | 1,265 | 3.1 | 30.9 |
| 7 | Influenza and pneumonia (J10-J18) | 1,236 | 3.0 | 30.2 |
| 8 | Alzheimer's disease (G30) | 1,013 | 2.5 | 24.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 813 | 2.0 | 19.9 |
| 10 | Septicemia (A40-A41) | 606 | 1.5 | 14.8 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 540 | 1.3 | 13.2 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 377 | 0.9 | 9.2 |
| 13 | Pneumonitis due to solids and liquids (J69) | 320 | 0.8 | 7.8 |
| 14 | Parkinson's disease (G20-G21) | 224 | 0.6 | 5.5 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 221 | 0.5 | 5.4 |
| | All other causes (Residual) | 5,903 | 14.5 | 144.2 |
| | Louisiana | | | |
| | All causes | 41,984 | 100.0 | 936.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,185 | 26.6 | 249.5 |
| 2 | Malignant neoplasms (C00-C97) | 9,441 | 22.5 | 210.6 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,595 | 6.2 | 57.9 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,115 | 5.0 | 47.2 |
| 5 | Diabetes mellitus (E10-E14) | 1,774 | 4.2 | 39.6 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 1,696 | 4.0 | 37.8 |
| 7 | Alzheimer's disease (G30) | 1,111 | 2.6 | 24.8 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 983 | 2.3 | 21.9 |
| 9 | Influenza and pneumonia (J10-J18) | 956 | 2.3 | 21.3 |
| 10 | Septicemia (A40-A41) | 820 | 2.0 | 18.3 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 607 | 1.4 | 13.5 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 499 | 1.2 | 11.1 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 365 | 0.9 | 8.1 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 364 | 0.9 | 8.1 |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 338 | 0.8 | 7.5 |
| | All other causes (Residual) | 7,135 | 17.0 | 159.2 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Maine | | | |
| | All causes | 12,694 | 100.0 | 980.6 |
| 1 | Malignant neoplasms (C00-C97) | 3,206 | 25.3 | 247.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,170 | 25.0 | 244.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 823 | 6.5 | 63.6 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 791 | 6.2 | 61.1 |
| 5 | Alzheimer's disease (G30) | 513 | 4.0 | 39.6 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 511 | 4.0 | 39.5 |
| 7 | Diabetes mellitus (E10-E14) | 404 | 3.2 | 31.2 |
| 8 | Influenza and pneumonia (J10-J18) | 317 | 2.5 | 24.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 232 | 1.8 | 17.9 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 166 | 1.3 | 12.8 |
| 11 | Septicemia (A40-A41) | 128 | 1.0 | 9.9 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 116 | 0.9 | 9.0 |
| 13 | Parkinson's disease (G20-G21) | 103 | 0.8 | 8.0 |
| 14 | Aortic aneurysm and dissection (I71) | 100 | 0.8 | 7.7 |
| 15 | Pneumonitis due to solids and liquids (J69) | 88 | 0.7 | 6.8 |
| | All other causes (Residual) | 2,026 | 16.0 | 156.5 |
| | Maryland | | | |
| | All causes | 43,970 | 100.0 | 805.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 12,008 | 27.3 | 220.0 |
| 2 | Malignant neoplasms (C00-C97) | 10,395 | 23.6 | 190.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,811 | 6.4 | 51.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,944 | 4.4 | 35.6 |
| 5 | Diabetes mellitus (E10-E14) | 1,519 | 3.5 | 27.8 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,332 | 3.0 | 24.4 |
| 7 | Influenza and pneumonia (J10-J18) | 1,121 | 2.5 | 20.5 |
| 8 | Septicemia (A40-A41) | 1,023 | 2.3 | 18.7 |
| 9 | Alzheimer's disease (G30) | 866 | 2.0 | 15.9 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 630 | 1.4 | 11.5 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 610 | 1.4 | 11.2 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 540 | 1.2 | 9.9 |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 477 | 1.1 | 8.7 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 443 | 1.0 | 8.1 |
| 15 | Pneumonitis due to solids and liquids (J69) | 412 | 0.9 | 7.5 |
| | All other causes (Residual) | 7,839 | 17.8 | 143.6 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Massachusetts | | | |
| | All causes | 56,928 | 100.0 | 885. |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 14,736 | 25.9 | 229. |
| <u>-</u> | Malignant neoplasms (C00-C97) | 13,914 | 24.4 | 216 |
| } | Cerebrovascular diseases (I60-I69) | 3,559 | 6.3 | 55 |
| ļ | Chronic lower respiratory diseases (J40-J47) | 2,745 | 4.8 | 42 |
| ; | Influenza and pneumonia (J10-J18) | 2,087 | 3.7 | 32 |
| 6 | Alzheimer's disease (G30) | 1,570 | 2.8 | 24 |
| • | Diabetes mellitus (E10-E14) | 1,423 | 2.5 | 22 |
| , | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,413 | 2.5 | 22 |
|) | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,297 | 2.3 | 20 |
| 0 | Septicemia (A40-A41) | 1,019 | 1.8 | 15 |
| 1 | Chronic liver disease and cirrhosis (K70,K73-K74) | 602 | 1.1 | 9 |
| 2 | Pneumonitis due to solids and liquids (J69) | 583 | 1.0 | 9 |
| 3 | Parkinson's disease (G20-G21) | 460 | 8.0 | 7 |
| 4 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 450 | 0.8 | 7 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 436 | 0.8 | 6 |
| | All other causes (Residual) | 10,634 | 18.7 | 165 |
| | Michigan | | | |
| | All causes | 87,795 | 100.0 | 873 |
| | Diseases of heart (I00-I09,I11,I13,I20-I51) | 26,659 | 30.4 | 265 |
| | Malignant neoplasms (C00-C97) | 19,985 | 22.8 | 198 |
| , | Cerebrovascular diseases (I60-I69) | 5,814 | 6.6 | 57 |
| | Chronic lower respiratory diseases (J40-J47) | 4,431 | 5.0 | 44 |
| | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,285 | 3.7 | 32 |
| i | Diabetes mellitus (E10-E14) | 2,785 | 3.2 | 27 |
| | Influenza and pneumonia (J10-J18) | 2,029 | 2.3 | 20 |
| | Alzheimer's disease (G30) | 1,958 | 2.2 | 19 |
| | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,618 | 1.8 | 16 |
| 0 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,106 | 1.3 | 11 |
| 1 | Chronic liver disease and cirrhosis (K70,K73-K74) | 993 | 1.1 | 9 |
| 2 | Septicemia (A40-A41) | 924 | 1.1 | ç |
| 3 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 723 | 8.0 | 7 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 696 | 8.0 | 6 |
| 5 | Parkinson's disease (G20-G21) | 637 | 0.7 | 6 |
| | All other causes (Residual) | 14,152 | 16.1 | 140 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------------|--------|------------------|---------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Minnesota | | | |
| | All causes | 38,510 | 100.0 | 767.2 |
| 1 | Malignant neoplasms (C00-C97) | 9,210 | 23.9 | 183.5 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8,602 | 22.3 | 171.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,706 | 7.0 | 53.9 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,971 | 5.1 | 39.3 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,928 | 5.0 | 38.4 |
| 6 | Diabetes mellitus (E10-E14) | 1,317 | 3.4 | 26.2 |
| 7 | Alzheimer's disease (G30) | 1,192 | 3.1 | 23.7 |
| 8 | Influenza and pneumonia (J10-J18) | 900 | 2.3 | 17.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 649 | 1.7 | 12.9 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 497 | 1.3 | 9.9 |
| 11 | Parkinson's disease (G20-G21) | 440 | 1.1 | 8.8 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 434 | 1.1 | 8.6 |
| 13 | Aortic aneurysm and dissection (I71) | 330 | 0.9 | 6.6 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 320 | 0.8 | 6.4 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior | 282 | 0.7 | 5.6 |
| | All other causes (Residual) | 7,732 | 20.1 | 154.0 |
| | Mississippi | | | |
| | All causes | 28,853 | 100.0 | 1,004.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 9,061 | 31.4 | 315.5 |
| 2 | Malignant neoplasms (C00-C97) | 6,069 | 21.0 | 211.3 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,926 | 6.7 | 67.1 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,642 | 5.7 | 57.2 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1,378 | 4.8 | 48.0 |
| 6 | Influenza and pneumonia (J10-J18) | 801 | 2.8 | 27.9 |
| 7 | Diabetes mellitus (E10-E14) | 671 | 2.3 | 23.4 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 580 | 2.0 | 20.2 |
| 9 | Alzheimer's disease (G30) | 574 | 2.0 | 20.0 |
| 10 | Septicemia (A40-A41) | 521 | 1.8 | 18.1 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 343 | 1.2 | 11.9 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 313 | 1.1 | 10.9 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 305 | 1.1 | 10.6 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 229 | 0.8 | 8.0 |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 222 | 0.8 | 7.7 |
| | All other causes (Residual) | 4,218 | 14.6 | 146.9 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Missouri | | | |
| | All causes | 55,940 | 100.0 | 986.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 16,708 | 29.9 | 294.5 |
| 2 | Malignant neoplasms (C00-C97) | 12,322 | 22.0 | 217.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,885 | 6.9 | 68.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,867 | 5.1 | 50.5 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,641 | 4.7 | 46.6 |
| 6 | Diabetes mellitus (E10-E14) | 1,625 | 2.9 | 28.6 |
| 7 | Influenza and pneumonia (J10-J18) | 1,619 | 2.9 | 28.5 |
| 8 | Alzheimer's disease (G30) | 1,187 | 2.1 | 20.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,076 | 1.9 | 19.0 |
| 10 | Septicemia (A40-A41) | 791 | 1.4 | 13.9 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 693 | 1.2 | 12.2 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 458 | 0.8 | 8.1 |
| 13 | Pneumonitis due to solids and liquids (J69) | 437 | 0.8 | 7.7 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 433 | 0.8 | 7.6 |
| 15 | Parkinson's disease (G20-G21) | 371 | 0.7 | 6.5 |
| | All other causes (Residual) | 8,827 | 15.8 | 155.6 |
| | Montana | | | |
| | All causes | 8,506 | 100.0 | 935.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,944 | 22.9 | 213.8 |
| 2 | Malignant neoplasms (C00-C97) | 1,911 | 22.5 | 210.1 |
| 3 | Cerebrovascular diseases (I60-I69) | 639 | 7.5 | 70.3 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 576 | 6.8 | 63.3 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 524 | 6.2 | 57.6 |
| 6 | Alzheimer's disease (G30) | 285 | 3.4 | 31.3 |
| 7 | Influenza and pneumonia (J10-J18) | 255 | 3.0 | 28.0 |
| 8 | Diabetes mellitus (E10-E14) | 210 | 2.5 | 23.1 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 184 | 2.2 | 20.2 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 127 | 1.5 | 14.0 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 105 | 1.2 | 11.5 |
| 12 | Pneumonitis due to solids and liquids (J69) | 79 | 0.9 | 8.7 |
| 13 | Septicemia (A40-A41) | 66 | 0.8 | 7.3 |
| 14 | Parkinson's disease (G20-G21) | 61 | 0.7 | 6.7 |
| 14 | Aortic aneurysm and dissection (I71) | 61 | 0.7 | 6.7 |
| | All other causes (Residual) | 1,479 | 17.4 | 162.6 |
| | | | | |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Nebraska | | | |
| | All causes | 15,738 | 100.0 | 910.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,242 | 27.0 | 245.3 |
| 2 | Malignant neoplasms (C00-C97) | 3,433 | 21.8 | 198.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,103 | 7.0 | 63.8 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 934 | 5.9 | 54.0 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 762 | 4.8 | 44.1 |
| 6 | Alzheimer's disease (G30) | 460 | 2.9 | 26.6 |
| 7 | Influenza and pneumonia (J10-J18) | 418 | 2.7 | 24.2 |
| 8 | Diabetes mellitus (E10-E14) | 393 | 2.5 | 22.7 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 277 | 1.8 | 16.0 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 201 | 1.3 | 11.6 |
| 11 | Septicemia (A40-A41) | 147 | 0.9 | 8.5 |
| 12 | Atherosclerosis (I70) | 143 | 0.9 | 8.3 |
| 13 | Parkinson's disease (G20-G21) | 142 | 0.9 | 8.2 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 142 | 0.9 | 8.2 |
| 15 | Pneumonitis due to solids and liquids (J69) | 140 | 0.9 | 8.1 |
| | All other causes (Residual) | 2,801 | 17.8 | 162.0 |
| | Nevada | | | |
| | All causes | 16,927 | 100.0 | 778.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,421 | 26.1 | 203.4 |
| 2 | Malignant neoplasms (C00-C97) | 3,937 | 23.3 | 181.1 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 1,174 | 6.9 | 54.0 |
| 4 | Cerebrovascular diseases (I60-I69) | 976 | 5.8 | 44.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 860 | 5.1 | 39.6 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 423 | 2.5 | 19.5 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 372 | 2.2 | 17.1 |
| 8 | Influenza and pneumonia (J10-J18) | 368 | 2.2 | 16.9 |
| 9 | Diabetes mellitus (E10-E14) | 343 | 2.0 | 15.8 |
| 10 | Septicemia (A40-A41) | 336 | 2.0 | 15.5 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 268 | 1.6 | 12.3 |
| 12 | Alzheimer's disease (G30) | 253 | 1.5 | 11.6 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 175 | 1.0 | 8.1 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 122 | 0.7 | 5.6 |
| 15 | Atherosclerosis (I70) | 108 | 0.6 | 5.0 |
| | All other causes (Residual) | 2,791 | 16.5 | 128.4 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | New Hampshire | | | |
| | All causes | 9,853 | 100.0 | 772.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,776 | 28.2 | 217.7 |
| 2 | Malignant neoplasms (C00-C97) | 2,529 | 25.7 | 198.3 |
| 3 | Cerebrovascular diseases (I60-I69) | 627 | 6.4 | 49.2 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 577 | 5.9 | 45.3 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 357 | 3.6 | 28.0 |
| 6 | Diabetes mellitus (E10-E14) | 311 | 3.2 | 24.4 |
| 6 | Alzheimer's disease (G30) | 311 | 3.2 | 24.4 |
| 8 | Influenza and pneumonia (J10-J18) | 237 | 2.4 | 18.6 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 141 | 1.4 | 11.1 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 132 | 1.3 | 10.4 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 105 | 1.1 | 8.2 |
| 12 | Septicemia (A40-A41) | 80 | 8.0 | 6.3 |
| 13 | Aortic aneurysm and dissection (I71) | 74 | 8.0 | 5.8 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 69 | 0.7 | 5.4 |
| 15 | Parkinson's disease (G20-G21) | 66 | 0.7 | 5.2 |
| | All other causes (Residual) | 1,461 | 14.8 | 114.6 |
| | New Jersey | | | |
| | All causes | 74,009 | 100.0 | 861.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 22,510 | 30.4 | 262.0 |
| 2 | Malignant neoplasms (C00-C97) | 17,827 | 24.1 | 207.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 4,016 | 5.4 | 46.8 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,885 | 3.9 | 33.6 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,599 | 3.5 | 30.3 |
| 6 | Diabetes mellitus (E10-E14) | 2,532 | 3.4 | 29.5 |
| 7 | Septicemia (A40-A41) | 1,986 | 2.7 | 23.1 |
| 8 | Influenza and pneumonia (J10-J18) | 1,973 | 2.7 | 23.0 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,662 | 2.2 | 19.3 |
| 10 | Alzheimer's disease (G30) | 1,522 | 2.1 | 17.7 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 762 | 1.0 | 8.9 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 730 | 1.0 | 8.5 |
| 13 | Pneumonitis due to solids and liquids (J69) | 628 | 0.8 | 7.3 |
| 14 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 553 | 0.7 | 6.4 |
| 15 | Parkinson's disease (G20-G21) | 524 | 0.7 | 6.1 |
| | All other causes (Residual) | 11,300 | 15.3 | 131.5 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|---------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | New Mexico | | | |
| | All causes | 14,344 | 100.0 | 773.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,360 | 23.4 | 181.1 |
| 2 | Malignant neoplasms (C00-C97) | 3,067 | 21.4 | 165.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,105 | 7.7 | 59.6 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 857 | 6.0 | 46.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 715 | 5.0 | 38.5 |
| 6 | Diabetes mellitus (E10-E14) | 582 | 4.1 | 31.4 |
| 7 | Influenza and pneumonia (J10-J18) | 372 | 2.6 | 20.1 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 349 | 2.4 | 18.8 |
| 9 | Alzheimer's disease (G30) | 325 | 2.3 | 17.5 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 317 | 2.2 | 17.1 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 220 | 1.5 | 11.9 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 161 | 1.1 | 8.7 |
| 13 | Septicemia (A40-A41) | 126 | 0.9 | 6.8 |
| 14 | Parkinson's disease (G20-G21) | 117 | 0.8 | 6.3 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 107 | 0.7 | 5.8 |
| | All other causes (Residual) | 2,564 | 17.9 | 138.2 |
| | New York | | | |
| | All causes | 158,118 | 100.0 | 825.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 56,672 | 35.8 | 295.8 |
| 2 | Malignant neoplasms (C00-C97) | 36,661 | 23.2 | 191.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 7,625 | 4.8 | 39.8 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 6,966 | 4.4 | 36.4 |
| 5 | Influenza and pneumonia (J10-J18) | 5,368 | 3.4 | 28.0 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,663 | 2.9 | 24.3 |
| 7 | Diabetes mellitus (E10-E14) | 3,934 | 2.5 | 20.5 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,465 | 1.6 | 12.9 |
| 9 | Septicemia (A40-A41) | 2,319 | 1.5 | 12.1 |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,980 | 1.3 | 10.3 |
| 11 | Alzheimer's disease (G30) | 1,803 | 1.1 | 9.4 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1,416 | 0.9 | 7.4 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,338 | 0.8 | 7.0 |
| 14 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,228 | 0.8 | 6.4 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 929 | 0.6 | 4.8 |
| | All other causes (Residual) | 22,751 | 14.4 | 118.8 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | North Carolina | | | |
| | All causes | 72,027 | 100.0 | 865.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 18,524 | 25.7 | 222.6 |
| 2 | Malignant neoplasms (C00-C97) | 16,210 | 22.5 | 194.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 5,259 | 7.3 | 63.2 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,700 | 5.1 | 44.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 3,674 | 5.1 | 44.2 |
| 6 | Diabetes mellitus (E10-E14) | 2,205 | 3.1 | 26.5 |
| 7 | Alzheimer's disease (G30) | 1,962 | 2.7 | 23.6 |
| 8 | Influenza and pneumonia (J10-J18) | 1,898 | 2.6 | 22.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,437 | 2.0 | 17.3 |
| 10 | Septicemia (A40-A41) | 1,161 | 1.6 | 14.0 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 986 | 1.4 | 11.9 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 735 | 1.0 | 8.8 |
| 13 | Pneumonitis due to solids and liquids (J69) | 722 | 1.0 | 8.7 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 652 | 0.9 | 7.8 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 644 | 0.9 | 7.7 |
| | All other causes (Residual) | 12,258 | 17.0 | 147.3 |
| | North Dakota | | | |
| | All causes | 5,892 | 100.0 | 929.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,623 | 27.5 | 255.9 |
| 2 | Malignant neoplasms (C00-C97) | 1,293 | 21.9 | 203.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 469 | 8.0 | 74.0 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 322 | 5.5 | 50.8 |
| 5 | Alzheimer's disease (G30) | 294 | 5.0 | 46.4 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 246 | 4.2 | 38.8 |
| 7 | Diabetes mellitus (E10-E14) | 214 | 3.6 | 33.7 |
| 8 | Influenza and pneumonia (J10-J18) | 162 | 2.7 | 25.5 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 91 | 1.5 | 14.4 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 63 | 1.1 | 9.9 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 57 | 1.0 | 9.0 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 53 | 0.9 | 8.4 |
| 13 | Parkinson's disease (G20-G21) | 50 | 0.8 | 7.9 |
| 13 | Pneumonitis due to solids and liquids (J69) | 50 | 0.8 | 7.9 |
| 15 | Septicemia (A40-A41) | 44 | 0.7 | 6.9 |
| | All other causes (Residual) | 861 | 14.6 | 135.8 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|---------|------------------|---------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Ohio | | | |
| | All causes | 109,766 | 100.0 | 961.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 31,388 | 28.6 | 274.8 |
| 2 | Malignant neoplasms (C00-C97) | 25,173 | 22.9 | 220.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 7,252 | 6.6 | 63.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 6,063 | 5.5 | 53.1 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,146 | 3.8 | 36.3 |
| 6 | Diabetes mellitus (E10-E14) | 3,846 | 3.5 | 33.7 |
| 7 | Alzheimer's disease (G30) | 2,599 | 2.4 | 22.8 |
| 8 | Influenza and pneumonia (J10-J18) | 2,487 | 2.3 | 21.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,027 | 1.8 | 17.7 |
| 10 | Septicemia (A40-A41) | 1,627 | 1.5 | 14.2 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,287 | 1.2 | 11.3 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,047 | 1.0 | 9.2 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 944 | 0.9 | 8.3 |
| 14 | Parkinson's disease (G20-G21) | 802 | 0.7 | 7.0 |
| 15 | Aortic aneurysm and dissection (I71) | 697 | 0.6 | 6.1 |
| | All other causes (Residual) | 18,381 | 16.7 | 160.9 |
| | Oklahoma | | | |
| | All causes | 35,502 | 100.0 | 1,016.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,230 | 31.6 | 321.4 |
| 2 | Malignant neoplasms (C00-C97) | 7,474 | 21.1 | 213.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,427 | 6.8 | 69.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,988 | 5.6 | 56.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,580 | 4.5 | 45.2 |
| 6 | Diabetes mellitus (E10-E14) | 1,064 | 3.0 | 30.5 |
| 7 | Influenza and pneumonia (J10-J18) | 914 | 2.6 | 26.2 |
| 8 | Alzheimer's disease (G30) | 754 | 2.1 | 21.6 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 501 | 1.4 | 14.3 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 500 | 1.4 | 14.3 |
| 11 | Septicemia (A40-A41) | 430 | 1.2 | 12.3 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 428 | 1.2 | 12.3 |
| 13 | Atherosclerosis (I70) | 330 | 0.9 | 9.4 |
| 14 | Pneumonitis due to solids and liquids (J69) | 222 | 0.6 | 6.4 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 221 | 0.6 | 6.3 |
| | All other causes (Residual) | 5,439 | 15.3 | 155.7 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|---------|------------------|---------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Oregon | | | |
| | All causes | 31,119 | 100.0 | 883.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,262 | 23.3 | 206.2 |
| 2 | Malignant neoplasms (C00-C97) | 7,249 | 23.3 | 205.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,645 | 8.5 | 75.1 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,845 | 5.9 | 52.4 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,397 | 4.5 | 39.7 |
| 6 | Alzheimer's disease (G30) | 1,124 | 3.6 | 31.9 |
| 7 | Diabetes mellitus (E10-E14) | 1,041 | 3.3 | 29.6 |
| 8 | Influenza and pneumonia (J10-J18) | 666 | 2.1 | 18.9 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 518 | 1.7 | 14.7 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 367 | 1.2 | 10.4 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 364 | 1.2 | 10.3 |
| 12 | Parkinson's disease (G20-G21) | 304 | 1.0 | 8.6 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 266 | 0.9 | 7.6 |
| 14 | Atherosclerosis (I70) | 211 | 0.7 | 6.0 |
| 15 | Aortic aneurysm and dissection (I71) | 198 | 0.6 | 5.6 |
| | All other causes (Residual) | 5,662 | 18.2 | 160.8 |
| | Pennsylvania | | | |
| | All causes | 130,223 | 100.0 | 1,055.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 38,852 | 29.8 | 315.0 |
| 2 | Malignant neoplasms (C00-C97) | 29,849 | 22.9 | 242.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 8,579 | 6.6 | 69.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 6,017 | 4.6 | 48.8 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,728 | 3.6 | 38.3 |
| 6 | Diabetes mellitus (E10-E14) | 3,708 | 2.8 | 30.1 |
| 7 | Influenza and pneumonia (J10-J18) | 2,957 | 2.3 | 24.0 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,944 | 2.3 | 23.9 |
| 9 | Alzheimer's disease (G30) | 2,823 | 2.2 | 22.9 |
| 10 | Septicemia (A40-A41) | 2,566 | 2.0 | 20.8 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,341 | 1.0 | 10.9 |
| 12 | Pneumonitis due to solids and liquids (J69) | 1,280 | 1.0 | 10.4 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,156 | 0.9 | 9.4 |
| 14 | Parkinson's disease (G20-G21) | 1,031 | 0.8 | 8.4 |
| 15 | Aortic aneurysm and dissection (I71) | 850 | 0.7 | 6.9 |
| | All other causes (Residual) | 21,542 | 16.5 | 174.6 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Rhode Island | | | |
| | All causes | 10,246 | 100.0 | 957. |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,109 | 30.3 | 290. |
| <u>-</u> | Malignant neoplasms (C00-C97) | 2,404 | 23.5 | 224 |
| } | Cerebrovascular diseases (I60-I69) | 605 | 5.9 | 56 |
| | Chronic lower respiratory diseases (J40-J47) | 521 | 5.1 | 48 |
| | Influenza and pneumonia (J10-J18) | 319 | 3.1 | 29 |
| | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 277 | 2.7 | 25 |
| | Alzheimer's disease (G30) | 264 | 2.6 | 24 |
| | Diabetes mellitus (E10-E14) | 263 | 2.6 | 24 |
| | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 142 | 1.4 | 13 |
| 0 | Septicemia (A40-A41) | 134 | 1.3 | 12 |
| 1 | Chronic liver disease and cirrhosis (K70,K73-K74) | 128 | 1.2 | 12 |
| 2 | Parkinson's disease (G20-G21) | 87 | 8.0 | 8 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 86 | 0.8 | 8 |
| 4 | Pneumonitis due to solids and liquids (J69) | 65 | 0.6 | 6 |
| 5 | Atherosclerosis (I70) | 64 | 0.6 | 6 |
| | All other causes (Residual) | 1,778 | 17.4 | 166 |
| | South Carolina | | | |
| | All causes | 37,736 | 100.0 | 918 |
| | Diseases of heart (I00-I09,I11,I13,I20-I51) | 9,659 | 25.6 | 235 |
| | Malignant neoplasms (C00-C97) | 8,333 | 22.1 | 202 |
| | Cerebrovascular diseases (I60-I69) | 2,822 | 7.5 | 68 |
| | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,972 | 5.2 | 48 |
| | Chronic lower respiratory diseases (J40-J47) | 1,889 | 5.0 | 46 |
| | Diabetes mellitus (E10-E14) | 1,112 | 2.9 | 2 |
| | Alzheimer's disease (G30) | 967 | 2.6 | 2 |
| | Influenza and pneumonia (J10-J18) | 910 | 2.4 | 2 |
| | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 781 | 2.1 | 19 |
|) | Septicemia (A40-A41) | 655 | 1.7 | 1 |
| | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 440 | 1.2 | 10 |
| <u> </u> | Chronic liver disease and cirrhosis (K70,K73-K74) | 386 | 1.0 | , |
| 3 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 362 | 1.0 | ; |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 326 | 0.9 | • |
| 5 | Pneumonitis due to solids and liquids (J69) | 309 | 0.8 | • |
| | All other causes (Residual) | 6,813 | 18.1 | 16 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | South Dakota | | | |
| | All causes | 6,898 | 100.0 | 906.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,937 | 28.1 | 254.5 |
| 2 | Malignant neoplasms (C00-C97) | 1,562 | 22.6 | 205.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 518 | 7.5 | 68. |
| 4 | Chronic lower respiratory diseases (J40-J47) | 383 | 5.6 | 50.3 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 348 | 5.0 | 45. |
| 6 | Influenza and pneumonia (J10-J18) | 240 | 3.5 | 31.5 |
| 7 | Diabetes mellitus (E10-E14) | 195 | 2.8 | 25.0 |
| 8 | Alzheimer's disease (G30) | 167 | 2.4 | 21.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 128 | 1.9 | 16.8 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 94 | 1.4 | 12.4 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 75 | 1.1 | 9.9 |
| 12 | Septicemia (A40-A41) | 69 | 1.0 | 9. |
| 13 | Aortic aneurysm and dissection (I71) | 61 | 0.9 | 8.0 |
| 14 | Parkinson's disease (G20-G21) | 58 | 0.8 | 7.6 |
| 15 | Pneumonitis due to solids and liquids (J69) | 42 | 0.6 | 5.5 |
| | All other causes (Residual) | 1,021 | 14.8 | 134.2 |
| | Tennessee | | | |
| | All causes | 56,606 | 100.0 | 976.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 16,226 | 28.7 | 279.9 |
| 2 | Malignant neoplasms (C00-C97) | 12,518 | 22.1 | 215.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,980 | 7.0 | 68.7 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 3,011 | 5.3 | 51.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,744 | 4.8 | 47.3 |
| 6 | Diabetes mellitus (E10-E14) | 1,749 | 3.1 | 30.2 |
| 7 | Influenza and pneumonia (J10-J18) | 1,710 | 3.0 | 29. |
| 3 | Alzheimer's disease (G30) | 1,299 | 2.3 | 22.4 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 778 | 1.4 | 13.4 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 611 | 1.1 | 10. |
| 11 | Septicemia (A40-A41) | 605 | 1.1 | 10.4 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 586 | 1.0 | 10. |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 530 | 0.9 | 9. |
| 14 | Pneumonitis due to solids and liquids (J69) | 485 | 0.9 | 8.4 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 467 | 0.8 | 8.′ |
| | All other causes (Residual) | 9,307 | 16.4 | 160.5 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|-------------------------------------------------------------------------------|---------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Texas | | | |
| | All causes | 155,524 | 100.0 | 714.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 43,452 | 27.9 | 199.5 |
| 2 | Malignant neoplasms (C00-C97) | 34,164 | 22.0 | 156.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 10,548 | 6.8 | 48.4 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 8,232 | 5.3 | 37.8 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 7,720 | 5.0 | 35.4 |
| 6 | Diabetes mellitus (E10-E14) | 5,654 | 3.6 | 26.0 |
| 7 | Alzheimer's disease (G30) | 3,793 | 2.4 | 17.4 |
| 8 | Influenza and pneumonia (J10-J18) | 3,673 | 2.4 | 16.9 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,311 | 1.5 | 10.6 |
| 10 | Septicemia (A40-A41) | 2,286 | 1.5 | 10.5 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,284 | 1.5 | 10.5 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,166 | 1.4 | 9.9 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,421 | 0.9 | 6.5 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1,255 | 0.8 | 5.8 |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,075 | 0.7 | 4.9 |
| | All other causes (Residual) | 25,490 | 16.4 | 117.0 |
| | Utah | | | |
| | All causes | 13,116 | 100.0 | 566.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,977 | 22.7 | 128.5 |
| 2 | Malignant neoplasms (C00-C97) | 2,376 | 18.1 | 102.6 |
| 3 | Cerebrovascular diseases (I60-I69) | 903 | 6.9 | 39.0 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 714 | 5.4 | 30.8 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 603 | 4.6 | 26.0 |
| 6 | Diabetes mellitus (E10-E14) | 514 | 3.9 | 22.2 |
| 7 | Influenza and pneumonia (J10-J18) | 424 | 3.2 | 18.3 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 340 | 2.6 | 14.7 |
| 9 | Alzheimer's disease (G30) | 303 | 2.3 | 13.1 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 184 | 1.4 | 7.9 |
| 11 | Parkinson's disease (G20-G21) | 144 | 1.1 | 6.2 |
| 12 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99 | 136 | 1.0 | 5.9 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 133 | 1.0 | 5.7 |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 124 | 0.9 | 5.4 |
| 15 | Septicemia (A40-A41) | 109 | 8.0 | 4.7 |
| | All other causes (Residual) | 3,132 | 23.9 | 135.2 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | | Percent of total | |
|-------------------|------------------------------------------------------------------------------|--------|------------------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Vermont | | | |
| | All causes | 5,075 | 100.0 | 823.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,370 | 27.0 | 222.2 |
| 2 | Malignant neoplasms (C00-C97) | 1,224 | 24.1 | 198.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 335 | 6.6 | 54.3 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 276 | 5.4 | 44.8 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 240 | 4.7 | 38.9 |
| 6 | Diabetes mellitus (E10-E14) | 174 | 3.4 | 28.2 |
| 7 | Alzheimer's disease (G30) | 163 | 3.2 | 26.4 |
| 8 | Influenza and pneumonia (J10-J18) | 111 | 2.2 | 18.0 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 92 | 1.8 | 14.9 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 66 | 1.3 | 10.7 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 53 | 1.0 | 8.6 |
| 12 | Aortic aneurysm and dissection (I71) | 48 | 0.9 | 7.8 |
| 13 | Septicemia (A40-A41) | 47 | 0.9 | 7.6 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 37 | 0.7 | 6.0 |
| 15 | Parkinson's disease (G20-G21) | 36 | 0.7 | 5.8 |
| | All other causes (Residual) | 803 | 15.8 | 130.2 |
| | Virginia | | | |
| | All causes | 57,196 | 100.0 | 784.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 14,952 | 26.1 | 205.0 |
| 2 | Malignant neoplasms (C00-C97) | 13,602 | 23.8 | 186.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,960 | 6.9 | 54.3 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,752 | 4.8 | 37.7 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,479 | 4.3 | 34.0 |
| 6 | Diabetes mellitus (E10-E14) | 1,558 | 2.7 | 21.4 |
| 7 | Influenza and pneumonia (J10-J18) | 1,480 | 2.6 | 20.3 |
| 8 | Alzheimer's disease (G30) | 1,368 | 2.4 | 18.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,236 | 2.2 | 16.9 |
| 10 | Septicemia (A40-A41) | 1,156 | 2.0 | 15.8 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 799 | 1.4 | 11.0 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 598 | 1.0 | 8.2 |
| 13 | Pneumonitis due to solids and liquids (J69) | 589 | 1.0 | 8.1 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 479 | 0.8 | 6.6 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 397 | 0.7 | 5.4 |
| | All other causes (Residual) | 9,791 | 17.1 | 134.2 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and State | Percent of total | | |
|-------------------|--------------------------------------------------------------------------------------------------------|------------------|--------|---------|
| | | Number | deaths | Rate |
| | Washington | | | |
| | All causes | 45,338 | 100.0 | 747.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,141 | 24.6 | 183.6 |
| 2 | Malignant neoplasms (C00-C97) | 10,858 | 23.9 | 178.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,753 | 8.3 | 61.8 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,721 | 6.0 | 44.8 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,203 | 4.9 | 36.3 |
| 6 | Alzheimer's disease (G30) | 2,195 | 4.8 | 36.2 |
| 7 | Diabetes mellitus (E10-E14) | 1,494 | 3.3 | 24.6 |
| 8 | Influenza and pneumonia (J10-J18) | 907 | 2.0 | 14.9 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 811 | 1.8 | 13.4 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 525 | 1.2 | 8.7 |
| 11 | Parkinson's disease (G20-G21) | 432 | 1.0 | 7.1 |
| 12 | Pneumonitis due to solids and liquids (J69) | 407 | 0.9 | 6.7 |
| 13 | Atherosclerosis (I70) | 356 | 0.8 | 5.9 |
| 14 | Septicemia (A40-A41) | 352 | 0.8 | 5.8 |
| 15 | Aortic aneurysm and dissection (I71) | 319 | 0.7 | 5.3 |
| | All other causes (Residual) | 6,864 | 15.1 | 113.1 |
| | West Virginia | | | |
| | All causes | 21,016 | 100.0 | 1,166.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,189 | 29.4 | 343.5 |
| 2 | Malignant neoplasms (C00-C97) | 4,652 | 22.1 | 258.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,260 | 6.0 | 69.9 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,228 | 5.8 | 68.2 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 956 | 4.5 | 53.1 |
| 6 | Diabetes mellitus (E10-E14) | 846 | 4.0 | 47.0 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 462 | 2.2 | 25.6 |
| 8 | Influenza and pneumonia (J10-J18) | 427 | 2.0 | 23.7 |
| 9 | Alzheimer's disease (G30) | 405 | 1.9 | 22.5 |
| 10 | Septicemia (A40-A41) | 295 | 1.4 | 16.4 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 276 | 1.3 | 15.3 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 209 | 1.0 | 11.6 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 175 | 0.8 | 9.7 |
| 14 | Atherosclerosis (I70) | 149 | 0.7 | 8.3 |
| 15 | Pneumonitis due to solids and liquids (J69) | 139 | 0.7 | 7.7 |
| | All other causes (Residual) | 3,348 | 15.9 | 185.8 |

¹ Rank based on number of deaths

^{...} Category not applicable

LCWK9. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2002

[Rates per 100,000 population.]

| | Cause of death (Based on the Tenth Revision, International Classification of | Percent of total | | |
|-------------------|------------------------------------------------------------------------------|------------------|--------|-------|
| Rank ¹ | Diseases, 1992) and State | Number | deaths | Rate |
| | Wisconsin | | | |
| | All causes | 46,981 | 100.0 | 863.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 12,923 | 27.5 | 237.5 |
| 2 | Malignant neoplasms (C00-C97) | 10,828 | 23.0 | 199.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,479 | 7.4 | 63.9 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,335 | 5.0 | 42.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,274 | 4.8 | 41.8 |
| 6 | Diabetes mellitus (E10-E14) | 1,353 | 2.9 | 24.9 |
| 7 | Alzheimer's disease (G30) | 1,344 | 2.9 | 24.7 |
| 8 | Influenza and pneumonia (J10-J18) | 1,291 | 2.7 | 23.7 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 852 | 1.8 | 15.7 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 627 | 1.3 | 11.5 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 437 | 0.9 | 8.0 |
| 12 | Septicemia (A40-A41) | 426 | 0.9 | 7.8 |
| 13 | Parkinson's disease (G20-G21) | 404 | 0.9 | 7.4 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 396 | 0.8 | 7.3 |
| 15 | Aortic aneurysm and dissection (I71) | 337 | 0.7 | 6.2 |
| | All other causes (Residual) | 7,675 | 16.3 | 141.1 |
| | Wyoming | | | |
| | All causes | 4,174 | 100.0 | 837.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,005 | 24.1 | 201.5 |
| 2 | Malignant neoplasms (C00-C97) | 859 | 20.6 | 172.2 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 324 | 7.8 | 65.0 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 289 | 6.9 | 58.0 |
| 5 | Cerebrovascular diseases (I60-I69) | 243 | 5.8 | 48.7 |
| 6 | Diabetes mellitus (E10-E14) | 145 | 3.5 | 29.1 |
| 7 | Influenza and pneumonia (J10-J18) | 135 | 3.2 | 27.1 |
| 8 | Alzheimer's disease (G30) | 122 | 2.9 | 24.5 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 105 | 2.5 | 21.1 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 70 | 1.7 | 14.0 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 43 | 1.0 | 8.6 |
| 12 | Septicemia (A40-A41) | 41 | 1.0 | 8.2 |
| 13 | Aortic aneurysm and dissection (I71) | 38 | 0.9 | 7.6 |
| 14 | Parkinson's disease (G20-G21) | 23 | 0.6 | 4.6 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 23 | 0.6 | 4.6 |
| | All other causes (Residual) | 709 | 17.0 | 142.2 |

¹ Rank based on number of deaths

^{...} Category not applicable