

Table 17. Reported AIDS cases, by age category, transmission category, and sex, 2005 and cumulative—United States and dependent areas

| Transmission category | Males | | | | Females | | | | Total | | | |
|---|---------------|------------|-------------------------|------------|---------------|------------|-------------------------|------------|---------------|------------|-------------------------|------------|
| | 2005 | | Cumulative ^a | | 2005 | | Cumulative ^a | | 2005 | | Cumulative ^b | |
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Adult or adolescent | | | | | | | | | | | | |
| Male-to-male sexual contact | 14,819 | 48 | 416,232 | 54 | — | — | — | — | 14,819 | 35 | 416,232 | 44 |
| Injection drug use | 4,168 | 13 | 159,676 | 21 | 2,047 | 19 | 65,534 | 36 | 6,215 | 15 | 225,210 | 24 |
| Male-to-male sexual contact and injection drug use | 1,742 | 6 | 62,940 | 8 | — | — | — | — | 1,742 | 4 | 62,940 | 7 |
| Hemophilia/coagulation disorder | 68 | 0 | 5,142 | 1 | 11 | 0 | 340 | 0 | 79 | 0 | 5,482 | 1 |
| High-risk heterosexual contact ^c | 3,110 | 10 | 46,533 | 6 | 4,918 | 45 | 79,872 | 44 | 8,028 | 19 | 126,405 | 13 |
| Sex with injection drug user | 389 | 1 | 11,371 | 1 | 751 | 7 | 25,310 | 14 | 1,140 | 3 | 36,681 | 4 |
| Sex with bisexual male | — | — | — | — | 223 | 2 | 4,876 | 3 | 223 | 1 | 4,876 | 1 |
| Sex with person with hemophilia | 3 | 0 | 81 | 0 | 13 | 0 | 493 | 0 | 16 | 0 | 574 | 0 |
| Sex with HIV-infected transfusion recipient | 13 | 0 | 529 | 0 | 27 | 0 | 767 | 0 | 40 | 0 | 1,296 | 0 |
| Sex with HIV-infected person, risk factor not specified | 2,705 | 9 | 34,552 | 5 | 3,904 | 36 | 48,426 | 27 | 6,609 | 16 | 82,978 | 9 |
| Receipt of blood transfusion, blood components, or tissue ^d | 73 | 0 | 5,190 | 1 | 87 | 1 | 4,144 | 2 | 160 | 0 | 9,334 | 1 |
| Other/risk factor not reported or identified ^e | 6,976 | 23 | 69,095 | 9 | 3,881 | 35 | 31,879 | 18 | 10,857 | 26 | 100,975 | 11 |
| Subtotal | 30,956 | 100 | 764,808 | 100 | 10,944 | 100 | 181,769 | 100 | 41,900 | 100 | 946,578 | 100 |
| Child (<13 yrs at diagnosis) | | | | | | | | | | | | |
| Hemophilia/coagulation disorder | 0 | 0 | 223 | 5 | 0 | 0 | 7 | 0 | 0 | 0 | 230 | 2 |
| Mother with documented HIV infection or 1 of the following risk factors | 45 | 92 | 4,274 | 89 | 41 | 93 | 4,363 | 95 | 86 | 92 | 8,637 | 91 |
| Injection drug use | 12 | 24 | 1,660 | 34 | 4 | 9 | 1,652 | 36 | 16 | 17 | 3,312 | 35 |
| Sex with injection drug user | 2 | 4 | 783 | 16 | 3 | 7 | 744 | 16 | 5 | 5 | 1,527 | 16 |
| Sex with bisexual male | 2 | 4 | 98 | 2 | 0 | 0 | 104 | 2 | 2 | 2 | 202 | 2 |
| Sex with person with hemophilia | 0 | 0 | 21 | 0 | 0 | 0 | 16 | 0 | 0 | 0 | 37 | 0 |
| Sex with HIV-infected transfusion recipient | 0 | 0 | 11 | 0 | 0 | 0 | 15 | 0 | 0 | 0 | 26 | 0 |
| Sex with HIV-infected person, risk factor not specified | 11 | 22 | 733 | 15 | 17 | 39 | 760 | 16 | 28 | 30 | 1,493 | 16 |
| Receipt of blood transfusion, blood components, or tissue | 1 | 2 | 69 | 1 | 0 | 0 | 82 | 2 | 1 | 1 | 151 | 2 |
| Has HIV infection, risk factor not specified | 17 | 35 | 899 | 19 | 17 | 39 | 990 | 21 | 34 | 37 | 1,889 | 20 |
| Receipt of blood transfusion, blood components, or tissue ^f | 0 | 0 | 244 | 5 | 0 | 0 | 142 | 3 | 0 | 0 | 386 | 4 |
| Other/risk factor not reported or identified ^g | 4 | 8 | 86 | 2 | 3 | 7 | 102 | 2 | 7 | 8 | 188 | 2 |
| Subtotal | 49 | 100 | 4,827 | 100 | 44 | 100 | 4,614 | 100 | 93 | 100 | 9,441 | 100 |
| Total | 31,005 | 100 | 769,635 | 100 | 10,988 | 100 | 186,383 | 100 | 41,993 | 100 | 956,019 | 100 |

^a From the beginning of the epidemic through 2005.

^b From the beginning of the epidemic through 2005. Includes 1 person of unknown sex.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d AIDS developed in 47 adults/adolescents after they received blood that had tested negative for HIV antibodies. AIDS developed in 14 additional adults after they received tissue, organs, or artificial insemination from HIV-infected donors. Four of the 14 received tissue or organs from a donor who was negative for HIV antibody at the time of donation.

^e Includes 36 adults/adolescents who were exposed to HIV-infected blood, body fluids, or concentrated virus in health care, laboratory, or household settings, as supported by seroconversion, epidemiologic, and/or laboratory evidence. One person was infected after intentional inoculation with HIV-infected blood. Includes an additional 535 persons who acquired HIV infection perinatally but who were more than 12 years of age when AIDS was diagnosed. These 535 persons are not counted in the values for the pediatric transmission category.

^f AIDS developed in 3 children after they received blood that had tested negative for HIV antibodies.

^g Includes 5 children who were exposed to HIV-infected blood, as supported by seroconversion, epidemiologic, and/or laboratory evidence: 1 child was infected after intentional inoculation with HIV-infected blood, and 4 children were exposed to HIV-infected blood in a household setting. Of the 188 children, 24 had sexual contact with an adult with, or at high risk for, HIV infection.