

**Table 6. Estimated numbers of AIDS cases in adult and adolescent Hispanics, by transmission category and place of birth, 2005—United States and dependent areas (Revised June 2007)**

Transmission category	Place of birth											
	United States		Central/South America		Cuba		Mexico		Puerto Rico		Total <sup>a</sup>	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact	1,229	44	443	51	94	52	875	59	254	17	3,602	42
Injection drug use	684	24	94	11	24	13	161	11	652	42	2,073	24
Male-to-male sexual contact and injection drug use	131	5	21	2	14	8	62	4	65	4	345	4
High-risk heterosexual contact <sup>b</sup>	716	26	292	34	50	27	367	25	554	36	2,545	29
Other <sup>c</sup>	48	2	11	1	0	0	23	2	10	1	109	1
<b>Total<sup>d</sup></b>	<b>2,809</b>	<b>100</b>	<b>861</b>	<b>100</b>	<b>182</b>	<b>100</b>	<b>1,488</b>	<b>100</b>	<b>1,536</b>	<b>100</b>	<b>8,674</b>	<b>100</b>

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

<sup>a</sup> Includes 246 persons whose place of birth is not among those listed and 1,553 persons whose place of birth is unknown.

<sup>b</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>c</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>d</sup> Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.