Table 9. Estimated numbers of persons living with HIV/AIDS at the end of 2005, by race/ethnicity, sex, and transmission category—33 states with confidential name-based HIV infection reporting (Revised June 2007)

Transmission category	White, not Hispanic		Black, not Hispanic		Hispanic		Asian/Pacific Islander		American Indian/Alaska Native		Total ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	104,184	77	66,130	48	33,729	55	1,551	67	900	61	207,810	61
Injection drug use	12,476	9	32,096	23	14,806	24	264	11	228	15	60,259	18
Male-to-male sexual contact and injection drug use	10,875	8	9,305	7	3,471	6	87	4	188	13	24,083	7
High-risk heterosexual contact ^b	6,436	5	30,066	22	8,432	14	375	16	143	10	45,835	13
Other ^C	1,518	1	1,531	1	414	1	32	1	18	1	3,538	1
Subtotal	135,490	100	139,128	100	60,851	100	2,308	100	1,477	100	341,524	100
Female adult or adolescent												
Injection drug use	8,120	33	19,358	24	5,532	29	107	16	164	29	33,501	26
High-risk heterosexual contact ^b	15,719	65	60,471	74	13,336	69	522	80	381	68	91,087	72
Other ^C	458	2	1,520	2	342	2	23	4	13	2	2,376	2
Subtotal	24,298	100	81,349	100	19,211	100	652	100	558	100	126,964	100
Child (<13 yrs at diagnosis)												
Perinatal	793	83	3,980	92	1,191	90	30	83	19	95	6,051	90
Other ^d	165	17	357	8	136	10	6	17	1	5	675	10
Subtotal	958	100	4,336	100	1,327	100	36	100	20	100	6,726	100
Total ^e	160,746		224,815		81,389		2,996		2,055		475,220	

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

Data include persons with a diagnosis of HIV infection (not AIDS), a diagnosis of HIV infection and a later diagnosis of AIDS, or concurrent diagnoses of HIV infection and AIDS.

See Technical Notes for the list of 33 states that have had laws or regulations requiring confidential name-based HIV infection reporting since at least 2001.

^a Includes 3,218 persons of unknown race or multiple races.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^C Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

e Includes 6 persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.