



Forest Insect
& Disease
Leaflet 154

U.S. Department
of Agriculture
Forest Service

Phomopsis Blight of Junipers

Glenn W. Peterson¹ and C. S. Hodges, Jr.²



F-702969

Phomopsis blight has been a serious problem for more than 75 years in nurseries producing juniper seedlings and grafts. *Phomopsis juniperovora* Hahn, the fungus causing this disease, is widespread in the United States (fig. 1).

¹ Plant Pathologist, U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station, Lincoln, Nebr.

² Plant Pathologist, U.S. Department of Agriculture, Forest Service, Pacific Southwest Forest and Range Experiment Station, Institute of Pacific Islands Forestry, Honolulu, Hawaii.

Description

Losses have been most severe in seedling and transplant beds of eastern redcedar and Rocky Mountain juniper. (See cover photo.) Other junipers are susceptible, as are some species in the genera *Chamaecyparis*, *Cupressus*, and *Thuja*. Arizona cypress seedlings have been seriously damaged in some Southern States.

Phomopsis juniperovora initially infects foliage, then spreads to and sometimes kills stem tissues. Newly developing needles are especially susceptible while they are still in

When junipers in landscape plantings become infected, they may become unsightly because of numerous dead branch tips (fig. 3). Older trees are seldom killed because only small-diameter stems are girdled. For this reason, *Phomopsis* blight does not cause significant damage in natural stands of junipers.

Total loss of first-year seedlings is common in epidemic years if control measures are not used. Losses are particularly high in areas where water tends to stand and in new seedlings adjacent to beds of infected stock. Some of the worst epidemics occur late in the growing season, when there is a late flush of growth on juniper seedlings.

Damage from drought may be confused with *Phomopsis* blight. In both cases, tips of branches may be killed. However, the demarcation

between green and dead tissues is sharp in *Phomopsis*-blighted seedlings and gradual in seedlings affected by drought. Damage from the lesser cornstalk borer can be distinguished from *Phomopsis* blight by the straw color of the dead tops and by the feeding wounds on the lower stem and taproot present in the former. Needle blight of junipers and other species in the Cupressaceae caused by *Cercospora sequoiae* Ell. & Ev. var. *juniperi* Ell. & Ev. can be easily distinguished from *Phomopsis* blight. *Cercospora* infection starts on the oldest needles of lower branches and spreads upward and outward, while *Phomopsis* infection starts at or near the tips of the shoots.

Recently, *Kabatina juniperi* Schneider & v. Arx has become a problem in juniper outplantings and in production of grafted junipers.



Figure 3.—Dead branch tips on eastern red-cedar caused by *Phomopsis juniperovora*.

F-702971

The fungus causes symptoms similar to those caused by *Phomopsis*. The tips of branches are killed for a short distance and the dark fruiting bodies (acervuli) develop on ashen-gray tissues, as is the case with *Phomopsis*. A specialist can recognize differences in spore size and fruiting bodies between this fungus and *Phomopsis*. The time of symptom development is also helpful in distinguishing between these two blights. Kabatina blight symptoms develop only when new growth begins in the spring, while *Phomopsis* blight symptoms develop anytime during the growing season.

Life Cycle

Spores produced in fruiting bodies (pycnidia) formed on leaves and stems of seedlings infected the previous year are the most important source of inoculum early in the growing season. Pycnidia with viable spores may develop within 3 to

4 weeks after seedlings become infected, but usually are not well developed until infected tissues have dried considerably. These spores are most commonly found on tissues that have turned ashen gray. The pycnidia are at first embedded in needles and stems, but partially erupt through the epidermis (fig. 4). Two types of spores (alpha and beta) develop in the same or different pycnidia (fig. 5) and are extruded in whitish tendrils. The fungus can produce spores for as long as 2 years in dead parts of infected plants.

Spores are dispersed primarily by rain splash. Infection is caused by alpha-spores; the filamentous beta-spores do not germinate. Only a short period of high humidity is needed for infection to occur; for example, seedlings exposed to 100 percent relative humidity (24° C) for only 7 hours can be



Figure 4.—*Pycnidia on leaves and branches.*

F-702972



F-531116

Figure 5.—*Alpha and beta (long) spores of Phomopsis juniperovora.*

come infected. Spore germination, germ-tube development, and infection are optimum near 24° C; however, disease development is enhanced by higher temperatures (32° C).

Control

Because susceptible new foliage and viable fungus spores are present throughout the growing season in juniper seedling beds, protective fungicides need to be applied regularly during this season. The only chemical currently registered for control of *Phomopsis* blight is benomyl. This chemical applied at 7- to 10-day intervals, combined with a vigorous schedule of roguing infected seedlings over the same interval, will give excellent control of *Phomopsis* blight.

Other actions can be taken to reduce losses. Sowing juniper seed adjacent to beds containing juniper stock should be avoided if possible. Poorly drained areas should be avoided because losses are often greater where water tends to stand. If overhead sprinklers are used, seedlings should be irrigated so that water on seedlings dries before nightfall. Because shading frames increase the length of time that moisture remains on foliage, they should not be used unless absolutely necessary. Junipers or other hosts of this fungus should not be used in nursery windbreaks or in landscape plantings on nursery grounds because they may be a source of inoculum (spores) for nursery stock. Such trees are more likely to be extensively infected if pruning re-

sults in the development of juvenile foliage.

There is considerable variation in susceptibility to *Phomopsis juniperovora* among junipers (fig. 6).

Research is seeking to determine if there is genetic resistance to *Phomopsis juniperovora* among and within progenies from select eastern redcedar trees.



F-702973

Figure 6.—*Eastern redcedars resistant to Phomopsis juniperovora adjacent to a severely infected juniper.*

References

- Hahn, Glenn Gardner.** Taxonomy, distribution, and pathology of *Phomopsis occulta* and *P. juniperovora*. Mycol. 35: 112-129; 1943.
- Hodges, C. S.; Green, H. J.** Survival in the plantation of eastern redcedar seedlings infected with Phomopsis blight in the nursery. Plant Dis. Rep. 45: 134-136; 1961.
- Otta, J. D.** Benomyl and thiophanate methyl control Phomopsis blight of eastern redcedar in a nursery. Plant Dis. Rep. 58: 476-477; 1974.
- Otta, J. D.** Alpha and beta spore occurrence in *Phomopsis juniperovora* pycnidia on *Juniperus virginiana* and in culture. Can. J. Bot. 56: 727-728; 1978.
- Otta, J. D.; Fiedler, D. J.; Lengkeek, V. H.** Effect of benomyl on *Phomopsis juniperovora* infection of *Juniperus virginiana*. Phytopathol. 70: 46-50; 1980.
- Peterson, Glenn W.** Cedar blight. In: Internationally dangerous forest tree diseases. Misc. Pub. 939. Washington, DC: U.S. Department of Agriculture; 1963: 105-106.
- Peterson, Glenn W.** Field survival and growth of Phomopsis-blighted and non-blighted eastern redcedar planting stock. Plant Dis. Rep. 49: 121-123; 1965.
- Peterson, Glenn W.** Chemical control of Phomopsis blight of junipers: a search for new methods. Tree Plant. Notes. 23(3): 3-4; 1972.
- Peterson, Glenn W.** Infection of *Juniperus virginiana* and *J. scopulorum* by *Phomopsis juniperovora*. Phytopathol. 63: 246-251; 1973.
- Peterson, Glenn W.; Sumner, D. R.; Norman, C.** Control of Phomopsis blight of eastern redcedar seedlings. Plant Dis. Rep. 49: 529-531; 1965.
- Peterson, Glenn W.** Phomopsis blight of junipers. In: Forest Nursery Diseases in the United States. Agric. Handb. 470. Washington, DC: U.S. Department of Agriculture; 1975: 76-79.
- Peterson, Glenn W.; Otta, J. D.** Controlling Phomopsis blight of junipers. Am. Nurseryman 149(5): 15, 75, 78, 80-82; 1979.
- Schoeneweiss, Donald F.** Susceptibility of evergreen hosts to the juniper blight fungus, *Phomopsis juniperovora*, under epidemic conditions. J. Am. Soc. Hortic. Sci. 94(6): 609-611; 1969.
- Smyly, W. B.; Filer, T. H., Jr.** Benomyl controls Phomopsis blight on Arizona cypress in a nursery. Plant Dis. Rep. 57: 59-61; 1973.

Pesticides used improperly can be injurious to man, animals, and plants. Follow the directions and heed all precautions on the labels. Store pesticides in original containers under lock and key—out of the reach of children and animals—and away from food and feed.

Apply pesticides so that they do not endanger humans, livestock, crops, beneficial insects, fish, and wildlife. Do not apply pesticides where there is danger of drift when honey bees or other pollinating insects are visiting plants, or in ways that may contaminate water or leave illegal residues.

Avoid prolonged inhalation of pesticide sprays or dusts; wear protective clothing and equipment, if specified on the label.

If your hands become contaminated with a pesticide, do not eat or drink until you have washed. In case a pesticide is swallowed or gets in the eyes, follow the first aid treatment given on the label, and get prompt medical attention. If a pesticide is spilled on your skin or clothing, remove clothing immediately and wash skin thoroughly.

NOTE: Some States have restrictions on the use of certain pesticides. Check your State and local regulations. Also, because registrations of pesticides are under constant review by the U.S. Environmental Protection Agency, consult your local forest pathologist, county agriculture agent, or State extension specialist to be sure the intended use is still registered.



Use Pesticides Safely
FOLLOW THE LABEL
U.S. DEPARTMENT OF AGRICULTURE