

References

- Aberg B, Ekman L, Falk R, Greitz U, Persson G, Snihs JO. Metabolism of methyl mercury (^{203}Hg) compounds in man. *Arch Environ Health* 1969;19:478-84.
- Adgate JL, Barr DB, Clayton CA, Eberly LE, Freeman, NCG, Lioy, PJ, et al. Measurement of children's exposure to pesticides: analysis of urinary metabolite levels in a probability-based sample. *Environ Health Perspect* 2001;109(6):583-90.
- Adibi JJ, Perera FP, Jedrychowski W, Camann DE, Barr D, Jacek R, et al. Prenatal exposures to phthalates among women in New York City and Krakow, Poland. *Environ Health Perspect* 2003;111(14):1719-22.
- Adlercreutz CH, Goldin BR, Gorbach SL, Hockerstedt KA, Watanabe S, Hamalainen EK, et al. Soybean phytoestrogens intake and cancer risk. *J Nutr* 1995a;125(3 Suppl):757S-70s.
- Adlercreutz H, Fotsis T, Bannwart C, Wahala K, Makela T, Brunow G, et al. Determination of urinary lignans and phytoestrogen metabolites, potential antiestrogens and anticarcinogens in urine of women on various habitual diets. *J Steroid Biochem* 1986;25:791-7.
- Adlercreutz H, Honjo H, Higashi A, Fotsis T, Hamalainen E, Hasegawa T, et al. Urinary excretion of lignans and isoflavone phytoestrogens in Japanese men and women consuming a traditional Japanese diet. *Am J Clin Nutr* 1991;54:1093-1100.
- Adlercreutz H, van der Wildt J, Kinzel J, Attalla H, Wahala K, Makela T, et al. Lignan and isoflavonoid conjugates in human urine. *J Steroid Biochem Mol Biol* 1995b;52(1) 97-103.
- Adonis M, Martinez V, Riquelme R, Ancic P, Gonzalez G, Tapia R, et al. Susceptibility and exposure biomarkers in people exposed to PAHs from diesel exhaust. *Toxicol Lett* 2003;144(1):3-15.
- Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological profile for Cadmium [online]. 1999. Available from URL: <http://www.atsdr.cdc.gov/toxprofiles/tp5.html>, 4/7/05.
- Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological profile for Colbalt [online]. 2004. Available from URL: <http://www.atsdr.cdc.gov/toxprofiles/tp33.html>, 4/7/05.
- Agency for Toxic Substances and Disease Registry (ATSDR). 2002 Sept. Toxicological profile for di(2-ethylhexyl) phthalate [online]. 2002. Available from URL: <http://www.atsdr.cdc.gov/toxprofiles/tp9.html>, 03/09/05.
- Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological profile for diethyl phthalate [online]. 1995. Available from URL: <http://www.atsdr.cdc.gov/toxprofiles/tp73.html>, 03/17/05.
- Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological profile for di-n-butyl phthalate [online]. 2001. Available from URL: <http://www.atsdr.cdc.gov/toxprofiles/tp135.html>, 03/17/05
- Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological profile for di-n-octyl phthalate [online]. 1997. Available from URL: <http://www.atsdr.cdc.gov/toxprofiles/tp95.html>, 03/17/05.
- Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological profile for pyrethrins and pyrethroids [online]. 2003. Available from URL: <http://www.atsdr.cdc.gov/toxprofiles/tp155.html>, 03/17/05.
- Ahlborg UG, Lindgren JE, Mercier M. Metabolism of pentachlorophenol. *Arch Toxicol* 1974;32(4):271-81.
- Akkina JE, Reif JS, Keefe TJ, Bachand AM. Age at natural menopause and exposure to organochlorine pesticides in Hispanic women. *J Toxicol Environ Health* 2004;67:1407-22.
- Albertazzi P, Pansini F, Bottazi M, Bonaccorsi G, De Aloysio D, Morton MS. Dietary soy supplementation and phytoestrogen levels. *Obstet Gynecol* 1999;94(2):229-31.
- Albro PW, Moore B. Identification of the metabolites of simple phthalate diesters in rat urine. *J Chromatogr* 1974;94:209-18.
- Albro PW, Corbett JT, Schroeder JL, Jordan S, Matthews HB. Pharmacokinetics, interactions with macromolecules and species differences in metabolism of DEHP. *Environ Health Perspect* 1982;45:19-25.
- Albro PW and Lavenhar SR. Metabolism of di(2-ethylhexyl) phthalate. *Drug Metab Rev* 1989;21:13-34.
- Alexander BH, Checkoway H, van Netten C, Muller CH, Ewers TG, Kaufman JD, et al. Semen quality of men employed at a lead smelter. *Occup Environ Med* 1996;53:411-6.
- Alexander CS. Cobalt-beer cardiomyopathy. A clinical and pathological study of twenty-eight cases. *Am J Med* 1972;53:395-417.
- Alexandersson R. Blood and urinary concentrations as estimators of cobalt exposure. *Arch Environ Health* 1988;43(4):299-303.
- Alexandrie AK, Warholm M, Carstensen U, Axmon A, Hagmar L, Levin JO, et al. CYP1A1 and GSTM1 polymorphisms affect urinary 1-hydroxypyrene levels after PAH exposure. *Carcinogenesis* 2000;21(4):669-76.

- Alfven T, Jarup L, Elinder CG. Cadmium and lead in blood in relation to low bone mineral density and tubular proteinuria. *Environ Health Perspect* 2002;110:699-702.
- Allain P, Berre S, Premel-Cabic A, Mauras Y, Cledes A, Cournot A. Urinary elimination of molybdenum by healthy subjects as determined by inductively coupled plasma mass spectrometry. *Magnesium and Trace Elements* 1991-1992;10(1):47-50.
- Ambrose AM, Yost DH. Pharmacologic and toxicologic studies of N,N-diethyltoluamide. II. N,N-Diethyl-o-toluamide and N,N-diethyl-p-toluamide. *Toxicol Appl Pharmacol* 1965;7(6):772-80.
- American Conference of Government Industrial Hygienists (ACGIH). Documentation of biological exposure indices. 7th edition. Cincinnati (OH): ACGIH Worldwide; 2001.
- Anderson HA, Falk C, Hanrahan L, Olson J, Burse VW, Needham LL, et al. Profiles of Great Lakes critical pollutants: a sentinel analysis of human blood and urine. The Great Lakes Consortium. *Environ Health Perspect* 1998;106(5):279-89.
- Anderson WA, Castle L, Scotter MJ, Massey RC, Springall C. A biomarker approach to measuring human dietary exposure to certain phthalate diesters. *Food Addit Contam* 2001;18(12):1068-74.
- Angerer J, Heinzow B, Schaller K H, Weltle D, Lehnert G. Determination of environmentally caused chlorophenol levels in urine of the general population. *Fresenius' Journal of Analytical Chemistry* 1992;342(4-5):433-8.
- Angerer J, Maass R, Heinrich R. Occupational exposure to hexachlorocyclohexane. VI. Metabolism of gamma-hexachlorocyclohexane in man. *Int Arch Occup Environ Health* 1983;52(1):59-67.
- Angerer J, Ritter A. Determination of metabolites of pyrethroids in human urine using solid-phase extraction and gas chromatography-mass spectrometry. *J Chromatogr B Biomed Sci Appl* 1997;695(2):217-26.
- Apostoli P, Baj A, Bavazzano P, Ganzi A, Neri A, Ronchi L, et al. Blood lead reference values: the results of an Italian polycentric study. *Sci Total Environ* 2002a;287:1-11.
- Apostoli P, Cortesi I, Mangili A, Elia G, Drago I, Gagliardi T, et al. Assessment of reference values for mercury in urine: the results of an Italian polycentric study. *Sci Total Environ* 2002b;289:13-24.
- Apostoli P, Schaller KH. Urinary beryllium--a suitable tool for assessing occupational and environmental beryllium exposure. *Int Arch Occup Environ Health* 2001;74:162-6.
- Aprea C, Betta A, Catenacci G, Lotti A, Magnaghi S, Barisano A, et al. Reference values of urinary 3,5,6-trichloro-2-pyridinol in the Italian population--validation of analytical method and preliminary results (multicentric study). *J AOAC Int* 1999;82:305-12.
- Aprea C, Sciarra G, Orsi D, Boccalon P, Sartorelli P, Sartorelli E. Urinary excretion of alkylphosphates in the general population (Italy). *Sci Total Environ* 1996;177:37-41.
- Aprea C, Strambi M, Novelli MT, Lunghini L, Bozzi N. Biologic monitoring of exposure to organophosphorus pesticides in 195 Italian children. *Environ Health Perspect* 2000;108:521-5.
- Arai Y, Uehara M, Sato Y, Kimura M, Eboshida A, Adlercreutz H, et al. Comparison of isoflavones among dietary intake, plasma concentration and urinary excretion for accurate estimation of phytoestrogen intake. *J Epidemiol* 2000;10:127-35.
- Araki S, Sato H, Yokoyama K, Murata K. Subclinical neurophysiological effects of lead: a review on peripheral, central, and autonomic nervous system effects in lead workers. *Am J Ind Med* 2000;37(2):193-204.
- Arbuckle TE, Burnett R, Cole D, Teschke K, Dosemeci M, Bencej C, et al. Predictors of herbicide exposure in farm applicators. *Int Arch Occup Environ Health* 2002;75(6):406-14.
- Arisawa K, Matsumura T, Tohyama C, Saito H, Satoh H, Nagai M, et al. Fish intake, plasma omega-3 polyunsaturated fatty acids, and polychlorinated dibenzo-*p*-dioxins/polychlorinated dibenzo-furans and co-planar polychlorinated biphenyls in the blood of the Japanese population. *Int Arch Occup Environ Health* 2003;76(3):205-15.
- Armitage AK, Dollery CT, George CF, Houseman TH, Lewis PJ, Turner DM. Absorption and metabolism of nicotine from cigarettes. *BMJ* 1975;4:313-6.
- Armstrong B, Hutchinson E, Unwin J, Fletcher T. Lung cancer risk after exposure to polycyclic aromatic hydrocarbons: a review and meta-analysis. *Environ Health Perspect* 2004;112(9):970-8.
- Aylward LL, Hays SM. Temporal trends in human TCDD body burden: decreases over three decades and implications for exposure levels. *J Expo Anal Environ Epidemiol* 2002;12(5):319-28.
- Baccarelli A, Mocarelli P, Patterson DG Jr, Bonzini M, Pesatori AC, Caporaso N, et al. Immunologic effects of dioxin: new results from Seveso and comparison with other studies. *Environ Health Perspect* 2002;110:1169-73.
- Bachthaler M, Lenhart M, Paetzl C, Feuerbach S, Link J, Manke C. Corrosion of tungsten coils after peripheral vascular embolization therapy: influence on outcome and tungsten load. *Catheterization and Cardiovascular Interventions* 2004;62:380-84.

- Bailly R, Lauwerys R, Buchet JP, Mahieu P, Konings J. Experimental and human studies on antimony metabolism: their relevance for the biological monitoring of workers exposed to inorganic antimony. *British Journal of Industrial Medicine* 1991;48:93-7.
- Baker DB, Loo S, Barker J. Evaluation of human exposure to the heptachlor epoxide contamination of milk in Hawaii. *Hawaii Med J* 1991;50(3):108-12,118.
- Baker SE, Barr DB, Driskell WJ, Beeson MD, Needham LL. Quantification of selected pesticide metabolites in human urine using isotope dilution high-performance liquid chromatography/tandem mass spectrometry. *J Expo Anal Environ Epidemiol* 2000;10:789-98.
- Baker SE, Olsson AO, Barr DB. Isotope dilution high-performance liquid chromatography-tandem mass spectrometry method for quantifying urinary metabolites of synthetic pyrethroid insecticides. *Arch Environ Contam Toxicol* 2004;46:281-8.
- Balluz L, Philen R, Ortega L, Rosales C, Brock J, Barr D, et al. Investigation of systemic lupus erythematosus in Nogales, Arizona. *Am J Epidemiol* 2001;154(11):1029-36.
- Barr DB, Wilder LC, Caudill SP, Gonzalez AJ, Needham LL, Pirkle JL. Urinary creatinine concentrations in the U.S. population: implications for urinary biologic monitoring measurements. *Environ Health Perspect* 2005;113(2):192-200.
- Barr DB, Silva MJ, Kato K, Reidy JA, Malek NA, Hurtz D, et al. Assessing human exposure to phthalates using monoesters and their oxidized metabolites as biomarkers. *Environ Health Perspect* 2003;111(9):1148-51.
- Barr DB, Turner WE, DiPietro E, McClure PC, Baker SE, Barr JR, et al. Measurement of p-nitrophenol in the urine of residents whose homes were contaminated with methyl parathion. *Environ Health Perspect Suppl* 2002;110(6):1085-91.
- Barregard L, Sallsten G, Conradi N. Tissue levels of mercury determined in a deceased worker after occupational exposure. *Int Arch Occup Environ Health* 1999;72:169-73.
- Barregard L, Sallsten G, Schutz A, Attewell R, Skerfving S, Jarvholm B. Kinetics of mercury in blood and urine after brief occupational exposure. *Arch Environ Health* 1992;7(3):176-84.
- Bartels MJ, Kastl PE. Analysis of 3,5,6-trichloropyridinol in human urine using negative-ion chemical ionization gas chromatography-mass spectrometry. *J Chromatogr* 1992;575(1):69-74.
- Bates MN, Buckland SJ, Garrett N, Ellis H, Needham LL, Patterson DG Jr, et al. Persistent organochlorines in the serum of the non-occupationally exposed New Zealand population. *Chemosphere* 2004b;54:1431-43.
- Bates MN, Fawcett J, Garrett N, Cutress T, Kjellstrom T. Health effects of dental amalgam exposure: a retrospective cohort study. *Int J Epidemiol* 2004a;33:1-9.
- Bazzano LA, He J, Muntner P, Vupputuri S, Whelton PK. Relationship between cigarette smoking and novel risk factors for cardiovascular disease in the United States. *Ann Intern Med* 2003;138(11):891-7.
- Beck H, Dross A, Mathar W. PCDD and PCDF exposure and levels in humans in Germany. *Environ Health Perspect* 1994;102 Suppl 1:173-85.
- Becker K, Kaus S, Krause C, Lepom P, Schulz C, Seiwert M, et al. German Environmental Survey 1998 (GerES III): environmental pollutants in blood of the German population. *International Journal of Hygiene and Environmental Health* 2002;205:297-308.
- Becker K, Seiwert M, Angerer J, Heger W, Koch HM, Nagorka R, et al. DEHP metabolites in urine of children and DEHP in house dust. *International Journal of Hygiene and Environmental Health* 2004;207(5):409-17.
- Becker K, Schulz C, Kaus S, Seiwert M, Seifert B. German environmental survey 1998 (GerES III): environmental pollutants in the urine of the German population. *International Journal of Hygiene and Environmental Health* 2003; 206:15-24.
- Bell JW, Veltri JC, Page BC. Human exposures to N,N-diethyl-m-toluamide insect repellents reported to the American Association of Poison Control Centers, 1993-1997. *International Journal of Toxicology* 2002;2:341-52.
- Benes B, Spevackova V, Smid J, Cejchanova M, Cerna M, Subrt P, et al. The concentration levels of Cd, Pb, Hg, Cu, Zn, and Se in blood of the population in the Czech Republic. *Cent Eur J Public Health* 2000;8(2):117-9.
- Benes B, Spevackova V, Smid J, Cejchanova M, Kaplanova E, Cerna M, et al. Determination of normal concentrations of Cd, Pb, Hg, Cu, Zn, and Se in urine of the population of the Czech Republic. *Cent Eur J Public Health* 2002;10:3-5.
- Benowitz NL. Cotinine as a biomarker of environmental tobacco smoke exposure. *Epidemiol Rev* 1996;18:188-204.
- Benowitz NL, Jacob P. Metabolism of nicotine to cotinine studied by a dual stable isotope method. *Clin Pharmacol Ther* 1994;56:483-93.
- Berman JD. Chemotherapy for leishmaniasis: Biochemical mechanisms, clinical efficacy, and future strategies. *Reviews in Infectious Diseases* 1988;10(3):560-86.
- Bertram HP, Kemper FH, Muller C. Hexachlorobenzene content in human whole blood and adipose tissue: experiences in environmental specimen banking. *IARC Sci Publ* 1986;77:173-82.

- Bhatnagar VK, Kashyap R, Zaidi SS, Kulkarni PK, Saiyed HN. Levels of DDT, HCH, and HCB residues in human blood in Ahmedabad, India. *Bull Environ Contam Toxicol* 2004;72:261-5.
- Bhattacharyya MH, Breitenstein BD, Metivier H, Muggenburg BA, Stradling GN, Volf V. Guidebook for the treatment of accidental internal radionuclide contamination of workers. In: Gerber GB, Thomas RG, eds. *Radiation protection dosimetry*. Vol. 41 (1). Kent (England): Nuclear Technology Publishing; 1992. p. 1-49.
- Bidstrup PL, Bonnell JA, Harvey DG, Lockett S. Chronic mercury poisoning in men repairing direct-current meters. *Lancet* 1951;2:856-61.
- Bittner AC Jr, Echeverria D, Woods JS, Aposhian HV, Naleway C, Martin MD, et al. Behavioral effects of low-level exposure to Hg⁰ among dental professionals: a cross-study evaluation of psychomotor effects. *Neurotoxicol Teratol* 1998;20(4):429-39.
- Bjornberg KA, Vahter M, Petersson-Grawe K, Glynn A, Cnattingius S, Darnerud PO, et al. Methyl mercury and inorganic mercury in Swedish pregnant women and in cord blood: influence of fish consumption. *Environ Health Perspect* 2003;111(4):637-41.
- Bloom MS, Vena JE, Swanson MK, Moysich KB, Olson JR. Profiles of ortho-polychlorinated biphenyl congeners, dichlorodiphenyldichloroethylene, hexachlorobenzene, and mirex among male Lake Ontario sportfish consumers: the New York State Angler Cohort Study. *Environ Res* 2005;97:178-94.
- Blount BC, Silva MJ, Caudill SP, Needham LL, Pirkle JL, Sampson EJ, et al. Levels of seven urinary phthalate metabolites in a human reference population. *Environ Health Perspect* 2000;108(10):979-82.
- Boffetta P, Hjourenkova N, Gustavsson P. Cancer risk from occupational and environmental exposure to polycyclic aromatic hydrocarbons. *Cancer Causes Control* 1997;8(3):444-72.
- Borja-Aburto VH, Hertz-Picciotto I, Rojas LM, Farias P, Rios C, Blanco J. Blood lead levels measured prospectively and risk of spontaneous abortion. *Am J Epidemiol* 1999;150(6):590-7.
- Bostrom CE, Gerde P, Hanberg A, Jernstrom B, Johansson C, Kyrklund T, et al. Cancer risk assessment, indicators, and guidelines for polycyclic aromatic hydrocarbons in the ambient air. *Environ Health Perspect* 2002;110Suppl 3:451-88.
- Botella B, Crespo J, Rivas A, Cerrillo I, Olea-Serrano MF, Olea N. Exposure of women to organochlorine pesticides in Southern Spain. *Environ Res* 2004;96:34-40.
- Bradberry, SM, Watt BE, Proudfoot AT, Vale JA. Mechanisms of toxicity, clinical features, and management of acute chlorophenoxy herbicide poisoning: a review. *J Toxicol Clin Toxicol* 2000;38:111-22.
- Braun WH, Blau GE, Chenoweth MB. The metabolism/pharmacokinetics of pentachlorophenol in man and a comparison with the rat and monkey. *Developments in Toxicology and Environmental Science* 1979;4:289-96.
- Brenniman GR, Levy, PS. Epidemiological study of barium in Illinois drinking water supplies. In: Calabrese EJ, ed. *Advances in modern toxicology*. Princeton (NJ): Princeton Scientific Publications; 1984. p. 231-49.
- Briegner H, Semisch CW, Stasney J, Piatnek DA. Industrial antimony poisoning. *Industrial Medicine and Surgery (Dec.)* 1954;521-3.
- Brockhaus A, Dolger R, Ewers U, Kramer U, Soddemann H, Wiegand H. Intake and health effects of thallium among a population living in the vicinity of a cement plant emitting thallium-containing dust. *Int Arch Occup Environ Health* 1981;48(4):375-89.
- Brokopp CD, Wyatt JI, Gabica J. Dialkyl phosphates in urine samples from pesticide formulators exposed to disulfoton and phorate. *Bull Environ Contam Toxicol* 1981;26(4):524-9.
- Bucher JR, Hailey JR, Roycroft JR, Haseman JK, Sills RC, Grumbein SL, et al. Inhalation toxicity and carcinogenicity studies of cobalt sulfate. *Toxicol Sci* 1999;49:56-67.
- Budavari S, ed. *The Merck index*. 11th ed. Rahway (NJ): Merck & Co., Inc; 1989. p. 308-9.
- Burns JE. Pesticides in people: organochlorine pesticide and polychlorinated biphenyl residues in biopsied human adipose tissue—Texas 1969-1972. *Pesticides Monitoring Journal* 1974;7:122-6.
- Byrne AR, Benedik L. Uranium content of blood, urine and hair of exposed and non-exposed persons determined by radiochemical neutron activation analysis, with emphasis on quality control. *Sci Total Environ* 1991;107:143-57.
- Byrne SL, Shurdut BA, Saunders DG. Potential chlorpyrifos exposure to residents following standard crack and crevice treatment. *Environ Health Perspect* 1998;106(11):725-31.
- Cabral J, Hall R, Rossi L, Bronczyk SA, Shubik P. Effects of long-term intake of DDT on rats. *Tumori* 1982;68:11-7.
- Calafat, AM, Needham LL, Silva MJ, Lambert G. Exposure to di-(2-ethylhexyl) phthalate among premature neonates in a neonatal intensive care unit. *Pediatrics* 2004;113:e429-e434.
- Calvert GM, Wille KK, Sweeney MH, Fingerhut MA, Halperin WE. Evaluation of serum lipid concentrations among U.S. workers exposed to 2,3,7,8-tetrachloro-dibenzo-p-dioxin. *Arch Environ Health* 1996;51(2):100-7.

- Canfield RL, Henderson CR, Cory-Slechta DA, Cox C, Jusko TA, Lanphear BP. Intellectual impairment in children with blood lead concentrations below 10 µg/dL. *N Engl J Med* 2003;348:1517-26.
- Caraballo R, Giovino G, Pechacek TF, Mowery PD, Richter PA, Strauss WJ, et al. Racial/ethnic differences in serum cotinine levels among adult U.S. cigarette smokers: the Third National Health and Nutrition Examination Survey, 1988-1991. *JAMA* 1998;280:135-40.
- Cardenas A, Roels H, Bernard AM, Barbon R, Buchet JP, Lauwerys RR, et al. Markers of early renal changes induced by industrial pollutants. I. Application to workers exposed to mercury vapour. *British Journal of Industrial Medicine* 1993;50:17-27.
- Cassidy A, Bingham S, Setchell KDR. Biological effects of a diet of soy protein rich in isoflavones on the menstrual cycle of premenopausal women. *Am J Clin Nutr* 1994;60(3):333-40.
- Center for the Evaluation of Risks to Human Reproduction. National Toxicology Program. National Institute of Environmental Health Sciences. [online]. 2000. Available from URL: <http://cerhr.niehs.nih.gov/news/phthalates/monographs.html>, 4/7/05.
- Centers for Disease Control. Preventing lead poisoning in young children: a statement by the Centers for Disease Control. Atlanta (GA): CDC; 1991.
- Centers for Disease Control and Prevention. Surveillance for elevated blood lead levels among children – United States, 1997-2001. *MMWR CDC Surveill Summ* September 12, 2003a;52(No. SS-10):1-21.
- Centers for Disease Control and Prevention. National Center for Environmental Health. Cancer Clusters. Churchill County (Fallon). Nevada Exposure Assessment. [online] 2003b. <http://www.cdc.gov/nceh/clusters/Fallon/study.htm>, 4/17/05.
- Chapman LJ, Sauter SL, Henning RA, Dodson VN, Reddan WG, Matthews CG. Differences in frequency of finger tremor in otherwise asymptomatic mercury workers. *British Journal of Industrial Medicine* 1990;47:838-43.
- Charles MJ, Schell MJ, Willman E, Gross HB, Lin Y, Sonnenberg S, et al. Organochlorines and 8-hydroxy-2-deoxyguanosine in cancerous and noncancerous breast tissue: do the data support the hypothesis that oxidative DNA damage caused by organochlorines affects breast cancer? *Arch Environ Contam Toxicol* 2001;41:386-95.
- Charlier C, Pitance F, Piomteux G. PCB residues in a breast cancer patient population. *Bull Environ Contam Toxicol* 2003;71:887-91.
- Chen Z, Zheng W, Custer LJ, Dai Q, Shu X, Jin F, et al. Usual dietary consumption of soy foods and its correlation with the excretion rate of isoflavonoids in overnight urine samples among Chinese women in Shanghai. *Nutr Cancer* 1999;33:82-7.
- Choudhury H, Harvey T, Thayer W, Lockwood T, Stiteler WM, Goodrum PE, et al. Urinary cadmium elimination as a biomarker of exposure for evaluating a cadmium dietary exposure – biokinetics model. *J Toxicol Environ Health* 2001;63(Pt A):321-50.
- Christensen JM, Poulsen OM. A 1982-1992 surveillance programme on Danish pottery painters. Biological levels and health effects following exposure to soluble or insoluble cobalt compounds in cobalt blue dyes. *Sci Total Environ* 1994;50(1-3):95-104.
- Chuang JC, Callahan PJ, Lyu CW, Wilson NK. Polycyclic aromatic hydrocarbon exposures of children in low-income families. *J Expo Anal Environ Epidemiol* 1999;9(2):85-98.
- Chuang CY, Lee CC, Chang YK, Sung FC. Oxidative DNA damage estimated by urinary 8-hydroxydeoxyguanosine: influence of taxi driving, smoking and areca chewing. *Chemosphere* 2003;52(7):1163-71.
- Cianciola ME, Echeverria D, Martin MD, Aposian HV, Woods JS. Epidemiologic assessment of measures used to indicate low-level exposure to mercury vapor (Hg⁰). *J Toxicol Environ Health* 1997; 52:19-33.
- Clement J, Okey A. Reproduction in female rats born to DDT-treated parents. *Bull Environ Contam Toxicol* 1974;12(3):373-7.
- Coldham NG, Dave M, Silvapathasundaram S, McDonnell DP, Connor C, Sauer MJ. Evaluation of a recombinant yeast cell estrogen screening assay. *Environ Health Perspect* 1997; 105:734-42.
- Cordasco EM, Stone FD. Pulmonary edema of environmental origin. *Chest* 1973;64(2):182-5.
- Cornwell T, Cohick W, Raskin I. Dietary phytoestrogens and health. *Phytochemistry* 2004;65(8):995-1016.
- Cotter LH. Paradichlorobenzene poisoning from insecticides. *New York State Journal of Medicine* 1953;53:1690-2.
- Cugell DW, Morgan WKC, Perkins DG, Rubin A. The respiratory effects of cobalt. *Arch Intern Med* 1990;150:177-83.
- Curl CL, Fenske RA, Kissel JC, Shirai JH, Moate TF, Griffith W, et al. Evaluation of take-home organophosphorus pesticide exposure among agricultural workers and their children. *Environ Health Perspect* 2002;110(12):787-92.
- Czerczak S, Gromiec JP. Nickel, ruthenium, rhodium, palladium, osmium, and platinum. In: Bingham E, Cohns B, Powell CH, eds. *Patty's toxicology*. 5th ed. New York: John Wiley & Sons; 2000. p. 289-380.

- Dang HS, Pullat VR, Pillai KC. Determining the normal concentration of uranium in urine and application of the data to its biokinetics. *Health Phys* 1992;62:562-6.
- Das YT, Taskar PK, Brown HD, Chattopadhyay SK. Exposure of professional pest control operator to dichlorvos (DDVP) and residue on house structures. *Toxicol Lett* 1983;17(1-2):95-9.
- Davies JE, Peterson JC. Surveillance of occupational, accidental, and incidental exposure to organophosphate pesticides using urine alkyl phosphate and phenolic metabolite measurements. *Ann NY Acad Sci* 1997;837:257-68.
- Davies JE, Freed VH, Enos HF, Duncan RC, Barquet A, Morgade C, et al. Reduction of pesticide exposure with protective clothing for applicators and mixers. *J Occup Med* 1982;24(6):464-68.
- Davison AG, Fayers PM, Taylor AJ, Venables KM, Darbyshire J, Pickering CA, et al. Cadmium fume inhalation and emphysema. *Lancet* 1988 Mar 26;1(8587):663-7.
- De Boeck M, Kirsch-Volders M, Lison D. Cobalt and antimony: genotoxicity and carcinogenicity. *Mutation Res* 2003;533:135-52.
- Dejmek J, Solansky I, Benes I, Lenicek J, Sram RJ. The impact of polycyclic aromatic hydrocarbons and fine particles on pregnancy outcome. *Environ Health Perspect* 2000;108(12):1159-64.
- den Tonkelaar I, Keinan-Boker L, Van't Veer P, Arts CJM, Adlercreutz H, Thijssen JHH, et al. Urinary phytoestrogens and postmenopausal breast cancer risk. *Cancer Epidemiol Biomarkers and Prev* 2001;10:223-8.
- Dernehl CU, Stead FM, Nau CA. Arsine, stibine, and hydrogen sulfide. *Industrial Medicine* 1944;13:361-2.
- Deutsche Forschungsgemeinschaft (DFG). List of MAK and BAT values, 2000. Report No. 35, Commission for the Investigation of Health Hazards of Chemical Compounds in the work area. Weinheim (Germany): Wiley-VCH; 2000.
- Dewailly E, Ayotte P, Bruneau S, Lebel G, Levallois P, Weber JP. Exposure of the Inuit population of Nunavik (Arctic Quebec) to lead and mercury. *Arch Environ Health* 2001;56(4):350-7.
- Diamond GL, Thayer WC, Choudhury H. Pharmacokinetic/pharmacodynamics (PK/PD) modeling of risks of kidney toxicity from exposure to cadmium: estimates of dietary risks in the U.S. population. *J Toxicol Environ Health* 2003;66(Pt A):2141-64.
- Dickel H, Radulescu M, Weyher I, Diepgen TL. Occupationally-induced "isolated cobalt sensitization." *Contact Dermatitis* 2001;45:246-7.
- Dirven HA, van der Broek PH, Jongeneelen FJ. Determination of four metabolites of the plasticizer di (2-ethylhexyl) phthalate in human urine samples. *Int Arch Occup Environ Health* 1993;64(8):555-60.
- Dixon RA, Ferreira D. Genistein. *Phytochemistry* 2002;60(3):205-11.
- Dorgan JF, Brock JW, Rothman N, Needham LL, Miller R, Stephenson HE Jr, et al. Serum organochlorine pesticides and PCBs and breast cancer risk: results from a prospective analysis (USA). *Cancer Causes Control* 1999;10(1):1-11.
- Draper WM, Street JC. Applicator exposure to 2,4-D, dicamba, and a dicamba isomer. *J Environ Sci Health B* 1982;17(4):321-39.
- Droste JHJ, Weyler JJ, Van Meerbeeck JP, Vermeire PA, van Sprundel MP. Occupational risk factors of lung cancer: a hospital based case-control study. *Occup Environ Med* 1999;56:322-7.
- Dua VK, Pant CS, Sharma VP. Determination of levels of HCH and DDT in soil, water and whole blood from bioenvironmental and insecticide-sprayed areas of malaria control. *Indian J Malariol* 1996;33(1):7-15.
- Duggan RE, Corneliussen PE. Dietary intake of pesticide chemicals in the United States. 3. June 1968–April 1970. *Pesticide Monitoring Journal* 1972;5(4):331-41.
- Durakovic A, Horan P, Dietz LA, Zimmerman I. Estimate of the time zero lung burden of depleted uranium in Persian Gulf War veterans by the 24-hour urinary excretion and exponential decay analysis. *Mil Med* 2003;168(8):600-5.
- Durham WF, Armstrong JF, Quinby GE. DDT and DDE content of complete prepared meals. *Arch Environ Health* 1965;11(5):641-7.
- Duty SM, Silva MJ, Barr DB, Brock JW, Ryan L, Chen ZY, Herrick RF, Christiani DC, Hauser R. Phthalate exposure and human semen parameters. *Epidemiology* 2003;14:269-277
- Duty SM, Calafat AM, Silva MJ, Brock JW, Ryan L, Chen Z, et al. The relationship between environmental exposure to phthalates and computer-aided sperm analysis motion parameters. *J Androl* 2004;25(2):293-302.
- Duty SM, Calafat AM, Silva MJ, Ryan L, Hauser R. Phthalate exposure and reproductive hormones in adult men. *Hum Reprod* 2005;20(3):604-10.
- Eisner MD. Environmental tobacco smoke exposure and pulmonary function among adults in NHANES III: impact on the general population and adults with current asthma. *Environ Health Perspect* 2002;110:765-70.

- Ejnik JW, Carmichael AJ, Hamilton MM, McDiarmid M, Squibb K, Boyd P, et al. Determination of the isotopic composition of uranium in urine by inductively coupled plasma mass spectrometry. *Health Phys* 2000;78:143-6.
- Elinder CG, Friberg L. Antimony. In: Friberg L, Nordberg GF, Vouk VB, eds. *Handbook on the toxicology of metals*. 2nd ed. New York: Elsevier; 1986. p. 26-42.
- Ensslin AS, Huber R, Pethran A, Rommelt H, Schierl R, Kulka U, et al. Biological monitoring of hospital pharmacy personnel occupationally exposed to cytostatic drugs: urinary excretion and cytogenetic studies. *Int Arch Occup Environ Health* 1997;70(3):205-8.
- Eriksson P, Archer T, Fredriksson A. Altered behaviour in adult mice exposed to a single low dose of DDT and its fatty acid conjugate as neonates. *Brain Res* 1990;514(1):141-2.
- Eskenazi B, Mocarelli P, Warner M, Samuels S, Vercellini P, Olive D, et al. Serum dioxin concentrations and endometriosis: a cohort study in Seveso, Italy. *Environ Health Perspect* 2002;110(7):629-34.
- Eskenazi B, Harley K, Bradman A, Weltzien E, Jewell NP, Barr DB, et al. Association of in utero organophosphate pesticide exposure and fetal growth and length of gestation in an agricultural population. *Environ Health Perspect* 2004;112(10):1116-24.
- Esteban E, Rubin C, Hill R, Olson D, Pearce K. Association between indoor residential contamination with methyl parathion and urinary para-nitrophenol. *J Expo Anal Environ Epidemiol* 1996;6(3):375-87.
- European Commission. Scientific Committee for Toxicity, Ecotoxicity and the Environment (CSTEE). Opinion on the results of a second risk assessment of: Bis(2-ethylhexyl) phthalate (DEHP) human health part [online]. 2004. Available from URL: http://europa.eu.int/comm/health/ph_risk/committees/sct/documents/out214_en.pdf, 03/17/05.
- Ewers U, Krause C, Schulz C, Wilhelm M. Reference values and human biological monitoring values for environmental toxins. Report on the work and recommendations of the Commission on Human Biological Monitoring of the German Federal Environmental Agency. *Int Arch Occup Environ Health* 1999;72(4):255-60.
- Ezaki T, Tsukahara T, Moriguchi J, Furuki K, et al. No clear-cut evidence for cadmium-induced renal tubular dysfunction among over 10,000 women in the Japanese general population: a nationwide large-scale survey. *Int Arch Occup Environ Health* 2003;76:186-96.
- Fabro S, McLachlan JA, Dames NM. Chemical exposure of embryos during the preimplantation stages of pregnancy: mortality rate and intrauterine development. *Am J Obstet Gynecol* 1984;148(7):929-38.
- Factor-Litvak P, Hasselgren G, Jacobs D, Begg M, Kline J, Geier J, et al. Mercury derived from dental amalgams and neuropsychologic function. *Environ Health Perspect* 2003;111:719-23.
- Falk C, Hanrahan L, Anderson HA, Kanarek MS, Draheim L, Needham LL. Body burden levels of dioxin, furans, and PCBs among frequent consumers of Great Lakes sport fish. The Great Lakes Consortium. *Environ Res* 1999;80(2 Pt 2):S19-S25.
- Farago ME, Kavanagh P, Blanks R, Kelly J, Kazantzis G, Thornton I, et al. Platinum concentrations in urban road dust and soil, and in blood and urine in the United Kingdom. *Analyst* 1998;123(3):451-4.
- Fenske RA, Lu C, Barr D, Needham L. Children's exposure to chlorpyrifos and parathion in an agricultural community in central Washington State. *Environ Health Perspect* 2002;110(5):549-53.
- Fernandez MA, Sanz P, Palomar M, Serra J, Gadea E. Fatal chemical pneumonitis due to cadmium fumes. *Occup Med* 1996;46:372-4.
- Fierens S, Mairesse H, Heilier JF, De Burbure C, Focant JF, Eppe G, et al. Dioxin/polychlorinated biphenyl body burden, diabetes and endometriosis: findings in a population-based study in Belgium. *Biomarkers* 2003;8(6):529-34.
- Finklea J, Priester LE, Creason JP, Hauser T, Hinnert T, Hammer DI. Polychlorinated biphenyl residues in human plasma expose a major urban pollution problem. *Am J Public Health* 1972;62(5):645-51.
- Frank R, Campbell RA, Sirons G J. Forestry workers involved in aerial application of 2,4-dichlorophenoxyacetic acid (2,4-D): exposure and urinary excretion. *Arch Environ Contam Toxicol* 1985;4:427-35.
- Franklin CA, Fenske RA, Greenhalgh R, Mathieu L, Denley HV, Leffingwell JT, et al. Correlation of urinary pesticide metabolite excretion with estimated dermal contact in the course of occupational exposure to Guthion. *J Toxicol Environ Health* 1981;7(5):715-31.
- Galletti M, D'Annibale L, Pinto V, Cremisini C. Uranium daily intake and urinary excretion: a preliminary study in Italy. *Health Phys* 2003;85:228-35.
- Garfitt SJ, Jones K, Mason HJ, Cocker J. Exposure to the organophosphate diazinon: data from a human volunteer study with oral and dermal doses. *Toxicol Lett* 2002;134:105-13.
- Geltman PL, Brown MJ, Cochran J. Lead poisoning among refugee children resettled in Massachusetts, 1995 to 1999. *Pediatrics* 2001;108(1):158-62.
- Genter MB. Magnesium, calcium, strontium, barium, and radium In: Bingham A, Cohressen B, Powell C, eds. *Patty's*

- toxicology. 5th ed. New York: John Wiley & Sons, Inc.; 2001. p. 221-52
- George DA. Permethrin and its two metabolite residues in seven agricultural crops. *J AOAC* 1985;68(6):1160-3.
- Geyer HJ, Schramm KW, Feicht EA, Behechti A, Steinberg C, Bruggemann R, et al. Half-lives of tetra-, penta-, hexa-, hepta-, and octachlorodibenzo-p-dioxin in rats, monkeys, and humans—a critical review. *Chemosphere* 2002;48(6):631-44.
- Glynn AW, Granath F, Aune M, Atuma S, Darnerud PO, Bjerselius R, et al. Organochlorines in Swedish women: determinants of serum concentrations. *Environ Health Perspect* 2003;111:349-55.
- Glynn AW, Wolk A, Aune M, Atuma S, Zettermark S, Maehle-Schmid M, et al. Serum concentrations of organochlorines in men: a search for markers of exposure. *Sci Total Environ* 2000; 263(1-3):197-208.
- Goen T, Gundel J, Schaller KH, Angerer J. The elimination of 1-hydroxypyrene in the urine of the general population and workers with different occupational exposures to PAH. *Sci Total Environ* 1995;163(1-3):195-201.
- Goldberg MA, Dunning SP, Bunn HF. Regulation of the erythropoietin gene: evidence that the oxygen sensor is a heme protein. *Science* 1988;242:1412-5.
- Grace PB, Taylor JI, Low Y, Luben RN, Mulligan AA, Botting NP, et al. Phytoestrogen concentrations in serum and spot urine as biomarkers for dietary phytoestrogen intake and their relation to breast cancer risk in European prospective investigation of cancer and nutrition—Norfolk. *Cancer Epidemiol Biomarkers and Prev* 2004;13:698-708.
- Grandjean P, Weihe P, Needham LL, Burse VW, Patterson DG Jr, Sampson EJ, et al. Relation of a seafood diet to mercury, selenium, arsenic, and polychlorinated biphenyl and other organochlorine concentrations in human milk. *Environ Res* 1995;71(1):29-38.
- Grether JK, Harris JA, Neutra R, Kizer KW: Exposure to aerial malathion application and the occurrence of congenital anomalies and low birthweight. *Am J Public Health* 1987;77:1009-10.
- Grimalt JO, Sunyer J, Moreno V, Amaral OC, Sala M, Rosell A, et al. Risk excess of soft-tissue sarcoma and thyroid cancer in a community exposed to airborne organochlorinated compound mixtures with a high hexachlorobenzene content. *Int J Cancer* 1994;56(2):200-3.
- Guallar E, Sanz-Gallardo MI, van't Veer P, Bode P, Aro A, Gomez-Aracena J, et al. Mercury, fish oils, and the risk of myocardial infarction. *New Engl J Med* 2002;347(22):1747-54.
- Gundel J, Mannschreck C, Buttner K, Ewers U, Angerer J. Urinary levels of 1-hydroxypyrene, 1-, 2-, 3-, and 4-hydroxyphenanthrene in females living in an industrial area of Germany. *Arch Environ Contam Toxicol* 1996;31(4):585-90.
- Gundel J, Schaller KH, Angerer J. Occupational exposure to polycyclic aromatic hydrocarbons in a fireproof stone producing plant: biological monitoring of 1-hydroxypyrene, 1-, 2-, 3- and 4-hydroxyphenanthrene, 3-hydroxybenz(a)anthracene and 3-hydroxybenzo-(a)-pyrene. *Int Arch Occup Environ Health* 2000;73(4):270-4.
- Gwiazda RH, Squibb K, McDiarmid M, Smith D. Detection of depleted uranium in urine of veterans from the 1991 Gulf War. *Health Phys* 2004;86:12-8.
- Halperin W, Vogt R, Sweeney MH, Shopp G, Fingerhut M, Petersen M. Immunological markers among workers exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Occup Environ Med* 1998;55(11):742-9.
- Hamilton EI, Sabbioni E, Van der Venne MT. Element reference values in tissues from inhabitants of the European community. VI. Review of elements in blood, plasma and urine and a critical evaluation of reference values for the United Kingdom population. *Sci Total Environ* 1994;158:165-90.
- Hanrahan LP, Falk C, Anderson HA, Draheim L, Kanarek MS, Olson J. Serum PCB and DDE levels of frequent Great Lakes sport fish consumers—a first look. *The Great Lakes Consortium. Environ Res* 1999;80(2 Pt 2):S26-S37.
- Hansen AM, Wallin H, Binderup ML, Dybdahl M, Autrup H, Loft S, et al. Urinary 1-hydroxypyrene and mutagenicity in bus drivers and mail carriers exposed to urban air pollution in Denmark. *Mutat Res* 2004;557(1):7-17.
- Hardt J, Angerer J. Gas chromatographic method with mass-selective detection for the determination of 2-isopropoxyphenol in human urine. *J Chromatogr B* 1999;723(1&2):139-45.
- Harris CA, Henttu P, Park MG, Sumpter JP. The estrogenic activity of phthalate esters in vitro. *Environ Health Perspect* 1997;105:105-8.
- Hattula T, Rahola T. The distribution and biological half-time of ²⁰³Hg in the human body according to a modified whole-body counting technique. *Environmental Physiology and Biochemistry* 1975;5:252-7.
- Hauser R, Meeker JD, Park S, Silva MJ, Calafat AM. Temporal variability of urinary phthalate metabolite levels in men of reproductive age [published erratum appears in *Environ Health Perspect* 2004;112(17):1740]. *Environ Health Perspect* 2004;112(17):1734-40.
- Hayes TB, Collins A, Lee M, Mendoza M, Noriega N, Stuart AA, et al. Hermaphroditic, demasculinized frogs after

- exposure to the herbicide atrazine at low ecologically relevant doses. Proceedings of the National Academy of Sciences USA Washington, DC: PNAS; 2002 Apr 16; 99(8):5476-80.
- Hayes WJ Jr. Mortality in 1969 from pesticides, including aerosols. Arch Environ Health 1976;31:61-72.
- Hedge AG, Thakker DM, Bhat IS. Long-term clearance of inhaled ⁶⁰Co. Health Phys 1979;36:732-4.
- Hellstrom L, Elinder CG, Dahlberg B, Lundberg M, et al. Cadmium exposure and end-stage renal disease. Am J Kidney Dis 2001;38:1001-8.
- Helzlsouer KJ, Alberg AJ, Huang HY, Hoffman SC, Strickland PT, Brock JW, et al. Serum concentrations of organochlorine compounds and the subsequent development of breast cancer. Cancer Epidemiol Biomarkers Prev 1999;8(6):525-32.
- Henson MC, Chedrese PJ. Endocrine disruption by cadmium, a common environmental toxicant with paradoxical effects on reproduction. Experimental Biology and Medicine 2004;229:383-92.
- Hernandez-Valero MA, Bondy ML, Spitz MR, Zahm SH. Evaluation of Mexican-American migrant farmworker work practices and organochlorine pesticide metabolites. Am J Ind Med 2001;40(5):554-60.
- Herr CE, Jankofsky M, Angerer J, Kuster W, Stilianakis NI, Gieler U, et al. Influences on human internal exposure to environmental platinum. J Expo Anal Environ Epidemiol 2003;13(1):24-30.
- Herrero C, Ozalla D, Sala M, Otero R, Santiago-Silva M, Lecha M, et al. Urinary porphyrin excretion in a human population highly exposed to hexachlorobenzene. Arch Dermatol 1999;135(4):400-4.
- Heudorf U, Angerer J. Internal exposure to PAHs of children and adults living in homes with parquet flooring containing high levels of PAHs in the parquet glue. Int Arch Occup Environ Health 2001a;74(2):91-101.
- Heudorf U, Angerer J. Metabolites of organophosphorous insecticides in urine specimens from inhabitants of a residential area. Environ Res 2001c; 86:80-7.
- Heudorf U, Angerer J. Metabolites of pyrethroid insecticides in urine specimens: current exposure in an urban population in Germany. Environ Health Perspect 2001d;109(3):213-7.
- Heudorf U, Angerer J. Urinary monohydroxylated phenanthrenes and hydroxypyrene--the effects of smoking habits and changes induced by smoking on monooxygenase-mediated metabolism. Int Arch Occup Environ Health. 2001b 74(3):177-83.
- Heudorf U, Angerer J, Drexler H. Current internal exposure to pesticides in children and adolescents in Germany: urinary levels of metabolites of pyrethroid and organophosphorus insecticides. Int Arch Occup Environ Health. 2004;77(1):67-72.
- Heudorf U, Angerer J, Drexler H. Polychlorinated biphenyls in the blood plasma: current exposure of the population in Germany. Rev Environ Health 2002;17(2):123-34.
- Hill RH Jr, Head SL, Baker S, Gregg M, Shealy DB, Bailey SL, et al. Pesticide residues in urine of adults living in the United States: reference range concentrations. Environ Res 1995;71:99-108.
- Hill RH Jr, To T, Holler JS, Fast DM, Smith SJ, Needham LL, et al. Residues of chlorinated phenols and phenoxy acid herbicides in the urine of Arkansas children. Arch Environ Contam Toxicol 1989;18(4):469-74.
- Hines CJ, Deddens JA. Determinants of chlorpyrifos exposures and urinary 3,5,6-trichloro-2-pyridinol levels among termiticide applicators. Ann Occup Hyg 2001;45:309-21.
- Hines CJ, Deddens JA, Striley CA, Biagini RE, Shoemaker DA, Brown KK, et al. Biological monitoring for selected herbicide biomarkers in the urine of exposed custom applications: application of mixed-effect models. Ann Occup Hyg 2003;47(6):503-17.
- Hodgert JH, Zacharewski TR, Hammond GL. Interactions between human plasma sex hormone-binding globulin and xenobiotic ligands. J Steroid Biochem Mol Biol 2000;75(2-3):167-76.
- Hoppin JA, Brock JW, Davis BJ, Baird DD. Reproducibility of urinary phthalate metabolites in first morning urine samples. Environ Health Perspect 2000;110(5):515-8.
- Horiguchi H, Oguma E, Sasaki S, Miyamoto K, Ikeda Y, Machida M, et al. Comprehensive study of the effects of age, iron deficiency, diabetes mellitus, and cadmium burden on dietary cadmium absorption in cadmium-exposed female Japanese farmers. Toxicol Appl Pharmacol 2004a; 196:114-23.
- Horiguchi H, Oguma E, Sasaki S, Miyamoto K, Ikeda, Yoko ; Machida, Munehito et al. Dietary exposure to cadmium at close to the current provisional tolerable weekly intake does not affect renal function among female Japanese farmers. Environ Res 2004b;95:20-31.
- Horn-Ross PL, Barnes S, Kirk M, Coward L, Parsonnet J, Hiatt RA. Urinary phytoestrogen levels in young women from a multiethnic population. Cancer Epidemiol Biomarkers Prev 1997;6:339-45.
- Hotz P, Buchet JP, Bernard A, Lison D, Lauwerys R. Renal effects of low-level environmental cadmium exposure: 5-year follow-up of a subcohort from the Cadmibel study. Lancet 1999;354:1508-13.

- Howe GR, Hirohata T, Hislop TG, Iscovich JM, Yuan JM, Katsouyanni K, et al. Dietary factors and risk of breast cancer: combined analysis of 12 case-control studies. *J Natl Cancer Inst* 1990;82:561-9.
- Hoyer AP, Grandjean P, Jorgensen T, Brock JW, Hartvig HB. Organochlorine exposure and risk of breast cancer. *Lancet* 1998;352(9143):1816-20.
- Hu H, Aro A, Payton M, Korrick S, Sparrow D, Weiss ST, et al. The relationship of bone and blood lead to hypertension. *JAMA* 1996;275(15):1171-6.
- Hunter DJ, Hankinson SE, Laden F, Colditz GA, Manson JE, Willett WC, et al. Plasma organochlorine levels and the risk of breast cancer. *N Engl J Med* 1997;337(18):1253-8.
- Hursh JB, Greenwood MR, Clarkson TW, Allen J, Demuth S. The effect of ethanol on the fate of mercury vapor inhaled by man. *J Pharmacol Exp Ther* 1980;214(3):520-7.
- Hutchins AM, Lampe JW, Martini MC, Campbell DR, Slavin JL. Vegetables, fruits and legumes: effect on isoflavonoid phytoestrogen and lignan excretion. *J Am Diet Assoc* 1995a;95:769-74.
- Hutchins AM, Slavin JL, Lampe JW. Urinary isoflavonoid phytoestrogen and lignan excretion after consumption of fermented and unfermented soy products. *J Am Diet Assoc* 1995b;95(5):545-51.
- Iavicoli I, Bocca B, Petrucci F, Senofonte O, Carelli G, Alimonti A, et al. Biomonitoring of traffic police officers exposed to airborne platinum. *Occup Environ Med* 2004 Jul;61(7):636-9.
- Iavicoli I, Caroli S, Alimonti A, Petrucci F, Carelli G. Biomonitoring of a worker population exposed to low antimony trioxide levels. *J Trace Elem Med Biol* 2002;16: 33-9.
- Ichikawa Y, Kusaka Y, Goto S. Biological monitoring of cobalt exposure, based on cobalt concentrations in blood and urine. *Int Arch Occup Environ Health* 1985;55(4):269-76.
- Immerman FW, Schaum JL. Nonoccupational Pesticide Exposure Study (NOPES). Report.[EPA/600/3-90/003; Order No. PB90-152224], U.S. Environmental Protection Agency: Washington (DC); 1990. p. 256.
- Ingram D, Sanders K, Kolybaba M, Lopez D. Case-control study of phyto-oestrogens and breast cancer. *Lancet* 1997;350(9083):990-4.
- Institute of Medicine. Board on Health Promotion and Disease Prevention. Veterans and Agent Orange: update 2002. Washington (DC): National Academy Press; 2003.
- Institute of Medicine. Products for tobacco exposure reduction. In: Clearing the smoke: assessing the science base for tobacco harm reduction. Washington (DC): National Academy Press; 2001b. p. 99.
- Institute of Medicine. Standing Committee on the Scientific Evaluation of Dietary Reference Intakes, Food and Nutrition Board. Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc: a report of the Panel on Micronutrients. Washington, (DC): National Academy Press; 2001a.
- International Agency for Research on Cancer (IARC). IARC Monographs. [online]. Available from URL: <http://www.iarc.fr>, 03/17/05.
- International Programme on Chemical Safety. Environmental health criteria monographs. Carbamate pesticides: a general introduction. Vol. 64. Geneva: World Health Organization; 1986. p. 23-43.
- International Programme on Chemical Safety. Environmental health criteria 97: deltamethrin. Section 1.1.4. [online] 1990. Available from URL: <http://www.inchem.org/documents/ehc/ehc/ehc97.htm#SubSectionNumber:1.1.4>, 02/7/05.
- Iversen BS, Menne C, White MA, Kristiansen J, Christensen JM, Sabbioni E. Inductively coupled plasma mass spectrometric determination of molybdenum in urine from a Danish population. *Analyst* 1998;123(1):81-5.
- Iwase A, Aiba M, Kira S. Respiratory nicotine absorption in non-smoking females during passive smoking. *Int Arch Occup Environ Health* 1991;63:139-43.
- Jacob J, Seidel A. Biomonitoring of polycyclic aromatic hydrocarbons in human urine. *J Chromatogr B* 2002;778(1-2):31-47.
- Jacobson JL, Jacobson SW. Intellectual impairment in children exposed to polychlorinated biphenyls in utero. *N Engl J Med* 1996;335:783-9.
- James RA, Hertz-Picciotto I, Willman E, Keller JA, Charles MJ. Determinants of serum polychlorinated biphenyls and organochlorine pesticides measured in women from the child health and development study cohort, 1963-1967. *Environ Health Perspect* 2002;110:617-24.
- Jarfelt K, Dalgaard M, Hass U, Borch J, Jacobsen H, Ladefoged O. Antiandrogenic effects in male rats perinatally exposed to a mixture of di(2-ethylhexyl) phthalate and di(2-ethylhexyl) adipate. *Reprod Toxicol* 2005;19(4):505-15.
- Jarup L, Hellstrom L, Alfven T, Carlsson MD, Grubb A, Persson B, et al. Low level exposure to cadmium and early kidney damage: the OSCAR study [published erratum appears in *Occup Environ Med* 2002;59:497]. *Occup Environ Med* 2000;57:668-72.

- Jarvis JQ, Hammond E, Meier R, Robinson C. Cobalt cardiomyopathy. A report of two cases from mineral assay laboratories and a review of the literature. *Journal of Occupational Medicine* 1992;34:620-6.
- Jauhiainen A, Kangas J, Laitinen S, Savolainen K. Biological monitoring of workers exposed to mevinphos in greenhouses. *Bull Environ Contam Toxicol* 1992;49(1):37-43.
- Jefferson JA, Escudero E, Hurtado ME, Pando J, Tapia R, Swenson ER, et al. Excessive erythrocytosis, chronic mountain sickness, and serum cobalt levels. *Lancet* 2002;359:407-8.
- Jemal A, Graubard BI, Devesa SS, Flegal KM. The association of blood lead level and cancer mortality among whites in the United States. *Environ Health Perspect* 2002;110(4):325-9.
- Jin T, Nordberg G, Ye T, Bo M, Wang H, Zhu G, et al. Osteoporosis and renal dysfunction in a general population exposed to cadmium in China. *Environ Res* 2004;96:353-9.
- Joblin S, Reynolds T, White R, Parker MG, Sumpter JP. A variety of environmentally persistent chemicals including some phthalate plasticizers are weakly estrogenic. *Environ Health Perspect* 1995;103:582-7.
- Johnson E, Shorter C, Bestervelt L, Patterson D, Needham L, Piper W, et al. Serum hormone levels in humans with low serum concentrations of 2,3,7,8-TCDD. *Toxicol Ind Health* 2001;17(4):105-12.
- Johnson GD, Harsy SG, Geronimo J, Wise JM. Ortho-phenylphenol and phenylhydroquinone residues in citrus fruit and processed citrus products after postharvest fungicidal treatments with sodium orthophenylphenate in California and Florida. *J Agric Food Chem* 2001;49(5):2497-502.
- Johnson MD, Kenney N, Stoica A, Hilkivi-Clarke L, Singh B, Chepko G, et al. Cadmium mimics the in vivo effects of estrogen in the uterus and mammary gland. *Nature Med* 2003;9:1081-4.
- Jones RD, Winter DP, Cooper AJ. Absorption study of pentachlorophenol in persons working with wood preservatives. *Hum Toxicol* 1986;5(3):189-94.
- Jongeneelen FJ. Biological monitoring of environmental exposure to polycyclic aromatic hydrocarbons; 1-hydroxypyrene in urine of people. *Toxicol Lett* 1994;72(1-3):205-11.
- Jonsson F, Sandborgh-Englund G, Johanson G. A compartmental model for the kinetics of mercury vapor in humans. *Toxicol Appl Pharmacol* 1999;155(2):161-8.
- Jonsson HT Jr, Keil JE, Gaddy RG, Loadholt CB, Hennigar GR, Walker EM Jr. Prolonged ingestion of commercial DDT and PCB; effects on progesterone levels and reproduction in the mature female rat. *Arch Environ Contam Toxicol* 1975-1976;3(4):479-90.
- Jourdan S, Moreno B, Pellegrino S, Petrarulo M, Marco R. Suicidal poisoning with barium chloride. *Forensic Sci Int* 2001;119:263-5.
- Jung D, Berg PA, Edler L, Ehrental W, Fenner D, Flesch-Janys D, et al. Immunologic findings in workers formerly exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin and its congeners. *Environ Health Perspect* 1998;106(Suppl 2):689-95.
- Juhler RK, Larsen SB, Myer O, Jensen ND, Spanò M, Giwercman A, et al. Human semen quality in relation to dietary pesticide exposure and organic diet. *Arch Environ Contam Toxicol* 1999;37:415-23.
- Kaiser R, Henderson AK, Daley WR, Naughton M, Khan MH, Rahman M, et al. Blood lead levels of primary school children in Dhaka, Bangladesh. *Environ Health Perspect* 2001;109(6):563-6.
- Kang D, Tepper A, Patterson DG Jr. Coplanar PCBs and the relative contribution of coplanar PCBs, PCDDs, and PCDFs to the total 2,3,7,8-TCDD toxicity equivalents in human serum. *Chemosphere* 1997;35(3):503-11.
- Kang JW, Cho SH, Kim H, Lee CH. Correlation of urinary 1-hydroxypyrene and 2-naphthol with total suspended particulates in ambient air in municipal middle-school students in Korea. *Arch Environ Health* 2002;57(4):377-82.
- Kanoh T, Fukuda M, Onozuka H, Kinouchi T, Ohnishi Y. Urinary 1-hydroxypyrene as a marker of exposure to polycyclic aromatic hydrocarbons in environment. *Environ Res* 1993;62(2):230-41.
- Karpas Z, Halicz L, Roiz J, Marko R, Katorza E, Lorber A, et al. Inductively coupled plasma mass spectrometry as a simple, rapid, and inexpensive method for determination of uranium in urine and fresh water: comparison with LIF. *Health Phys* 1996;71(6):879-85.
- Karpas Z, Lorber A, Elish E, Kol R, Roiz Y, Marko R, et al. Uptake of ingested uranium after low "acute intake." *Health Phys* 1998;74(3):337-45.
- Karr SC, Lampe JW, Hutchins AM, Slavin JL. Urinary isoflavonoid excretion in humans is dose dependent at low to moderate levels of soy-protein consumption. *Am J Clin Nutr* 1997;66:46-51.
- Kato K, Silva MJ, Reidy JA, Hurtz D, Malek NA, Needham LL, et al. Mono(2-ethyl-5-hydroxyhexyl) phthalate and mono(2-ethyl-5-oxohexyl) phthalate as biomarkers for human exposure assessment to di-(2-ethylhexyl) phthalate. *Environ Health Perspect* 2004;112(3):327-30.
- Kentner M, Leinemann M, Schaller KH, Weltle D, Lenert G. External and internal antimony exposure in starter battery production. *Int Arch Occup Environ Health* 1995;67:119-23.

- Kerper LE, Ballatori N, Clarkson TW. Methyl mercury transport across the blood-brain barrier by an amino acid carrier. *Am J Physiol* 1992;262:RF 761-RF 5.
- Kessler W, Numtip W, Grote K, Csanády G, Chahoud I, Filser J (2004). Blood burden of di(2-ethylhexyl)phthalate (DEHP) and its primary metabolite mono(2-ethylhexyl) phthalate (MEHP) in pregnant and non-pregnant rats and marmosets. *Toxicol Appl Pharmacol* 2004;195:142-53.
- Kikuchi Y, Nomiyama T, Kumagai N, Dekio F, Uemura T, Takebayashi T, et al. Uptake of cadmium in meals from the digestive tract of young non-smoking Japanese female volunteers. *J Occup Health* 2003;45:43-52.
- Kim MK, Chung BC, Yu VY, Nam JH, Lee HC, Huh KB, et al. Relationships of urinary phyto-oestrogen excretion to BMD in postmenopausal women. *Clin Endocrinol (Oxf)* 2002;56(3):321-8.
- Kim R, Rotnitzky A, Sparrow D, Weiss ST, Wager C, Hu H. A longitudinal study of low-level lead exposure and impairment of renal function: the Normative Aging Study. *JAMA* 1996;275:1177-81.
- Kim YD, Lee CH, Nan HM, Kang JW, Kim H. Effects of genetic polymorphisms in metabolic enzymes on the relationships between 8-hydroxydeoxyguanosine levels in human leukocytes and urinary 1-hydroxypyrene and 2-naphthol concentrations. *J Occup Health* 2003;45(3):160-7.
- Kingman A, Albertini T, Brown LJ. Mercury concentrations in urine and whole blood associated with amalgam exposure in a U.S. military population. *J Dent Res* 1998;77:461-71.
- Kirkman LM, Lampe JW, Campbell D, Martini M, Slavin J. Urinary lignan and isoflavonoid excretion in men and women consuming vegetable and soy diets. *Nutr Cancer* 1995;24:1-12.
- Kisker C, Schindelin H, Rees DC. Molybdenum-cofactor-containing enzymes: structure and mechanism. *Annual Review of Biochemistry* 1997;66:233-67.
- Kitamura K, Kikuchi Y, Watanabe S, Waechter G, Sakurai H, Takada T. Health effects of chronic exposure to polychlorinated dibenzo-P-dioxins (PCDD), dibenzofurans (PCDF) and coplanar PCB of municipal waste incinerator workers. *J Epidemiol* 2000;10(4):262-70.
- Klaassen CD, Liu J, Choudhuri S. Metallothionein: an intracellular protein to protect against cadmium toxicity. *Annual Review of Pharmacology and Toxicology* 1999;39:267-94.
- Klein KO. Isoflavones, soy-based infant formulas, and relevance to endocrine function. *Nutr Rev* 1998;56:193-204.
- Knowles LM, Zigrossi DA, Tauber RA, Hightower C, Milner JA. Flavonoids suppress androgen-independent human prostate tumor proliferation. *Nutr Cancer* 2000;38(1):116-22.
- Koch HM, Drexler H, Angerer J. Internal exposure of nursery-school children and their parents and teachers to di(2-ethylhexyl)phthalate (DEHP). *International Journal of Hygiene and Environmental Health* 2004;207(1):15-22.
- Koch HM, Hardt J, Angerer J. Biological monitoring of exposure of the general population to the organophosphorus pesticides chlorpyrifos and chlorpyrifos-methyl by determination of their specific metabolite 3,5,6-trichloro-2-pyridinol. *International Journal of Hygiene and Environmental Health* 2001;204:175-80.
- Koch HM, Rossbach B, Drexler H, Angerer J. Internal exposure of the general population to DEHP and other phthalates - determination of secondary and primary phthalate monoester metabolites in urine. *Environ Res* 2003;93:177-85.
- Kohli JD, Khanna RN, Gupta BN, Dhar MM, Tandon JS, Sircar KP. Absorption and excretion of 2,4-dichlorophenoxyacetic acid in man. *Xenobiotica* 1974b;4:97-100.
- Kogevinas M. Human health effects of dioxins: cancer, reproductive and endocrine system effects. *Hum Reprod Update* 2001;7(3):331-9.
- Kohli JD, Khanna RN, Gupta BN, Dhar MM, Tandon JS, Sircar KP. Absorption and excretion of 2,4,5-trichlorophenoxy acetic acid in man. *Archives of International Pharmacodynamic Therapeutics* 1974a; 210:250-5.
- Kolmodin-Hedman B, Erne K. Estimation of occupational exposure to phenoxy acids (2,4-D and 2,4,5-T). *Arch Toxicol Suppl* 1980;4:318-21.
- Komaromy-Hiller G, Ash KO, Costa R, Howerton K. Comparison of representative ranges based on U.S. patient population and literature reference intervals for urinary trace elements. *Clin Chim Acta* 2000;296(1-2):71-90.
- Koo JW, Parham F, Kohn MC, Masten SA, Brock JW, Needham LL, et al. The association between biomarker-based exposure estimates for phthalates and demographic factors in a human reference population. *Environ Health Perspect* 2002;110(4):405-10.
- Koopman-Esseboom C, Huisman M, Touwen BC, Boersma ER, Brouwer A, Sauer PJ, et al. Newborn infants diagnosed as neurologically abnormal with relation to PCB and dioxin exposure and their thyroid-hormone status. *Dev Med Child Neurol* 1997;39(11):785.
- Korn EL, Graubard BI. Confidence intervals for proportions with small expected number of positive counts estimated from survey data. *Survey Methodology* 1998; 24:193-201.
- Korrick SA, Hunter DJ, Rotnitzky A, Hu H, Speizer FE. Lead and hypertension in a sample of middle-aged women. *Am J Public Health* 1999;89:330-5.

- Kotsopoulos D, Dalais FS, Liang YL, McGrath BP, Teede HJ. The effects of soy protein-containing phytoestrogens on menopausal symptoms in postmenopausal women. *Climacteric* 2000;3(3):161-7.
- Koval'skiy GA, Yarovaya GA, Shmavonyan DM. Changes of purine metabolism in man and animals under conditions of molybdenum biogeochemical provinces. *Zhurnal Obshchey Biologii* 1961;22(3):179-91.
- Kozlowski LT, Mehta NY, Sweeney CT, Schwartz SS, Vogler GP, Jarvis MJ, et al. Filter ventilation and nicotine content of tobacco in cigarettes from Canada, the United Kingdom, and the United States. *Tob Control* 1998;7:369-75.
- Kramer RE, Wellman SE, Rockhold RW, Baker RC. Pharmacokinetics of methyl parathion: a comparison following single intravenous, oral or dermal administration. *Journal of Biomedical Science* 2002;9:311-20.
- Kraus T, Schramel P, Schaller KH, Zobelein P, Weber A, Angerer J. Exposure assessment in the hard metal manufacturing industry with special regard to tungsten and its compounds. *Occup Environ Med* 2001;58(10):631-4.
- Kreijkamp-Kaspers S, Kok L, Grobbee DE, de Haan EHF, Aleman A, Lampe JW, et al. Effect of soy protein containing isoflavones on cognitive function, bone mineral density and plasma lipids in postmenopausal women. *JAMA* 2004;292:65-74.
- Krieger RI, Dinoff TM. Malathion deposition, metabolite clearance, and cholinesterase status of date dusters and harvesters in California. *Arch Environ Contam Toxicol* 2000;38(4):546-63.
- Kriss JP, Carnes WH, Gross RT. Hypothyroidism and thyroid hyperplasia in patients treated with cobalt. *JAMA* 1955;157:117-21.
- Kristiansen J, Christensen JM, Iversen BS, Sabbioni E. Toxic trace element reference levels in blood and urine: influence of gender and lifestyle factors. *Sci Total Environ* 1997;204:147-60.
- Kuhn K, Wieseler B, Leng G, Idel H. Toxicokinetics of pyrethroids in humans: consequences for biological monitoring. *Bull Environ Contam Toxicol* 1999;62:101-8.
- Kuo CT, Chen HW, Chen JL. Determination of 1-hydroxypyrene in children urine using column-switching liquid chromatography and fluorescence detection. *J Chromatogr B* 2004;805(2):187-93.
- Kurtio P, Auvinen A, Salonen L, Saha H, Pekkanen J, Makelainen I, et al. Renal effects of uranium in drinking water. *Environ Health Perspect* 2002;110(4):337-42.
- Kutz FW, Cook BT, Carter-Pokras OD, Brody D, Murphy RS. Selected pesticide residues and metabolites in urine from a survey of the U.S. general population. *J Toxicol Environ Health* 1992;37(2):277-91.
- Kutz FW, Yobs AR, Strassman SC. Racial stratification of organochlorine insecticide residues in human adipose tissue. *Journal of Occupational Medicine* 1977;19(9):619-22.
- Kutz FW, Wood PH, Bottimore DP. Organochlorine pesticides and polychlorinated biphenyls in human adipose tissue. *Reviews of Environmental Contamination and Toxicology* 1991;120:1-82.
- Kuusimaki L, Peltonen Y, Mutanen P, Peltonen K, Savela K. Urinary hydroxy-metabolites of naphthalene, phenanthrene and pyrene as markers of exposure to diesel exhaust. *Int Arch Occup Environ Health* 2004;77(1):23-30.
- Lake BG, Phillips JC, Linnell JC, Gangolli SD. The in vitro hydrolysis of some phthalate diesters by hepatic and intestinal preparations from various species. *Toxicol Appl Pharmacol* 1977;39:239-48.
- Lampe JW, Gustafson DR, Hutchins AM, Martini MG, Li S, Wahala K, et al. Urinary isoflavonoid and lignan excretion on a western diet: relation to soy, vegetable and fruit intake. *Cancer Epidemiol Biomarkers and Prev* 1999;8:699-707.
- Lauwerys RB, Hoet P. Industrial chemical exposure: guidelines for biological monitoring. 3rd ed. Boca Raton (FL): Lewis Publishers; 2001. p. 87-95, 54-68, 500-11.
- Lanphear BP, Dietrich K, Auinger P, Cox C. Cognitive deficits with blood lead concentrations < 10 µg/dL in US children and adolescents. *Public Health Rep* 2000;115:521-9.
- Lebel G, Dodin S, Ayotte P, Marcoux S, Ferron LA, Dewailly E. Organochlorine exposure and the risk of endometriosis. *Fertil Steril* 1998;69(2):221-8.
- Lee HP, Gourley L, Duffy SW, Esteve J, Lee J, Day NE. Dietary effects on breast-cancer risk in Singapore. *Lancet* 1991;337:1197-1200.
- Lee KH, Ichiba M, Zhang J, Tomokuni K, Hong YC, Ha M, et al. Multiple biomarkers study in painters in a shipyard in Korea. *Mutat Res* 2003;540(1):89-98.
- Leggett RW. Age-specific kinetic model of lead metal in humans. *Environ Health Perspect* 1993;101(7):598-616.
- Leng G, Lewalter J. Role of individual susceptibility in risk assessment of pesticides. *Occup Environ Med* 1999;56(7):449-53.
- Leng G, Ranft U, Sugiri D, Hadnagy W, Berger-Preiss, Idel H. Pyrethroids used indoors—biological monitoring of exposure to pyrethroids following an indoor pest control operation. *International Journal of Hygiene and Environmental Health* 2003;206:85-92.

- Lhotka C, Szekeres T, Steffan I, Zhuber K, Zweymuller K. Four-year study of cobalt and chromium blood levels in patients managed with two different metal-on-metal total hip replacements. *J Orthop Res* 2003;21(2):189-95.
- Li N, Hao M, Phalen RF, Hinds WC, Nel AE. Particulate air pollutants and asthma. A paradigm for the role of oxidative stress in PM-induced adverse health effects. *Clin Immunol* 2003;109(3):250-65.
- Life Sciences Research Office (LSRO). Review and analysis of the literature on the potential adverse health effects of dental amalgam. Bethesda (MD): LSRO; July 2004.
- Lilis R, Blumberg WE, Fischbein A, Diamond S, Anderson HA, et al. Lead effects among secondary smelter workers with blood lead levels below 80 µg/100 ml. *Arch Environ Health* 1977;32(6):256-66.
- Linna A, Oksa P, Palmroos P, Roto P, Laippala P, Uitti J. Respiratory health of cobalt production workers. *American Journal of Industrial Medicine* 2003;44:124-32.
- Linnainmaa M, Kiilunen M. Urinary cobalt as a measure of exposure in the wet sharpening of hard metal and stellite blades. *Int Arch Occup Environ Health* 1997;69(3):193-200.
- Lisi P. Co-sensitivity between cobalt and other transition metals. *Contact Dermatitis* 2003;48:172-3.
- Lison D, Buchet JP, Swennen B, Molders J, Lauwerys R. Biological monitoring of workers exposed to cobalt metal, salt, oxides, and hard metal dust. *Occup Environ Med* 1994;51(7):447-50.
- Liss GM, Albro PW, Hartle RW, Stringer WT. Urine phthalate determinations as an index of occupational exposure to phthalic anhydride and di (2-ethylhexyl) phthalate. *Scand J Work Environ Health* 1985;11(5):381-7.
- Lloyd J. Long-term mortality study of steelworkers. V. Respiratory cancer in coke plant workers. *J Occup Med* 1971;13:53-68.
- Longnecker MP, Ryan JJ, Gladen BC, Schecter AJ. Correlations among human plasma levels of dioxin-like compounds and polychlorinated biphenyls (PCBs) and implications for epidemiologic studies. *Arch Environ Health* 2000;55(3):195-200.
- Longnecker MP, Wolff MS, Gladen BC, Brock JW, Grandjean P, Jacobson JL, et al. Comparison of polychlorinated biphenyl levels across studies of human neurodevelopment. *Environ Health Perspect* 2003;111(1):65-70.
- Lorber M. A pharmacokinetic model for estimating exposure of Americans to dioxin-like compounds in the past, present, and future. *Sci Total Environ* 2002;288(1-2):81-95.
- Lovekamp-Swan T, Davis BJ. Mechanisms of phthalate ester toxicity in the female reproductive system. *Environ Health Perspect* 2003;111(2):139-45.
- Lu LJ, Grady JJ, Marshall MV, Ramanujam VM, Anderson KE. Altered time course of urinary daidzein and genistein excretion during chronic soya diet in healthy male subjects. *Nutr Cancer* 1995;24:311-23.
- Lu PL, Chen ML, Mao IF. Urinary 1-hydroxypyrene levels in workers exposed to coke oven emissions at various locations in a coke oven plant. *Arch Environ Health* 2002;57(3):255-61.
- Lu C, Bravo R, Caltabiano LM, Irish RM, Weerasekera G, Barr DB. The presence of dialkylphosphates in fresh fruit juices: implication for organophosphorus pesticide exposure and risk assessments. *J Toxicol Environ Health A*. 2005;68(3):209-27.
- Lucena RA, Allam MF, Costabeber IH, Villarejo ML, Navajas RF. Breast cancer risk factors: PCB congeners. *Eur J Cancer Prev* 2001;10:117-9.
- Luedersdorf R, Fuchs A, Mayer P, Skulsukai G, Schacke G. Biological assessment of exposure to antimony and lead in the glass-producing industry. *Int Arch Occup Environ Health* 1987;59:469-74.
- Lundberg C. Effect of DDT on cytochrome P-450 and oestrogen-dependent functions in mice. *Environmental Physiology & Biochemistry* 1974; 4:200-4.
- Luotamo M, Jarvisalo J, Aitio A. Assessment of exposure to polychlorinated biphenyls: analysis of selected isomers in blood and adipose tissue. *Environ Res* 1991;54(2):121-34.
- MacIntosh DL, Needham LL, Hammerstrom KA, Ryan PB. A longitudinal investigation of selected pesticide metabolites in urine. *J Expo Anal Environ Epidemiol* 1999;9(5):494-501.
- Mahaffey KR, Clickner RP, Bodurow CC. Blood organic mercury and dietary mercury intake: National Health and Nutrition Examination Survey, 1999 and 2000. *Environ Health Perspect* 2004;112(5):562-70.
- Mahaffey KR, Mergler D. Blood levels of total and organic mercury in residents of the upper St. Lawrence River Basin, Quebec: association with age, gender, and fish consumption. *Environ Res* 1998;77(2):104-14.
- Mannino DM, Holguin F, Greves HM, Savage-Brown A, Stock AL, Jones RL. Urinary cadmium levels predict lower lung function in current and former smokers: data from the Third National Health and Nutrition Examination Survey. *Thorax* 2004;59:194-8.
- Mantere P, Hänninen H, Hernberg S, Luukkonen R. A prospective follow-up study on psychological effects in workers exposed to low levels of lead. *Scand J Work Environ Health* 1984;10:43-50.

- Marcus RL. Investigation of a working population exposed to thallium. *Journal for the Society of Occupational Medicine*. 1985;35(1):4-9.
- Martin SA Jr, Harlow SD, Sowers MF, Longnecker MP, Garabrant D, Shore DL, et al. DDT metabolite and androgens in African-American farmers. *Epidemiology* 2002;1:454-8.
- Mascagni P, Consonni D, Bregante G, Chiappino G, Toffoletto F. Olfactory function in workers exposed to moderate airborne cadmium levels. *Neurotoxicology* 2003;24:717-24.
- Maskarinec G, Singh S, Meng L, Franke AA. Dietary soy intake and urinary isoflavone excretion among women from a multiethnic population. *Cancer Epidemiol Biomarkers and Prev* 1998;7:613-9.
- Masuda Y. Fate of PCDF/PCB congeners and change of clinical symptoms in patients with Yusho PCB poisoning for 30 years. *Chemosphere* 2001;43(4-7):925-30.
- Masuda Y, Schecter A, Papke O. Concentrations of PCBs, PCDFs and PCDDs in the blood of Yusho patients and their toxic equivalent contribution. *Chemosphere* 1998;37(9-12):1773-80.
- Matsuura N, Uchiyama T, Tada H, Nakamura Y, Kondo N, Morita M. Effects of dioxins and polychlorinated biphenyls (PCBs) on thyroid function in infants born in Japan—the second report from research on environmental health. *Chemosphere* 2001;45(8):1167-71.
- McDiarmid MA, Squibb K, Engelhardt SM. Biologic monitoring for urinary uranium in Gulf War I veterans. *Health Phys* 2004;87:51-6.
- McDowell MA, Dillon CF, Osterloh J, Bolger PM, Pellizzari E, Fernando R, et al. Hair mercury levels in U.S. children and women of childbearing age: reference range data from NHANES 1999-2000. *Environ Health Perspect* 2004;112(11):1165-71.
- McKee RH, Butala JH, David RM, Gans G. NTP center for the evaluation of risks to human reproduction reports on phthalates: addressing the data gaps [review]. *Reprod Toxicol* 2004;18(1):1-22.
- McKee RH, El Hawari M, Stoltz M, Pallas F, Lington AW. Absorption, disposition and metabolism of Di-isononyl phthalate (DINP) in F-344 rats. *J Appl Toxicol* 2002;22:293-302.
- Melnick RL. Is peroxisome proliferation an obligatory precursor step in the carcinogenicity of di (2-ethylhexyl) phthalate (DEHP)? *Environ Health Perspect* 2001;109(5):437-42.
- Merlo F, Andreassen A, Weston A, Pan CF, Haugen A, Valerio F, et al. Urinary excretion of 1-hydroxypyrene as a marker for exposure to urban air levels of polycyclic aromatic hydrocarbons. *Cancer Epidemiol Biomarkers Prev* 1998;7(2):147-55.
- Michalek JE, Akhtar FZ, Arezzo JC, Garabrant DH, Albers JW. Serum dioxin and peripheral neuropathy in veterans of Operation Ranch Hand. *Neurotoxicology* 2001;22:479-90.
- Michalek JE, Akhtar FZ, Kiel JL. Serum dioxin, insulin, fasting glucose, and sex hormone-binding globulin in veterans of Operation Ranch Hand. *J Clin Endocrinol Metab* 1999;84(5):1540-3.
- Miettinen JK, Rahola T, Hattula T, Rissanen K, Tillander M. Elimination of ²⁰³Hg-methyl mercury in man. *Annals of Clinical Research* 1971;3(2):116-22.
- Milligan SR, Balasubramanian AV, Kalita JC. Relative potency of xenobiotic estrogens in an acute in vivo mammalian assay. *Environ Health Perspect* 1998;106(1):23-6.
- Ming-Hsin H, Shao-Chi C, Ju-Sun P, Kuo-Juie Y, Cheng-Wei L, Chia-Yu H, et al. Mechanism and treatment of cardiac arrhythmias in tartar emetic intoxication. *Chin Med J* 1958;76(2):103-15.
- Minoia C, Gatti A, Aprea C, Ronchi A, Sciarra G, Turci R, et al. Inductively coupled plasma mass spectrometric determination of molybdenum in urine. *Rapid Comm Mass Spectrom* 2002; 16:1313-9.
- Minoia C, Sabbioni E, Apostoli P, Pietra R, Pozzoli L, Gallorini M, et al. Trace element reference values in tissues from inhabitants of the European community I. A study of 46 elements in urine, blood, and serum of Italian subjects. *Sci Total Environ* 1990;95:89-105.
- Misra UK, Gawdi G, Pizzo SV. Induction of mitogenic signaling in the ILN prostate cell line on exposure to submicromolar concentrations of cadmium. *Cell Signal* 2003;15:1059-70.
- Mocarelli P, Needham LL, Marocchi A, Patterson DG Jr, Brambilla P, Gerthoux PM, et al. Serum concentrations of 2,3,7,8-tetrachlorodibenzo-p-dioxin and test results from selected residents of Seveso, Italy. *J Toxicol Environ Health* 1991;32(4):357-66.
- Morgan DP, Hetzler HL, Slach EF, Lin LI. Urinary excretion of parantrophenol and alkyl phosphates following ingestion of methyl or ethyl parathion by human subjects. *Arch Environ Contam Toxicol* 1977;6(2-3):159-73.
- Morgan DP, Roberts RJ, Walter AW, Stockdale EM. Anemia associated with exposure to lindane. *Arch Environ Health* 1980;35(5):307-10.
- Morgan MK, Sheldon LS, Croghan CW, Jones PA, Robertson GL, Chuang JC, et al. Exposures of preschool children to chlorpyrifos and its degradation product 3,5,6-trichloro-2-pyridinol in their everyday environments. *J Expo Anal Environ*

- Epidemiol. 2004 Sep 15; [Epub ahead of print]
doi:10.1038/sj.jea.7500406
- Moriguchi J, Ezaki T, Tsukahara T, Furuki K, Fukui Y, Okamoto S, et al. Alpha1-Microglobulin as a promising marker of cadmium-induced tubular dysfunction, possibly better than beta2-microglobulin. *Toxicol Lett.* 2004;148(1-2):11-20.
- Mosconi G, Bacis M, Vitali MT, Leghissa P, Sabbioni E. Cobalt excretion in urine: results of a study on workers producing diamond grinding tools and on a control group. *Sci Total Environ* 1994;150(1-3):133-9.
- Moulin JJ, Wild P, Romazini S, Lasfargues G, Peltier A, Bozec C, et al. Lung cancer risk in hard-metal workers. *Am J Epidemiol* 1998;148:241-8.
- Mukherjee S, Palmer LJ, Kim JY, Aeschliman DB, Houk RS, Woodin MA, et al. Smoking status and occupational exposure affects oxidative DNA injury in boilermakers exposed to metal fume and residual oil fly ash. *Cancer Epidemiol Biomarkers Prev* 2004;13(3):454-60.
- Muntner P, Vupputyuri S, Coresh J, Batuman V. Blood lead and chronic kidney disease in the general United States population: results from NHANES III. *Kidney Int* 2003;63:104-50.
- Murkies A, Dalais FS, Briganti EM, Burger HG, Healy DL, Wahlqvist ML. Phytoestrogens and breast cancer in postmenopausal women: a case control study. *Menopause* 2000;7(5):289-96.
- Nachlas MM, Seligman AM. Evidence for the specificity of esterase and lipase by the use of three chromogenic substrates. *J Biol Chem* 1949;181:343-55.
- Nagel SC, vom Saal FS, Welshons WV. The effective free fraction of estradiol and xenoestrogens in human serum measured by whole cell uptake assays: physiology of delivery modifies estrogenic activity. *Proc Soc Exp Biol Med* 1998;217:300-9.
- Nan HM, Kim H, Lim HS, Choi JK, Kawamoto T, Kang JW, et al. Effects of occupation, lifestyle and genetic polymorphisms of CYP1A1, CYP2E1, GSTM1 and GSTT1 on urinary 1-hydroxypyrene and 2-naphthol concentrations. *Carcinogenesis* 2001;22(5):787-93.
- Nash D, Magder L, Lustberg M, Sherwin R, Rubin R, Kaufmann R, et al. Blood lead, blood pressure, and hypertension in perimenopausal and postmenopausal women. *JAMA* 2003;289(12):1523-31.
- National Research Council. Committee on Measuring Lead in Critical Populations. Measuring lead exposure in infants, children and other sensitive populations. Washington (DC): National Academy Press; 1993. p. 31-98.
- National Research Council (Platinum Group Metals). Medical and biologic effects of environmental pollutants. Washington (DC): National Academy of Sciences; 1977. p. 232.
- National Research Council. Toxicologic effects of methylmercury. Washington (DC): National Academy of Sciences; 2000.
- National Toxicology Program. Lead and lead compounds. In: Report on carcinogens. 11th ed. [online]. 2004. Available from URL: <http://ntp.niehs.nih.gov/ntp/roc/eleventh/profiles/s101lead.pdf>, 02/27/05.
- National Toxicology Program. Report on carcinogens. 10th ed. Washington (DC): U.S. Department of Health and Human Services, Public Health Service; December 2002.
- National Toxicology Program. Report on Carcinogens. 11th ed. [online]. 2004. Available from URL: <http://ntp.niehs.nih.gov/ntp/roc/toc11.html>, 02/18/05.
- National Toxicology Program. TR-301. Toxicology and carcinogenesis studies of o-phenylphenol alone and with 7,12-dimethylbenz(a)anthracene in Swiss CD-1 mice (dermal studies) [online]. 1986. Available at URL: <http://ntp.niehs.nih.gov/index.cfm?objectid=07078EB9-E19D-AC1B-07D4A27130861AC1>, 03/17/05.
- National Toxicology Program. TR-462. Toxicology and carcinogenesis studies of molybdenum trioxide (CAS No. 1313-27-5) in F344 Rats and B6C3F1 Mice (inhalation studies) [online]. 1997. Available from URL: <http://ntp.niehs.nih.gov/index.cfm?objectid=070A72C6-E9C1-AE67-4BFAB97AD0F427E8>, 03/17/05.
- Nebert DW, Dalton TP, Okey AB, Gonzalez FJ. Role of aryl hydrocarbon receptor-mediated induction of the CYP1 enzymes in environmental toxicity and cancer. *J Biol Chem* 2004;279(23):23847-50.
- Needham LL, Burse VW, Head SL, Korver MP, McClure PC, Andrews JS Jr, et al. Adipose tissue/serum partitioning of chlorinated hydrocarbon pesticides in humans. *Chemosphere* 1990;20:975-80.
- Negri E, Bosetti C, Fattore E, La Vecchia C. Environmental exposure to polychlorinated biphenyls (PCBs) and breast cancer: a systemic review of the epidemiological evidence. *Eur J Cancer Prev* 2003;12:509-16.
- Nesbitt PD, Lam Y, Thompson LU. Human metabolism of mammalian lignan precursors in raw and processed flaxseed. *Am J Clin Nutr* 1999;69(3):549-55.
- Nicholls J, Lasley BL, Nakajima ST, Setchell KD, Schneeman BO. Effects of soy consumption on gonadotropin secretion and acute pituitary responses to gonadotropin-releasing hormone in women. *J Nutr* 2002;132(4):708-14.

- Nicolaou G, Pietra R, Sabioni E, Mosconi G, Cassina G, Seghizzi P. Multielement determination of metals in biological specimens of hard-metal workers: a study carried out by neutron activation analysis. *Journal of Trace Elements and Electrolytes in Health and Disease* 1987;(2):73-7.
- Nielsen J, Akesson B, Skerfving S. Phthalate ester exposure—air levels and health of workers processing polyvinylchloride. *American Industrial Hygiene Association Journal* 1985;46(11):643-7.
- Nigam SK, Karnik AB, Majumder SK, Visweswariah K, Raju GS, Bai KM, et al. Serum hexachlorocyclohexane residues in workers engaged at a HCH manufacturing plant. *Int Arch Occup Environ Health* 1986;57(4):315-20.
- Nikov GN, Eshete M, Rajnarayanan RV, Alworth WL. Interactions of synthetic estrogens with human estrogen receptors. *J Endocrinol* 2001;170(1):137-45.
- Nishijo M, Nakagawa H, Honda R, Tanebe K, et al. Effects of maternal exposure to cadmium on pregnancy outcome and breast milk. *Occup Environ Med* 2002;59:394-7.
- Nolan RJ, Rick DL, Freshour NL, Saunders JH. Chlorpyrifos: pharmacokinetics in human volunteers. *Toxicol Appl Pharmacol* 1984;73:8-15.
- Noonan CW, Sarasua SM, Campagna D, Kathman SJ, Lybarger JA, Mueller PW. Effects of exposure to low levels of environmental cadmium on renal biomarkers. *Environ Health Perspect* 2002;110:151-5.
- Nordberg GF, Nordberg M. Biological monitoring of cadmium. In: Clarkson TW, Friberg L, Nordberg GF, Sager PR, eds. *Biological monitoring of toxic metals*. New York: Plenum Press; 2001. p. 151-68.
- Occupational Safety and Health Administration. 29 CFR 1910.1025. Washington, D.C. Available from URL: http://www.osha.gov/pls/oshaweb/owadisp.show_document?p_table=STANDARDS&p_id=10033, 4/07/05.
- Occupational Safety and Health Administration. 29 CFR 1910.1027. Washington, D.C. Available from URL: http://www.osha.gov/pls/oshaweb/owadisp.show_document?p_table=STANDARDS&p_id=10035, 4/7/05.
- Occupational Safety and Health Administration [online]. Available from URL: <http://www.osha.gov/comp-links.html>, 03/17/05.
- O'Flaherty EJ. Physiologically based models for bone-seeking elements. IV. Kinetics of lead disposition in humans. *Tox Appl Pharmacol* 1993;118:16-29.
- Olsson AO, Nguyen JV, Sadowski MA, Barr DB. A liquid chromatography/electrospray ionization-tandem mass spectrometry method for quantification of specific organophosphorus pesticide biomarkers in human urine. *Analytical and Bioanalytical Chemistry* 2003;376:808-15.
- Olsson IM, Bensryd I, Lundh T, Ottosson H, Skerfving S, Oskarsson A. Cadmium in blood and urine – impact of sex, age, dietary intake, iron status, and former smoking – association of renal effects. *Environ Health Perspect* 2002;110:1185-90.
- Okubo T, Suzuki T, Yokoyama Y, Kano K, Kano I. Estimation of estrogenic and anti-estrogenic activities of some phthalate diesters and monoesters by MCF-7 cell proliferation assay in vitro. *Biol Pharm Bull* 2003;26(8):1219-24.
- Onstot JD, Stanley JS. Identification of SARA compounds in adipose tissue. U.S. Environmental Protection Agency (U.S. EPA), Office of Toxic Substances; 1989. EPA-560/5-89-003.
- Orloff KG, Mistry K, Chapp P, Metcalf S, Marino R, Shelly T, et al. Human exposure to uranium in groundwater. *Environ Res* 2004;94:319-26.
- Oskarsson A, Schultz A, Skerfving S, Hallen IP, Ohlin B, Lagerkvist BJ. Total and inorganic mercury in breast milk and blood in relation to fish consumption and amalgam fillings in lactating women. *Arch Environ Health* 1996;51(3):234-41.
- Otake T, Yoshinaga J, Yanagisawa Y. Exposure to phthalate esters from indoor environment. *J Expo Anal Environ Epidemiol* 2004;14(7):524-8.
- Pagnotto LD, Walkley JE. Urinary dichlorophenol as an index of para-dichlorobenzene exposure. *American Journal of Industrial Hygiene* 1965;26:137-42.
- Papke O. PCDD/PCDF: human background data for Germany, a 10-year experience. *Environ Health Perspect* 1998;106 (Suppl 2):723-31.
- Parkinson A, Safe SH, Robertson LW, Thomas PE, Ryan DE, Reik LM, et al. Immunochemical quantitation of cytochrome P-450 isozymes and epoxide hydrolase in liver microsomes from polychlorinated or polybrominated biphenyl-treated rats. A study of structure-activity relationships. *J Biol Chem* 1983;258(9):5967-76.
- Parks LG, Ostby JS, Lambright CR, Abbott BD, Klinefelter GR, Barlow NJ, et al. The plasticizer diethylhexyl phthalate induces malformations by decreasing fetal testosterone synthesis during sexual differentiation in the male rat. *Toxicol Sci* 2000;58:339-49.
- Paschal DC, Burse V, Gunter EW, Pirkle JL, Sampson EJ, Miller DT, et al. Exposure of the U.S. population aged 6 years and older to cadmium: 1988-1994. *Arch Environ Contam Toxicol* 2000;38:377-83.
- Paschal DC, Ting BG, Morrow JC, Pirkle JL, Jackson RJ, Sampson EJ, et al. Trace metals in urine of United States

- residents: reference range concentrations. *Environ Res* 1998;76(1):53-9.
- Patterson DG, Holler HS, Smith SJ, Liddle JA, Sampson EJ, Needham LL. Human adipose data for 2,3,7,8-tetrachlorodibenzo-p-dioxin in certain U.S. samples. *Chemosphere* 1986;15:2055-60.
- Patterson DG Jr, Todd GD, Turner WE, Maggio V, Alexander LR, Needham LL. Levels of non-ortho-substituted (coplanar), mono- and di-ortho-substituted polychlorinated biphenyls, dibenzo-p-dioxins, and dibenzofurans in human serum and adipose tissue. *Environ Health Perspect* 1994;102 (Suppl 1):195-204.
- Pavuk M, Cerhan JR, Lynch CF, Kocan A, Petrik J, Chovancova J. Case-control study of PCBs, other organochlorines and breast cancer. *J Expo Anal Environ Epidemiol* 2003;13:267-75.
- Pavuk M, Cerhan JR, Lynch CF, Schecter A, Petrik J, Chovancova J, et al. Environmental exposure to PCBs and cancer incidence in eastern Slovakia. *Chemosphere* 2004;54:1509-20.
- Payton M, Hu H, Sparrow D, Weiss ST. Low-level lead exposure and renal function in the Normative Aging Study. *Am J Epidemiol* 1994;140:821-29.
- Peck CC, Albro PW. Toxic potential of the plasticizer di (2-ethylhexyl) phthalate in the context of its disposition and metabolism in primates and man. *Environ Health Perspect* 1982;45:11-7.
- Perera FP, Rauh V, Tsai WY, Kinney P, Camann D, Barr D, et al. Effects of transplacental exposure to environmental pollutants on birth outcomes in a multiethnic population. *Environ Health Perspect* 2003;111(2):201-5.
- Perry MJ, Christiani DC, Mathew J, Degenhardt D, Tortorelli J, Strauss J, et al. Urinalysis of atrazine exposure in farm pesticide applicators. *Toxicology and Industrial Health* 2000;16(7-8): 285-90.
- Pesch A, Wilhelm M, Rostek U, Schmitz N, Weishoff-Houben M, Ranft U, et al. Mercury concentrations in urine, scalp hair, and saliva in children from Germany. *J Expo Anal Environ Epidemiol* 2002;12(4):252-8.
- Peters HA, Gocmen A, Cripps DJ, Bryan GT, Dogramaci I. Epidemiology of hexachlorobenzene-induced porphyria in Turkey: clinical and laboratory follow-up after 25 years. *Arch Neurol* 1982;39(12):744-9.
- Pethran A, Schierl R, Hauff K, Grimm CH, Boos KS, Nowak D. Uptake of antineoplastic agents in pharmacy and hospital personnel. Part 1: monitoring of urinary concentrations. *Int Arch Occup Environ Health* 2003;76(1):5-10
- Pirkle JL, Flegal KM, Bernert JT, Brody DJ, Etzel RA, Maurer KR. Exposure of the U.S. population to environmental tobacco smoke: the Third National Health and Nutrition Examination Survey, 1988-1991. *JAMA* 1996;275:1233-40.
- Pirkle JL, Kaufmann RB, Brody DJ, Hickman T, Gunter EW, Paschal DC. Exposure of the U.S. population to lead: 1991-1994. *Environ Health Perspect* 1998;106:745-50.
- Pisati G, Zedda S. Outcome of occupational asthma due to cobalt hypersensitivity. *Sci Total Environ.* 1994;150(1-3):167-71.
- Porta M, Zumeta E. Implementing the Stockholm Treaty on Persistent Organic Pollutants. *Occup Environ Med* 2002;59:651-3.
- Radomski JL, Astolfi E, Deichmann WB, Rey AA. Blood levels of organochlorine pesticides in Argentina: occupationally and nonoccupationally exposed adults, children and newborn infants. *Toxicol Appl Pharmacol* 1971;20(2):186-93.
- Rahola T, Hattula T, Korolainen A, Miettinen JK. Elimination of free and protein-bound ionic mercury ($^{203}\text{Hg}^{2+}$) in man. *Annals of Clinical Research* 1973;5:214-9.
- Ramesh A, Walker SA, Hood DB, Guillen MD, Schneider K, Weyand EH. Bioavailability and risk assessment of orally ingested polycyclic aromatic hydrocarbons. *International Journal of Toxicology* 2004;23(5):301-33.
- Ray DE, Forshaw PJ. Pyrethroid insecticides: poisoning syndromes, synergies, and therapy. *J Toxicol Clin Toxicol* 2000;38:95-101.
- Redmond CK, Strobino BR, Cypess RH. Cancer experience among coke by-product workers. *Ann N Y Acad Sci* 1976;271:102-15.
- Renes LE. Antimony poisoning in industry. *Industrial Hygiene and Occupational Medicine* 1953;99-108.
- Rice DC, Schoeny R, Mahaffey K. Methods and rationale for derivation of a reference dose for methyl mercury by the U.S. EPA. *Risk Anal* 2003; 23(1):107-15.
- Rice DC. The US EPA reference dose for methyl mercury: sources of uncertainty. *Environ Res* 2004;95:406-13.
- Richter ED, Kowalski M, Leventhal A, Grauer F, Marzouk J, Brenner S, et al. Illness and excretion of organophosphate metabolites four months after household pest extermination. *Arch Environ Health* 1992;47(2):135-8.
- Rier S, Foster WG. Environmental dioxins and endometriosis. *Seminars in Reproductive Medicine* 2003;21:145-54.

- Ritchie KA, Gilmour WH, Macdonald EB, Burke FJ, McGowan DA, Dale IM, et al. Health and neuropsychological functioning of dentists exposed to mercury. *Occup Environ Med* 2002;59(5):287-93.
- Robbins PJ, Cherniack MG. Review of the biodistribution and toxicity of the insect repellent N,N-diethyl-m-toluamide (DEET). *J Toxicol Environ Health* 1986;18(4):503-25.
- Roels HA, Boeckx M, Ceulemans E, Lauwerys RR. Urinary excretion of mercury after occupational exposure to mercury vapor and influence of the chelating agent meso-2,3-dimercaptosuccinic acid (DMSA). *British Journal of Industrial Medicine* 1991;48:247-53.
- Roels HA, Hoet P, Lison D. Usefulness of biomarkers of exposure to inorganic mercury, lead, or cadmium in controlling occupational and environmental risks of nephrotoxicity. *Ren Fail* 1999;21(3-4):251-62.
- Roggi C, Minoia C, Sciarra GF, Apostoli P, Maccarini L, Magnaghi S, et al. Urinary 1-hydroxypyrene as a marker of exposure to pyrene: an epidemiological survey on a general population group. *Sci Total Environ* 1997;199(3):247-54.
- Rugman FP. Aplastic anaemia associated with organochlorine pesticide: case reports and review of evidence. *J Clin Pathol* 1990;42(3):98-101.
- Safe SH, Pallaroni L, Yoon K, Gaido K, Ross S, Saville B, et al. Toxicology of environmental estrogens. *Reprod Fertil Dev* 2001;13(4):307-15.
- Sala M, Ribas-Fito N, Cardo E, de Muga ME, Marco E, Mazon C, et al. Levels of hexachlorobenzene and other organochlorine compounds in cord blood: exposure across placenta. *Chemosphere* 2001;43:895-901.
- Sala M, Sunyer J, Otero R, Santiago-Silva M, Camps C, Grimalt J. Organochlorine in the serum of inhabitants living near an electrochemical factory. *Occup Environ Med* 1999;56(3):152-8.
- Sallsten G, Barregard L, Schutz A. Decrease in mercury concentration in blood after long term exposure: a kinetic study of chloralkali workers. *British Journal of Industrial Medicine* 1993;50(9):814-21.
- Sanchez-Sicilia L, Seto DS, Nakamoto S, Kolff WJ. Acute mercurial intoxication treated by hemodialysis. *Ann Intern Med* 1963;59(5):692-706.
- Sandborgh-Englund G, Elinder CG, Langworth S, Schutz A, Ekstrand J. Mercury in biological fluids after amalgam removal. *J Dent Res* 1998;77(4):615-24.
- Sanzo JM, Dorronsoro M, Amiano P, Amurrio A, Aguinagalde FX, Azpiri MA. Estimation and validation of mercury intake associated with fish consumption in an EPIC cohort of Spain. *Public Health Nutr* 2001;4(5):981-8.
- SAS Institute, Inc. SAS/STAT User's Guide. Version 8. Cary, NC: SAS Institute, Inc.; 1999.
- Satarug S, Baker JR, Reilly PE, Moore MR, Williams DJ. Cadmium levels in the lung, liver, kidney cortex, and urine samples from Australians without occupational exposure to metals. *Arch Environ Health* 2002;57(1):69-77.
- Sauerhoff MW, Braun WH, Blau GE, Gehring PJ. The fate of 2,4-dichlorophenoxyacetic acid (2,4-D) following oral administration to man. *Toxicology* 1977;8:3-11.
- Schaller KH, Manke G, Raithel HJ, Buhlmeyer G, Schmidt M, Valentin H. Investigations of thallium-exposed workers in cement factories. *Int Arch Occup Environ Health* 1980;47(3):223-31.
- Scheepers PT, Coggon D, Knudsen LE, Anzion R, Autrup H, Bogovski S, et al. Biomarkers for occupational diesel exhaust exposure monitoring (BIOMODEM)—a study in underground mining. *Toxicol Lett* 2002;134(1-3):305-17.
- Schettgen T, Heudorf U, Drexler H, Angerer J. Pyrethroid exposure of the general population—is this due to diet? *Toxicology Lett* 2002a;134(1-3):141-5.
- Schettgen T, Koch HM, Drexler H, Angerer J. New gas chromatographic-mass spectrometric method for the determination of urinary pyrethroid metabolites in environmental medicine. *Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences* 2002b;778(1-2):121-30.
- Schierl R. Urinary platinum levels associated with dental gold alloys. *Arch Environ Health* 2001;56(3):283-6.
- Schierl R, Fries HG, van de Weyer C, Fruhmann G. Urinary excretion of platinum from platinum-industry workers. *Occup Environ Med* 1998;55(2):138-4.
- Schnorr TM, Lawson CC, Whelan EA, Dankovic DA, Deddens JA, Piacitelli LA, et al. Spontaneous abortion, sex ratio, and paternal occupational exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Environ Health Perspect* 2001;109(11):1127-32.
- Schober SE, Sinks TH, Jones RL, Bolger PM, McDowell M, Osterloh J. Blood mercury levels in US children and women of childbearing age, 1999-2000. *JAMA* 2003; 289:1667-74.
- Schramel P, Wendler I, Angerer J. The determination of metals (antimony, bismuth, lead, cadmium, mercury, palladium, platinum, tellurium, thallium, tin and tungsten) in urine samples by inductively coupled plasma-mass spectrometry. *Int Arch Occup Environ Health* 1997;69(3):219-23.
- Schwartz BS, Lee BK, Lee GS, Stewar WF, Lee SS, Hwang KY, et al. Association of blood lead, dimercaptosuccinic acid-chelatable lead, and tibia lead with neurobehavioral test scores in South Korean lead workers. *Am J Epidemiol* 2001;153(5):453-64.

- Schwartz J. Lead, blood pressure and cardiovascular disease in men. *Arch Environ Health* 1995; 50:31-7.
- Schwartz J, Landrigan PJ, Baker EL, Orenstein WA, von Lindern IH. Lead induced anemia: dose-response relationships and evidence for a threshold. *Am J Public Health* 1990;80:165-8.
- Seifert B, Becker K, Helm D, Krause C, Schulz C, Seiwert M. The German Environmental Survey 1990/1992 (GerES II): reference concentrations of selected environmental pollutants in blood, urine, hair, house dust, drinking water and indoor air. *J Expo Anal Environ Epidemiol* 2000;10:552-65.
- Selim S, Hartnagel RE Jr, Osimitz TG, Gabriel KL, Schoenig GP. Absorption, metabolism, and excretion of N,N-diethyl-m-toluamide following dermal application to human volunteers. *Fundam Appl Toxicol* 1995;25:95-100.
- Seow A, Shi CY, Franke AA, Hankin JH, Lee H, Yu MC. Isoflavonoid levels in spot urine are associated with frequency of dietary soy intake in a population-based sample of middle-aged and older Chinese in Singapore. *Cancer Epidemiol Biomarkers Prev* 1998;7:135-40.
- Serdar B, Waidyanatha S, Zheng Y, Rappaport SM. Simultaneous determination of urinary 1- and 2-naphthols, 3- and 9-phenanthrols, and 1-pyrenol in coke oven workers. *Biomarkers* 2003a;8(2):93-109.
- Serdar B, Egeghy PP, Waidyanatha S, Gibson R, Rappaport SM. Urinary biomarkers of exposure to jet fuel (JP-8). *Environ Health Perspect* 2003b;111(14):1760-4.
- Setchell KD, Brown NM, Desai P, Zimmer-Nechemias L, Wolfe BE, Brashear WT, et al. Bioavailability of pure isoflavones in healthy humans and analysis of commercial soy isoflavone supplements. *J Nutr* 2001;131(4 Suppl):1362S-75S.
- Setchell KD, Brown NM, Lydeking-Olsen E. The clinical importance of the metabolite equol—a clue to the effectiveness of soy and its isoflavones. *J Nutr* 2002;132(12):3577-84.
- Setchell KD, Cassidy A. Dietary isoflavones: biological effects and relevance to human health. *J Nutr* 1999;129(3):758S-67S.
- Setchell KD, Zimmer-Nechemias L, Cai J, Heubi JE. Exposure of infants to phyto-oestrogens from soy-based infant formula. *Lancet* 1997;350(9070):23-7.
- Shealy DB, Barr JR., Ashley DL, Patterson DG Jr, Camann DE, Bond AE. Correlation of environmental carbaryl measurements with serum and urinary 1-naphthol measurements in a farmer applicator and his family. *Environ Health Perspect* 1997;105:510-3.
- Shealy DB, Bonin MA, Wooten JV, Ashley DL, Needham LL. Application of an improved method for the analysis of pesticides and their metabolites in the urine of farmer applicators and their families. *Environment International* 2004; 22:661-75.
- Sherlock J, Hislop D, Newton G, Topping G, Whittle K. Elevation of mercury in human blood from controlled ingestion of methyl mercury in fish. *Human Toxicology* 1984;2:117-31.
- Shirakawa T, Kusaka Y, Fujimura N, Goto S, Kato M, Heki S, et al. Occupational asthma from cobalt sensitivity in workers exposed to hard metal dust. *Chest* 1989;95:29-37.
- Sievers E, Schleyerbach U, Schaub J. Molybdenum in infancy: methodical investigation of urinary excretion. *J Trace Elem Med Biol* 2001;15(2-3):149-54.
- Silva MJ, Barr DB, Reidy JA, Malek NA, Hodge CC, Caudill SP, et al. Urinary levels of seven phthalate metabolites in the U.S. population from the National Health and Nutrition Examination Survey (NHANES) 1999-2000 [published erratum appears in *Environ Health Perspect* 2004; 112(5):A270]. *Environ Health Perspect* 2004;112(3):331-8.
- Simpson GR, Higgins V, Chapman J, Birmingham S. Exposure of council and forestry workers to 2,4,5-T. *Med J Aust* 1978;18;2(11):536-7.
- Siwinska E, Mielzynska D, Bubak A, Smolik E. The effect of coal stoves and environmental tobacco smoke on the level of urinary 1-hydroxypyrene. *Mutat Res* 1999;445(2):147-53.
- Siwinska E, Mielzynska D, Kapka L. Association between urinary 1-hydroxypyrene and genotoxic effects in coke oven workers. *Occup Environ Med.* 2004;61(3):e10.
- Slavin JL, Karr SC, Hutchins AM, Lampe JW. Influence of soybean processing, habitual diet, and soy dose on urinary isoflavonoid excretion. *Am J Clin Nutr* 1998;68(6 Suppl):1492S-5S.
- Smallwood AW, DeBord KE, Lowry LK. N,N'-diethyl-m-toluamide (m-DET): analysis of an insect repellent in human urine and serum by high-performance liquid chromatography. *J Anal Toxicol* 1992;16(1):10-3.
- Smith JC, Allen PV, Turner MD, Most B, Fisher HL, Hall LL. The kinetics of intravenously administered methyl mercury in man. *Toxicol Appl Pharmacol* 1994;128(2):251-6.
- Smith JC, Farris FF. Methyl mercury pharmacokinetics in man: a reevaluation. *Toxicol Appl Pharmacol* 1996;37:245-52.
- Smith PJ, Langolf GD, Goldberg J. Effects of occupational exposure to elemental mercury on short term memory. *British Journal of Industrial Medicine* 1983;40:413-9.
- Smith RG, Vorwald AJ, Patil LS, Mooney TF. Effects of exposure to mercury in the manufacture of chlorine. *American Industrial Hygiene Association Journal* 1970;31:687-700.

- Smith T, Edmonds CJ, Barnaby CF. Absorption and retention of cobalt in man by whole-body counting. *Health Phys* 1972;22:359-67.
- Soderlund DM, Clark JM, Sheets LP, Mullin LS, Piccirillo VJ, Sargent D, et al. Mechanisms of pyrethroid neurotoxicity: implications for cumulative risk assessment. *Toxicology* 2002;171:3-59.
- Soldin OP, Hanak B, Soldin SJ. Blood lead concentrations in children: new ranges. *Clin Chim Acta* 2003;327:109-13.
- Staessen JA, Buchet JP, Ginocchio G, Lauwerys R, Lijnen P, Roels H, et al. Public health implications of environmental exposure to cadmium and lead: an overview of epidemiological studies in Belgium. *J Cardiovasc Risk* 1996;3:26-41.
- Staessen J, Roels HA, Emelianov D, Kuznetsova T, Thijs L, Vangronsveld J, et al. Environmental exposure to cadmium, forearm bone density, and risk of fractures: prospective population study. *Lancet* 1999;353:1140-4.
- Staessen JA, Roels H, Lauwerys RR, Amery A. Low-level lead exposure and blood pressure. *J Hum Hypertens* 1995;9:303-27.
- Stattin P, Adlercreutz H, Tenkanen L, Jellum E, Lumme S, Hallmans G, et al. Circulating enterolactone and prostate cancer risk: a Nordic nested case-control study. *Int J Cancer* 2002;99(1):124-9.
- Stehr-Green, PA. Demographic and seasonal influences on human serum pesticide residue levels. *J Toxicol Environ Health* 1989;27:405-21.
- Stern AH, Smith AE. An assessment of the cord blood: maternal blood methyl mercury ratio: implications for risk assessment. *Environ Health Perspect* 2003;111(12):1465-70.
- Strom BL, Schinnar R, Ziegler EE, Barnhart KT, Sammel MD, Macones GA, et al. Exposure to soy-based formula in infancy and endocrinological and reproductive outcomes in young adulthood. *JAMA* 2001;286:807-14.
- Sturgeon SR, Brock JW, Potischman N, Needham LL, Rothman N, Brinton LA, et al. Serum concentrations of organochlorine compounds and endometrial cancer risk (United States). *Cancer Causes Control* 1998;9(4):417-24.
- SUDAAN User's Manual, Release 8.0, Research Triangle Park (NC): Research Triangle Institute; 2001.
- Suwazono Y, Kobayashi E, Okubo Y, Nogawa K, Kido T, Nakagawa H. Renal effects of cadmium exposure in cadmium nonpolluted areas in Japan. *Environ Res* 2000;84 (Section A):44-55.
- Suzuki T, Hongo T, Yoshinaga J, Imai H, Nakazawa M, Matsuo N, et al. The hair-organ relationship in mercury concentration in contemporary Japanese. *Arch Environ Health* 1993;48(4):221-9.
- Swennen B, Buchet JP, Stanescu D, Lison D, Lauwerys R. Epidemiological survey of workers exposed to cobalt oxides, cobalt salts, and cobalt metal. *British Journal of Industrial Medicine* 1993;50(9):835-42.
- Szanişzlo J, Ungvary G. Polycyclic aromatic hydrocarbon exposure and burden of outdoor workers in Budapest. *J Toxicol Environ Health A* 2001;62(5):297-306.
- Takahashi W, Parks LH. Organochlorine pesticide residues in human tissues, Hawaii, 1968-1980. *Hawaii Med J* 1982; 41(9):250-1.
- Takamiya K. Monitoring of urinary alkyl phosphates in pest-control operators exposed to various organophosphorus insecticides. *Bull Environ Contam Toxicol* 1994;52(2):190-5.
- Taylor JK. Quality assurance of chemical measurements. Chelsea (MI): Lewis Publishing; 1987.
- Teede HJ, Dalais FS, Kotsopoulos D, Liang YL, Davis S, McGrath BP. Dietary soy has both beneficial and potentially adverse cardiovascular effects: a placebo-controlled study in men and postmenopausal women. *J Clin Endocrinol Metab* 2001;86(7):3053-60.
- Telisman S, Cvitkovic P, Jurasovic J, Pizent A, Gavella M, Rocic B. Semen quality and reproductive endocrine function in relation to biomarkers of lead, cadmium, zinc, and copper in men. *Environ Health Perspect*. 2000;108(1):45-53.
- Tepper A, Burt S, Piacitelli L, Patterson DG Jr. Coplanar PCBs and the relative contribution of coplanar PCBs, PCDDs, and PCDFs to the total 2,3,7,8-TCDD toxicity equivalents in human serum. *Chemosphere* 1997;35(3):503-11.
- Thomas D, Goldhaber M, Petitti D, Swan SH, Rappaport E, Hertz-Picciotto I. Reproductive outcome in women exposed to malathion. *Am J Epidemiol* 1990;132(4):794-5.
- Thomssen H, Hoffmann B, Schank M, Hohler T, Thabe H, Meyer zum Buschenfelde KH, et al. Cobalt-specific T lymphocytes in synovial tissue after an allergic reaction to a cobalt alloy joint prosthesis. *J Rheumatol* 2001;28(5):1121-8.
- Timchalk C, Selim S, Sangha G, Bartels MJ. The pharmacokinetics and metabolism of ¹⁴C/¹³C-labeled ortho-phenylphenol formation following dermal application to human volunteers. *Hum Exp Toxicol*. 1998;17(8):411-7.
- Ting BG, Paschal DC, Jarrett JM, Pirkle JL, Jackson RJ, Sampson EJ, et al. Uranium and thorium in urine of United States residents: reference range concentrations. *Environ Res* 1999;81:45-51.
- Toriba A, Chetiyankornkul T, Kizu R, Hayakawa K. Quantification of 2-hydroxyfluorene in human urine by

- column-switching high performance liquid chromatography with fluorescence detection. *Analyst* 2003;128(6):605-10.
- To-Figueras J, Sala M, Otero R, Barrot C, Santiago-Silva M, Rodamilans M, et al. Metabolism of hexachlorobenzene in humans: association between serum levels and urinary metabolites in a highly exposed population. *Environ Health Perspect* 1997;105(1):78-83.
- Tsai HT, Wu MT, Hauser R, Rodrigues E, Ho CK, Liu CL, et al. Exposure to environmental tobacco smoke and urinary 1-hydroxypyrene levels in preschool children. *Kaohsiung J Med Sci* 2003;19(3):97-104.
- Tsai PJ, Shih TS, Chen HL, Lee WJ, Lai CH, Liou SH. Urinary 1-hydroxypyrene as an indicator for assessing the exposures of booth attendants of a highway toll station to polycyclic aromatic hydrocarbons. *Environ Sci Technol* 2004;38(1):56-61.
- Tsongas TA, Meglen RR, Walravens PA, Chappell WR. Molybdenum in the diet: an estimate of average daily intake in the United States. *Am J Clin Nutr* 1980;33(5):1103-7.
- Tunnessen WW, McMahon KJ, Baser M. Acrodynia: exposure to mercury from fluorescent light bulbs. *Pediatrics* 1987;79:786-9.
- Turnlund JR, Keyes WR, Peiffer GL. Molybdenum absorption, excretion, and retention studied with stable isotopes in young men at five intakes of dietary molybdenum. *Am J Clin Nutr* 1995;62(4):790-6.
- Uehara M, Lapcik O, Hampl R, Al-maharik N, Makela T, Wahala K, et al. Rapid analysis of phytoestrogens in human by time-resolved fluoroimmunoassay. *Journal of Steroid Biochemistry and Molecular Biology (Oxford)* 2000b;72:273-82.
- Uehara M, Arai Y, Watanabe S, Adlercreutz H. Comparison of plasma and urinary phytoestrogens in Japanese and Finnish women by time-resolved fluoroimmunoassay. *Biofactors* 2000a;12:217-25.
- Umweltbundesamt [Federal Environmental Agency]. German Environmental Survey 1998 (GerES III). Federal Ministry for the Environment, Nature Conservation and Nuclear Safety. Germany. 2002 [online]. 2002. Available from URL: <http://www.umweltbundesamt.de/survey-e/us98/blut.htm>, 02/22/05.
- U.S. Department of Health and Human Services (U.S. DHHS). Public Health Service. Subcommittee on Risk Management. Dental amalgam: a scientific review and recommended Public Health Service strategy for research, education, and regulation. Washington (DC): DHHS; January 1993.
- U.S. Environmental Protection Agency (U.S. EPA). Deposition of air pollutants to the Great Waters – 3rd Report to Congress, 2000 [online]. Available from URL: <http://www.epa.gov/air/oaqps/gr8water/2ndrpt/chap2.pdf>, 2/22/05.
- U.S. Environmental Protection Agency (U.S. EPA). Hexachlorobenzene health advisory. EPA/820K87110. Washington (DC): U.S. EPA; March 1987.
- U. S. Environmental Protection Agency (U.S. EPA). Methyl mercury. Integrated Risk Information System (IRIS). Revised 07/27/2001 [online]. Available from URL: www.epa.gov/iris/subst/0073.htm, 3/03/05.
- U. S. Environmental Protection Agency (U.S. EPA). Cadmium compounds . Hazards summary. [online]. 1992. <http://www.epa.gov/ttn/atw/hlthef/cadmium.html>, 4/26/05.
- U.S. Environmental Protection Agency (U.S. EPA). Molybdenum [online]. 1993. Available from URL; <http://www.epa.gov/iris/subst/0425.htm>, 03/18/05.
- U.S. Environmental Protection Agency (U.S. EPA). Nonoccupational Pesticide Exposure Study (NOPES). EPA/600/3-90/003. Washington (DC): U.S. EPA; January 1990.
- U.S. Environmental Protection Agency (U.S. EPA). Office of Pesticide Programs. EPA R.E.D. Facts: Alachlor. Washington (DC): U.S. EPA; December 1998.
- U.S. Environmental Protection Agency (U.S. EPA). Office of Pesticide Programs. Interim Re-Registration Eligibility Decision: Atrazine. Case No. 0062. Washington (DC): U.S. EPA; January 2003a.
- U.S. Environmental Protection Agency (U.S. EPA). Office of Prevention, Pesticides and Toxic Substances, EPA Office of Pesticide Programs, Environmental Fate and Effects Division. White paper on potential developmental effects of atrazine on amphibians. Washington (DC): U.S. EPA; May 2003b.
- U.S. Environmental Protection Agency (U.S. EPA). Office of Pesticide Programs. EPA R.E.D. Facts: Metolachlor. Washington (DC): U.S. EPA; April 1995.
- U. S. Environmental Protection Agency (U.S. EPA). Office of Prevention Pesticides and Toxic Substances. Reregistration Eligibility Decision (RED): DEET. EPA 738-R-98-010. Washington (DC): U.S. EPA; September 1998. p. 1-118.
- U.S. Environmental Protection Agency (U.S. EPA), Office of Pesticide Programs. Status summary of the organophosphate review process [online]. Available from URL: <http://www.epa.gov/pesticides/op/status.htm>, 4/8/05.
- U. S. Environmental Protection Agency (U.S. EPA). Office of Prevention Pesticides and Toxic Substances. Pesticide industry sales and usage - 2000 and 2001 market estimates. Washington (DC): U.S. EPA; May, 2004 [also online] http://www.epa.gov/oppbead1/pestsales/01pestsales/market_estimates2001.pdf, 5/16/05.

- U.S. Geological Survey (U.S.GS). Cadmium. Mineral commodity summary [online]. 2004. Available from URL: <http://minerals.usgs.gov/minerals/pubs/commodity/cadmium/admimcs04.pdf>, 03/18/05.
- U.S. Nuclear Regulatory Commission (U.S. NRC). U.S. Nuclear Regulatory Commission (NRC) Guide 8.22–Bioassay at uranium mills. Washington (DC): NRC; July 1978.
- Vahter M, Mottet NK, Friberg L, Lind B, Shen DD, Burbacher T. Speciation of mercury in the primate blood and brain following long-term exposure to methyl mercury. *Tox Appl Pharmacol* 1994;124:221-9.
- Valentin-Blasini L, Blount B, Caudill S, Needham L. Urinary and serum concentrations of seven phytoestrogens in a human reference population subset. *J Expos Anal Environ Epidemiol* 2003;13:276-82.
- Van Oostdam JC, Dewailly E, Gilman A, Hansen JC, Odland JO, Chashchin V, et al. Circumpolar maternal blood contaminant survey, 1994-1997 organochlorine compounds. *Sci Total Environ* 2004;330:55-70.
- van Wijnen JH, Slob R, Jongmans-Liedekerken G, van de Weerd RH, Woudenberg F. Exposure to polycyclic aromatic hydrocarbons among Dutch children. *Environ Health Perspect* 1996;104(5):530-4.
- Vaughan GT, Florence TM. Platinum in the human diet, blood, hair and excreta. *Sci Total Environ* 1992;111(1):47-58.
- Waalkes MP. Cadmium carcinogenesis. *Mutat Res* 2003;533:107-20.
- Waliszewski SM, Pardio VT, Chantiri JN, Infanzon RM, Rivera J. Organochlorine pesticide residues in adipose tissue of Mexicans. *Sci Total Environ* 1996;181(2):125-31.
- Walker KC, Goette MB, Batchelor GS. Pesticide residues in foods: dichlorodiphenyltrichloroethane and dichlorodiphenyldichloroethylene content of prepared meals. *J Agric Food Chem* 1954;2:1034-7.
- Ward EM, Schulte P, Grajewski B, Andersen A, Patterson DG Jr, Turner W, et al. Serum organochlorine levels and breast cancer: a nested case-control study of Norwegian women. *Cancer Epidemiol Biomarkers Prev* 2000;9:1357-67.
- Wariishi M, Suzuki Y, Nishiyama K. Chlordane residues in normal human blood. *Bull Environ Contam Toxicol* 1986;36(5):635-43.
- Warner M, Eskenazi B, Mocarelli P, Gerthoux PM, Samuels S, Needham, et al. Serum dioxin concentrations and breast cancer risk in the Seveso women's health study. *Environ Health Persp* 2002;110(7):625-8.
- Werrin M. Chemical food poisoning. *Quarterly Bulletin of the Association of Food and Drug Officials* 1962; 27:38-45.
- Whincup PH, Gilg JA, Emberson JR, Jarvis MJ, Feyerabend C, Bryant A, et al. Passive smoking and risk of coronary heart disease and stroke: prospective study with cotinine measurement. *BMJ* 2004;329(7459):200-5.
- White MA, Sabbioni E. Trace element reference values in tissues from inhabitants of the European Union. X. A study of 13 elements in blood and urine of a United Kingdom population. *Sci Total Environ* 1998;216:253-70.
- Whitemore RW, Immerman FW, Camann DE, Bond AE, Lewis RG, Schaum JL. Non-occupational exposures to pesticides for residents of two U.S. cities. *Arch Environ Contam Toxicol* 1994;26(1):47-59.
- Wilhelm M, Ewers U, Schulz C. Revised and new reference values for some persistent organic pollutants (POPs) in blood for human biomonitoring in environmental medicine. *Int J Hyg Environ Health* 2003;206:223-9.
- Wilhelm M, Ewers U, Schulz C. Revised and new reference values for some trace elements in blood and urine for human biomonitoring in environmental medicine. *Int J Hyg Environ Health* 2004;207(1):69-73.
- Williams RL, Bernard CE, Krieger RI. Human exposure to indoor residential cyfluthrin residues during a structured activity program. *J Expos Anal Environ Epidemiol* 2003;13(2):112-9.
- Wilson HK. Related biological monitoring values for occupational exposure: a United Kingdom perspective. *Int Arch Occup Environ Health* 1999;72(4):274-8.
- Wilson NK, Chuang JC, Lyu C, Menton R, Morgan MK. Aggregate exposures of nine preschool children to persistent organic pollutants at day care and at home. *J Expo Anal Environ Epidemiol* 2003;13(3):187-202.
- Wolff MS, Berkowitz GS, Brower S, Senie R, Bleiweiss IJ, Tartert P, et al. Organochlorine exposures and breast cancer risk in New York City women. *Environ Res* 2000;84:151-61.
- Wones RG, Stadler BL, Frohman, LA. Lack of effect of drinking water barium on cardiovascular risk factor. *Environ Health Perspect* 1990;85:355-9.
- World Health Organization (WHO). International Agency for Research on Cancer (IARC). Polychlorinated dibenzo-p-dioxins and dibenzofurans. In: IARC monographs on the evaluation of carcinogenic risk to humans. Vol. 69. Geneva: WHO; 1997.
- World Health Organization (WHO). International Programme on Chemical Safety [online]. Available from URL: <http://www.inchem.org>, 03/18/05.
- World Health Organization (WHO). International Programme on Chemical Safety. Environmental Health Criteria 134. [online]. 1992. <http://www.inchem.org/documents/ehc/ehc/ehc134.htm>.

World Health Organization (WHO). Molybdenum. In: Trace elements in human nutrition and health. Geneva: WHO; 1996. p. 144-54.

Xu X, Duncan AM, Wangen KE, Kurzer MS. Soy consumption alters endogenous estrogen metabolism in postmenopausal women. *Cancer Epidemiol Biomarkers Prev* 2000;9(8):781-6.

Yang M, Kim S, Lee E, Cheong HK, Chang SS, Kang D, et al. Sources of polycyclic aromatic hydrocarbon exposure in non-occupationally exposed Koreans. *Environ Mol Mutagen* 2003;42(4):250-7.

Yoshida T, Andoh K, Fukuhara M. Urinary 2,5-dichlorophenol as biological index for p-dichlorobenzene exposure in the general population. *Arch Environ Contam Toxicol* 2002;43(4):481-5.

Zacharewski TR, Meek MD, Clemons JH, Wu ZF, Fielden MR, Matthews JB. Examination of the in vitro and in vivo estrogenic activities of eight commercial phthalate esters. *Toxicological Sci* 1998;46:282-93.

Zeitz P, Orr MF, Kaye WE. Public health consequences of mercury spills: hazardous substances emergency events surveillance system, 1993-1998. *Environ Health Perspect* 2002;110:129-32.

Zhang YL, Zhao YC, Wang JX, Zhu HD, Liu QF, Fan YG, et al. Effect of environmental exposure to cadmium on pregnancy outcome and fetal growth: a study on healthy pregnant women in China. *J Environ Sci Health B* 2004;39:2507-15.

Zheng W, Dai Q, Custer LJ, Shu XO, Wen WQ, Jin F, et al. Urinary excretion of isoflavonoids and the risk of breast cancer. *Cancer Epidemiol Biomarkers Prev* 1999;8(1):35-40.

Zschiesche W, Schaller KH, Weltle D. Exposure to soluble barium compounds: an interventional study in arc welders. *Int Arch Occup Environ Health* 1992;64(1):13-23.