

## MDMA (Ecstasy)

MDMA (3,4 methylenedioxymethamphetamine)<sup>†</sup> is a synthetic, psychoactive drug that is chemically similar to the stimulant methamphetamine and the hallucinogen mescaline. It produces an energizing effect as well as feelings of euphoria, emotional warmth, and distortions in time perception and tactile experiences.

### ***How is MDMA Abused?***

MDMA is taken orally as a capsule or tablet. It was initially popular among white adolescents and young adults in the nightclub scene or at weekend-long dance parties known as raves. However, the profile of the typical MDMA user has changed, and the drug now affects a broader range of ethnic groups. MDMA is also popular among urban gay males—some report using MDMA as part of a multiple-drug experience that includes marijuana, cocaine, methamphetamine, ketamine, and other legal and illegal substances.

### ***How does MDMA Affect the Brain?***

MDMA exerts its primary effects in the brain on neurons that use the chemical serotonin to communicate with other neurons. MDMA binds to the serotonin transporter, which is responsible for removing serotonin from the synapse

(or space between adjacent neurons) when the signal between neurons needs to be terminated. MDMA also causes excessive release of serotonin and has similar but less potent effects on neurons that contain dopamine and norepinephrine. The serotonin system plays an important role in regulating mood, aggression, sexual activity, sleep, and sensitivity to pain.

MDMA can produce confusion, depression, sleep problems, drug craving, and severe anxiety. These problems can occur soon after taking the drug or, sometimes, even days or weeks after taking MDMA. In addition, chronic users of MDMA perform more poorly than nonusers on certain types of cognitive or memory tasks, although some of these effects may be due to the use of other drugs in combination with MDMA. Research in animals indicates that MDMA can be harmful to the brain—one study in nonhuman primates showed that exposure to MDMA for only 4 days caused damage to serotonin nerve terminals that was still evident 6 to 7 years later.<sup>1</sup> Although similar neurotoxicity has not been shown definitively in humans, the wealth of animal research indicating MDMA's damaging properties strongly suggests that MDMA is not a safe drug for human consumption. This is currently an area of active research.

## **Addictive Potential**

For some people, MDMA can be addictive.<sup>2</sup> A survey of young adult and adolescent MDMA users found that 43 percent of those who reported ecstasy use met the accepted diagnostic criteria for dependence, as evidenced by continued use despite knowledge of physical or psychological harm, withdrawal effects, and tolerance (or diminished response).<sup>3</sup> These results are consistent with those of similar studies in other countries that also suggest a high rate of MDMA dependence among users.<sup>4</sup> MDMA abstinence-associated withdrawal symptoms include fatigue, loss of appetite, depressed feelings, and trouble concentrating.<sup>2</sup>

## **What Other Adverse Effects does MDMA Have on Health?**

MDMA can also be dangerous to overall health and, on rare occasions, lethal. MDMA can have many of the same physical effects as other stimulants such as cocaine and amphetamines. These include increases in heart rate and blood pressure, which present risks of particular concern for people with circulatory problems or heart disease; and other symptoms such as muscle tension, involuntary teeth clenching, nausea, blurred vision, faintness, and chills or sweating.

In high doses, MDMA can interfere with the body's ability to regulate temperature. On rare but unpredictable occa-

sions, this can lead to a sharp increase in body temperature (hyperthermia), which can result in liver, kidney, and cardiovascular system failure, and death. MDMA can interfere with its own metabolism (breakdown within the body); therefore, potentially harmful levels can be reached by repeated MDMA administration within short periods of time.

Other drugs that are chemically similar to MDMA, such as MDA (methylenedioxyamphetamine, the parent drug of MDMA) and PMA (paramethoxyamphetamine, associated with fatalities in the United States and Australia),<sup>5</sup> are sometimes also sold as ecstasy. These drugs can be neurotoxic or create additional health risks to the user. Furthermore, ecstasy tablets may be adulterated with other substances, such as ephedrine (a stimulant); dextromethorphan (DXM, a cough suppressant); ketamine (an anesthetic used mostly by veterinarians); caffeine; cocaine; and methamphetamine. Although the combination of MDMA with one or more of these drugs may be inherently dangerous, users might also combine these with other substances such as marijuana and alcohol, putting themselves at even higher risk.

## **What Treatment Options Exist?**

There are no specific treatments for MDMA abuse and addiction. The most effective treatments for drug abuse and addiction in general are cognitive-

behavioral interventions that are designed to help modify the patient's thinking, expectancies, and behaviors, and to increase skills in coping with life's stressors. Drug abuse recovery support groups may be effective in combination with behavioral interventions to support long-term, drug-free recovery. There are currently no pharmacological treatments for addiction to MDMA.

**How Widespread is MDMA Abuse?**

**Monitoring the Future (MTF) Survey<sup>††</sup>**

Between 2005 and 2007, past-year abuse of MDMA increased among 12th-graders, from 3.0 percent to 4.5 percent; and between 2004 and 2007, past-year abuse of MDMA increased among 10th-graders, from 2.4 percent to 3.5 percent.

<b>MDMA Use by Students 2007 Monitoring the Future Survey</b>			
	<b>8th Grade</b>	<b>10th Grade</b>	<b>12th Grade</b>
<b>Lifetime<sup>†††</sup></b>	2.3%	5.2%	6.5%
<b>Past Year</b>	1.5	3.5	4.5
<b>Past Month</b>	0.6	1.2	1.6

For the third year in a row, the younger students surveyed reported a slightly weaker attitude about the risks associated with MDMA use. The proportion of 8th-graders who perceived risk of harm from using MDMA "occasionally" decreased significantly, from 52.0 percent in 2006 to 48.6 percent in 2007; this proportion also fell significantly among 10th-graders, from 71.3 percent in 2006 to 68.2 percent in 2007. Perceived risk of MDMA use remained unchanged for 12th-graders from 2006 to 2007.

**National Survey on Drug Use and Health (NSDUH)<sup>††††</sup>**

In 2006, an estimated 528,000 people (0.2 percent of the population) in the United States age 12 or older used MDMA in the month prior to being surveyed. Lifetime use increased among individuals aged 12 years or older, from 4.3 percent in 2002 to 5.0 percent in 2006; however, past-year use of ecstasy decreased from 1.3 percent to 0.9 percent during the same 5-year period. Approximately 860,000 Americans used ecstasy for the first time in 2006, which is a significant increase from the 615,000 first-time users reported in 2005. Most (70.1 percent) of these new users were 18 or older; and among past-year initiates aged 12 to 49, the average age at initiation in 2006 was 20.6 years.

† For more information, please visit [www.clubdrugs.org](http://www.clubdrugs.org) and [www.teens.drugabuse.gov](http://www.teens.drugabuse.gov).

For street terms searchable by drug name, street term, cost and quantities, drug trade, and drug use, visit [www.whitehousedrugpolicy.gov/streetterms/default.asp](http://www.whitehousedrugpolicy.gov/streetterms/default.asp).

†† These data are from the 2006 Monitoring the Future survey, funded by the National Institute on Drug Abuse, National Institutes of Health, Department of Health and Human Services, and conducted annually by the University of Michigan's Institute for Social Research. The survey has tracked 12th-graders' illicit drug use and related attitudes since 1975; in 1991, 8th- and 10th-graders were added to the study. The latest data are online at [www.drugabuse.gov](http://www.drugabuse.gov).

††† "Lifetime" refers to use at least once during a respondent's lifetime. "Past year" refers to use at least once during the year preceding an individual's response to the survey. "Past month" refers to use at least once during the 30 days preceding an individual's response to the survey.

†††† NSDUH (formerly known as the National Household Survey on Drug Abuse) is an annual survey of Americans age 12 and older conducted by the Substance Abuse and Mental Health Services Administration. Copies of the latest survey are available at [www.samhsa.gov](http://www.samhsa.gov) and from NIDA at 877-643-2644.

## References

<sup>1</sup> Ricaurte GA, McCann UD. Experimental studies on 3,4-methylenedioxymethamphetamine (MDMA, "ecstasy") and its potential to damage brain serotonin neurons. *Neurotox Res* 3(1):85–99, 2001.

<sup>2</sup> Stone AL, Storr CL, Anthony JC. Evidence for a hallucinogen dependence syndrome developing soon after onset of hallucinogen use during adolescence. *Int J Methods Psychiatr Res* 15:116–130, 2006.

<sup>3</sup> Cottler LB, Womack SB, Compton WM, Ben-Abdallah A. Ecstasy abuse and dependence among adolescents and young adults: Applicability and reliability of DSM-IV criteria. *Human Psychopharmacol* 16:599–606, 2001.

<sup>4</sup> Leung KS, Cottler LB. Ecstasy and other club drugs: A review of recent epidemiological studies. *Curr Opin Psychiatry* 21:234–241, 2008.

<sup>5</sup> Kraner JC, McCoy DJ, Evans MA, Evans LE, Sweeney BJ. Fatalities caused by the MDMA-related drug para-methoxyamphetamine (PMA). *J Anal Toxicol* 25(7):645–648, 2001.