

NDC 0002-3235-60 (PU3235) — Bottles of 60

NDC 0002-3235-33 (PU3235) — (ID†100) Blisters

The 30 mg* capsule has an opaque white body and opaque blue cap, and is imprinted with “30 mg” on the body and “LILLY 3240” on the cap:

NDC 0002-3240-30 (PU3240) — Bottles of 30

NDC 0002-3240-90 (PU3240) — Bottles of 90

NDC 0002-3240-04 (PU3240) — Bottles of 1000

NDC 0002-3240-33 (PU3240) — (ID†100) Blisters

The 60 mg* capsule has an opaque green body and opaque blue cap, and is imprinted with “60 mg” on the body and “LILLY 3237” on the cap:

NDC 0002-3237-30 (PU3237) — Bottles of 30

NDC 0002-3237-90 (PU3237) — Bottles of 90

NDC 0002-3237-04 (PU3237) — Bottles of 1000

NDC 0002-3237-33 (PU3237) — (ID†100) Blisters

1084 * equivalent to duloxetine base.

1085 †Identi-Dose® (unit dose medication, Lilly).

1086

1087 Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room
1088 Temperature].

1089 Literature revised June 28, 2007

**Eli Lilly and Company
Indianapolis, IN 46285, USA**

1090

1091

1092

1093

1094

www.Cymbalta.com

Copyright © 2004, 2007, Eli Lilly and Company. All rights reserved.

PV 5904 AMP

PRINTED IN USA

1095

Medication Guide

1096

Antidepressant Medicines, Depression and other Serious Mental 1097 Illnesses, and Suicidal Thoughts or Actions

1098

1099 Read the Medication Guide that comes with your or your family member’s antidepressant
1100 medicine. This Medication Guide is only about the risk of suicidal thoughts and actions with
1101 antidepressant medicines. **Talk to your, or your family member’s, healthcare provider
1102 about:**

1102

- all risks and benefits of treatment with antidepressant medicines

1103

- all treatment choices for depression or other serious mental illness

1104

What is the most important information I should know about antidepressant 1105 medicines, depression and other serious mental illnesses, and suicidal thoughts 1106 or actions?

1107

1. Antidepressant medicines may increase suicidal thoughts or actions in some
1108 children, teenagers, and young adults within the first few months of treatment.

1109 **2. Depression and other serious mental illnesses are the most important causes of**
 1110 **suicidal thoughts and actions. Some people may have a particularly high risk of**
 1111 **having suicidal thoughts or actions.** These include people who have (or have a family
 1112 history of) bipolar illness (also called manic-depressive illness) or suicidal thoughts or
 1113 actions.

1114 **3. How can I watch for and try to prevent suicidal thoughts and actions in myself or a**
 1115 **family member?**

1116 • Pay close attention to any changes, especially sudden changes, in mood, behaviors,
 1117 thoughts, or feelings. This is very important when an antidepressant medicine is
 1118 started or when the dose is changed.

1119 • Call the healthcare provider right away to report new or sudden changes in mood,
 1120 behavior, thoughts, or feelings.

1121 • Keep all follow-up visits with the healthcare provider as scheduled. Call the
 1122 healthcare provider between visits as needed, especially if you have concerns about
 1123 symptoms.

1124 **Call a healthcare provider right away if you or your family member has any of the**
 1125 **following symptoms, especially if they are new, worse, or worry you:**

1126 • thoughts about suicide or dying

1127 • attempts to commit suicide

1128 • new or worse depression

1129 • new or worse anxiety

1130 • feeling very agitated or restless

1131 • panic attacks

1132 • trouble sleeping (insomnia)

1133 • new or worse irritability

1134 • acting aggressive, being angry, or violent

1135 • acting on dangerous impulses

1136 • an extreme increase in activity and talking (mania)

1137 • other unusual changes in behavior or mood

1138 **What else do I need to know about antidepressant medicines?**

1139 • **Never stop an antidepressant medicine without first talking to a healthcare provider.**
 1140 Stopping an antidepressant medicine suddenly can cause other symptoms.

1141 • **Antidepressants are medicines used to treat depression and other illnesses.** It is
 1142 important to discuss all the risks of treating depression and also the risks of not treating it.
 1143 Patients and their families or other caregivers should discuss all treatment choices with the
 1144 healthcare provider, not just the use of antidepressants.

- 1145 • **Antidepressant medicines have other side effects.** Talk to the healthcare provider about
1146 the side effects of the medicine prescribed for you or your family member.
- 1147 • **Antidepressant medicines can interact with other medicines.** Know all of the medicines
1148 that you or your family member takes. Keep a list of all medicines to show the healthcare
1149 provider. Do not start new medicines without first checking with your healthcare provider.
- 1150 • **Not all antidepressant medicines prescribed for children are FDA approved for use in**
1151 **children.** Talk to your child's healthcare provider for more information.
- 1152 *This Medication Guide has been approved by the US Food and Drug Administration for*
1153 *all antidepressants.*
- 1154 Patient Information revised June 21, 2007

PV 5083 AMP