FRONT

AVALIDE®
(irbesartan-hydrochlorothiazide) Tablets

Rx only

GLUE PANEL

AVALIDE®
(irbesartan-hydrochlorothiazide)
Tablets

GLUE PANEL

USE IN PREGNANCY

When used in pregnancy during the second and third trimesters, drugs that act directly on the renin-angiotensin system can cause injury and even death to the developing fetus. When pregnancy is detected, AVALIDE should be discontinued as soon as possible. (See **WARNINGS: Fetal/Neonatal Morbidity and Mortality**.)

DESCRIPTION

CLINICAL PHARMACOLOGY
Mechanism of Action

INDICATIONS AND USAGE

CONTRAINDICATIONS

WARNINGS
Fetal/Neonatal Morbidity and Mortality

PRECAUTIONS

ADVERSE REACTIONS

OVERDOSAGE

DOSAGE AND ADMINISTRATION

HOW SUPPLIED

Distributed by:
Bristol-Myers Squibb Sanofi-Synthelabo Partnership
New York, NY 10016

Bristol-Myers Squibb Company   sanofi-synthelabo

BACK

GLUE AREA

VENDOR VERIFICATION I 2 OF 5 BARCODE AREA 1/2" x 1/4"

BMS 128 BARCODE AREA 1-1/2" x 1/2" WITH UP TO 7pt. HUMAN READABLE TYPE

NO TEXT AREA

IC MARK AREA 3/8"

B-MS DRAWING #: 022920A-01