



Children's
ElixSure[®]
IB

Ibuprofen Oral Suspension

Berry Flavor

Drug Facts (continued)

Ask a doctor or pharmacist before use if the child is

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if

- the child feels faint, vomits blood, or has bloody or black stools. These are signs of stomach bleeding.
- stomach pain or upset gets worse or lasts
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- do not give longer than 10 days, unless directed by a doctor (see Warnings)
- find the right dose in the chart below. If possible use weight to dose; otherwise use age.
- replace bottle cap tightly to maintain child resistance
- repeat dose every **6-8 hours**, if needed
- do not use more than **4 times a day**
- **Attention:** use this product only with the enclosed teaspoon and do not overfill. The enclosed teaspoon holds 5 mL when filled level to the top rim and 2.5 mL when leveled at the 1/2 teaspoon mark. Level by mildly shaking or tapping the spoon.

age (yrs.)	weight (lbs.)	dose (tsp.)
under 2 years	under 24	ask a doctor
2 - 3	24 - 35	1
4 - 5	36 - 47	1 1/2
6 - 8	48 - 59	2
9 - 10	60 - 71	2 1/2
11	72 - 95	3

Drug Facts (continued)

Other information

- keep carton for complete information
- store at 20°-25°C (68°-77°F)
- see bottom of carton or bottle for lot no. and exp. date
- this product is safety sealed. Do not use if seal is broken or missing.

Inactive ingredients butylparaben, carbomer 934P, FD&C yellow #6, glycerin, natural and artificial flavor, poloxamer 188, propylene glycol, purified water, sodium hydroxide, sorbitol (crystalline), sucralose liquid concentrate

Questions or comments?

Call 1-800-345-0032 (Monday through Friday, 8 am to 6 pm EST) www.elixsure.com



Distributed by Alterna LLC, Morristown, NJ 07960
©2005 Alterna LLC MADE IN CANADA.
Subject to some or all of U.S. Patent Nos. 5,881,926; 6,071,523; 6,102,254; 6,355,258; 6,399,079; 6,656,482; 6,745,919; D 483,478 and Publication Nos. 2003/0235618; 2004/0258716 and 2005/0042237.
ElixSure®, ElixSure(Spoon Design)® and NonSpill® are registered trademarks of Taro Pharmaceuticals U.S.A., Inc. used under license by Alterna LLC.

Children's
ElixSure[®]
IB

Ibuprofen Oral Suspension
(NSAID), 100 mg/5 mL

PAIN RELIEVER **FEVER REDUCER**



Spill Resistant!

See new warnings information

- Lasts up to 8 Hours
- Aspirin Free
- No Shaking Needed

Berry Flavor

4 fl oz (120 mL)

Dosage teaspoon included

Drug Facts

Active ingredient

(in each 5 mL teaspoonful)
Ibuprofen 100 mg (NSAID)*...Pain reliever/fever reducer
* nonsteroidal anti-inflammatory drug

Purposes

- relieves minor aches and pains due to:
 - the common cold
 - flu
 - sore throat
 - headaches
 - toothaches
 - reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if the child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- with feeding tubes

Ask a doctor before use if the child has

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- not been drinking fluids
- lost a lot of fluid due to vomiting or diarrhea
- high blood pressure
- heart or kidney disease
- taken a diuretic



PPK-4389-3
1005-3
M200

Lot:

Exp:



T 129



NO VARNISH

↖ ↗

LOT & EXP. CODING

Drug Facts

Active ingredient (in each 5 mL teaspoonful)
Ibuprofen 100 mg (NSAID)*. Pain reliever/fever reducer
* nonsteroidal anti-inflammatory drug

Purposes
• relieves minor aches and pains due to:
• the common cold • flu • sore throat • headaches • toothaches
• reduces fever

Uses temporarily

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
• hives • facial swelling • asthma (wheezing) • shock • skin reddening
• rash • blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if the child:
• has had stomach ulcers or bleeding problems
• takes a blood thinning (anticoagulant) or steroid drug
• takes other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
• takes more or for a longer time than directed
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use
• if the child has ever had an allergic reaction to any other pain reliever/fever reducer
• right before or after heart surgery

Drug Facts (continued)
• with feeding tubes
Ask a doctor before use if the child has
• problems or serious side effects from taking pain relievers or fever reducers
• stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
• ulcers
• bleeding problems
• not been drinking fluids
• lost a lot of fluid due to vomiting or diarrhea
• high blood pressure
• heart or kidney disease
• taken a diuretic
Ask a doctor or pharmacist before use if the child is
• taking any other drug containing an NSAID (prescription or nonprescription)
• taking a blood thinning (anticoagulant) or steroid drug
• under a doctor's care for any serious condition
• taking any other drug
When using this product
• take with food or milk; if stomach upset occurs
• long term continuous use may increase the risk of heart attack or stroke
Stop use and ask a doctor if
• the user feels faint, vomits blood, or has bloody or black stools. These are signs of stomach bleeding.
• stomach pain or upset gets worse or lasts
• the user does not get any relief within first day (24 hours) of treatment
• fever or pain gets worse or lasts more than 3 days

Drug Facts (continued)
• redness or swelling is present in the painful area
• any new symptoms appear
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
• this product does not contain directions or complete warnings for adult use
• do not give more than directed
• do not give longer than 10 days, unless directed by a doctor (see Warnings)
• find the right dose in the chart below. If possible use weight to dose; otherwise use age.
• replace bottle cap tightly to maintain child resistance
• repeat dose every 6-8 hours, if needed
• do not use more than 4 times a day
Attention: use this product only with the teaspoon provided and do not overfill. The provided teaspoon holds 5 mL when filled level to the top rim and 2.5 mL when leveled at the 1/2 teaspoon mark. Level by mildly shaking or tapping the spoon.


age (yrs)	weight (lbs.)	dose (tsp.)
under 2 years	under 24	ask a doctor
2 - 3	24 - 35	1
4 - 5	36 - 47	1 1/2
6 - 8	48 - 59	2
9 - 10	60 - 71	2 1/2
11	72 - 95	3

Other information
• keep carton for complete information • store at 20°-25°C (68°-77°F) • see bottom of bottle for lot no. and exp. date

Inactive ingredients butylparaben, carbomer 934P, FD&C yellow #6, glycerin, natural and artificial flavor, poloxamer 188, propylene glycol, purified water, sodium hydroxide, sorbitol (crystalline), sucrose liquid concentrate

Questions or comments?
Call 1-800-345-0032 (Monday through Friday, 8 am to 6 pm EST) www.elixsure.com

Scaled at 80%



Distributed by: Alterna LLC, Morristown, NJ 07960
©2005 Alterna LLC
MADE IN CANADA.
Subject to some or all of U.S. Patent Nos. 5,881,926; 6,071,523; 6,102,254; 6,355,258; 6,399,079; 6,656,482; 6,745,919; D 483,478 and Publication Nos. 2003/0235618; 2004/0258716 and 2005/0042237. ElixSure®, ElixSure(Spoon Design)® and Nonspii® are registered trademarks of Taro Pharmaceuticals U.S.A., Inc. used under license by Alterna LLC.

PPK - 4393 - 2
1105 - 2

T 147

Children's
ElixSure™
IB

Ibuprofen Oral Suspension
(NSAID), 100 mg/5 mL

PAIN RELIEVER **FEVER REDUCER**

Spill Resistant!

- Lasts up to 8 Hours
- Aspirin Free
- No Shaking Needed

Berry Flavor 

Physician Sample Only

for complete labeling information read & keep
accompanying package insert

1 fl oz (28 mL) PPK - 4392 - 2 FB 1005 - 2

Active ingredient (in each 5 mL teaspoonful)
Ibuprofen 100 mg (NSAID)*
* nonsteroidal anti-inflammatory drug

Purposes Pain reliever/fever reducer

Directions • **This product does not contain directions or complete warnings for adult use.** • **do not give more than directed**

- do not give longer than 10 days, unless directed by a doctor (see Warnings)
- find right dose in the chart below
- repeat dose every **6-8 hours**, if needed
- do not use more than **4 times a day**

Attention: use this product only with the teaspoon provided and do not overfill. The teaspoon provided holds 5 mL when filled level to top rim and 2.5 mL when leveled at the 1/2 teaspoon mark. Level by mildly shaking or tapping the spoon.

age (yrs.)	weight (lbs.)	dose (tsp.)
under 2 years	under 24	ask a doctor
2 - 3	24 - 35	1
4 - 5	36 - 47	1 1/2
6 - 8	48 - 59	2
9 - 10	60 - 71	2 1/2
11	72 - 95	3

Distributed by
Alterna LLC, Morristown, NJ 07960
©2005 Alterna LLC MADE IN CANADA.
PPK - 4392 - 2BB 1005 - 2

Children's
ElixSure[®]
IB

Ibuprofen Oral Suspension (NSAID), 100 mg/5 mL

PAIN RELIEVER FEVER REDUCER



Drug Facts

Active ingredient (in each 5 mL teaspoonful)

Ibuprofen 100 mg (NSAID)* pain reliever/fever reducer
* nonsteroidal anti-inflammatory drug

Purposes

Uses temporarily

- relieves minor aches and pains due to:
 - the common cold
 - flu
 - sore throat
 - headaches
 - toothaches
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding.

The chance is higher if the child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- with feeding tubes

Ask a doctor before use if the child has

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain

148mm

105mm

- ulcers
- bleeding problems
- not been drinking fluids
- lost a lot of fluid due to vomiting or diarrhea
- high blood pressure
- heart or kidney disease
- taken a diuretic

Ask a doctor or pharmacist before use if the child is

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if

- the child feels faint, vomits blood, or has bloody or black stools. These are signs of stomach bleeding.
- stomach pain or upset gets worse or lasts
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- do not give longer than 10 days, unless directed by a doctor (see Warnings)
- find the right dose in the chart below. If possible use weight to dose; otherwise use age.
- repeat dose every **6-8 hours**, if needed
- do not use more than **4 times a day**

Attention: use this product only with the teaspoon provided and do not overfill. The provided teaspoon holds 5 mL when filled level to the top rim and 2.5 mL when leveled at the 1/2 teaspoon mark. Level by mildly shaking or tapping the spoon.

age (yrs.)	weight (lbs.)	dose (tsp.)
under 2 years	under 24	ask a doctor
2 - 3	24 - 35	1
4 - 5	36 - 47	1 1/2
6 - 8	48 - 59	2
9 - 10	60 - 71	2 1/2
11	72 - 95	3

Other information

- keep this insert for complete information
- store at 20°-25°C (68°-77°F)
- see bottom of bottle for lot no. and exp. date

Inactive ingredients butylparaben, carbomer 934P, FD&C yellow #6, glycerin, natural and artificial flavor, poloxamer 188, propylene glycol, purified water, sodium hydroxide, sorbitol (crystalline), sucralose liquid concentrate

Questions or comments?

Call 1-800-345-0032 (Monday through Friday, 8 am to 6 pm EST)

www.elixsure.com

Distributed by: Alterra LLC, Morristown, NJ 07960

©2005 Alterra LLC

MADE IN CANADA.

PPK-5289-0 1105-0

Berry Flavor Children's

ElixSure[®] IB

Ibuprofen Oral Suspension (NSAID), 100 mg/5 mL

PAIN RELIEVER FEVER REDUCER

Directions **4 fl oz (120 mL)**

• do not give more than directed • do not give longer than 10 days, unless directed by a doctor (see Warnings) • find the right dose in the chart below. If possible, use weight to dose; otherwise use age. • replace bottle cap tightly to maintain child resistance • repeat dose every **6-8 hours**, if needed • do not use more than **4 times a day**

Attention: use this product only with the enclosed teaspoon and do not overfill. The enclosed teaspoon holds 5 mL when filled level to top rim and 2.5 mL when leveled at the 1/2 tsp. mark. Level by mildly shaking or tapping the spoon.

age (yrs.)	weight (lbs.)	dose (tsp.)
under 2 years	under 24	ask a doctor
2 - 3	24 - 35	1
4 - 5	36 - 47	1 1/2
6 - 8	48 - 59	2
9 - 10	60 - 71	2 1/2
11	72 - 95	3

PPK - 4388 - 1FB 1005 - 1

57mm

Keep carton. See carton for important new warnings and for complete information.

Active ingredient (in each 5 mL): Ibuprofen 100 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Uses temporarily: • relieves minor aches and pains due to: • the common cold • flu • sore throat • headaches • toothaches
• reduces fever

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: • hives • facial swelling • asthma (wheezing) • shock • skin reddening • rash • blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Do not use • if the child has ever had an allergic reaction to any other pain reliever/fever reducer • right before or after heart surgery • with feeding tubes

See carton for important warnings and save it for complete information.

Do not use if imprinted safety band is torn or missing. Store at 20--25°C (68--77°F). See bottom of bottle for lot no. and exp. date.

Questions or comments? Call toll free 1-800-345-0032 or ask your pharmacist, doctor or healthcare professional.

Distributed by Alterra LLC, Morristown, NJ 07960
©2005 Alterra LLC MADE IN CANADA. PPK - 4388 - 1BB 1005 - 1

54mm