

Do not use if printed foil seal under bottle cap is open or torn.

Active ingredient (in each tablet) **Purpose**
Famotidine 10 mg Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

LOT
[]

EXP.
[]

XXXXXXXXXX

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

NDC 16837-873-61

Pepcid[®]

Famotidine Tablets 10mg Acid Reducer

AC

Prevents & Relieves
Heartburn due to
Acid Indigestion

JUST ONE
PER DOSE!

60 CHEWABLE
MINT FLAVORED TABLETS

Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating, or dizziness** • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • **frequent chest pain** • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** • adults and children 12 years and over: • **do not swallow tablet whole: chew completely** • to relieve symptoms, **chew** 1 tablet before swallowing • to prevent symptoms, **chew** 1 tablet before swallowing at any time from **15 to 60 minutes** before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor. **Other information** • Phenylketonurics: Contains Phenylalanine 1.4 mg per chewable tablet

Dist. by: **Johnson & Johnson** • MERCK
CONSUMER PHARMACEUTICALS CO., FORT WASHINGTON, PA 19034 USA
©Registered trademark of Merck & Co., Inc.

150% OF SIZE

Do not use if printed foil seal under bottle cap is open or torn.

Active ingredient (in each tablet) **Purpose**
Famotidine 10 mg Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

LOT
[]

EXP.
[]

XXXXXXXXXX

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

NDC 16837-873-61

Pepcid[®]

Famotidine Tablets 10mg Acid Reducer

AC

Prevents & Relieves
Heartburn due to
Acid Indigestion

JUST ONE
PER DOSE!

60 CHEWABLE
MINT FLAVORED TABLETS

Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating, or dizziness** • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • **frequent chest pain** • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** • adults and children 12 years and over: • **do not swallow tablet whole: chew completely** • to relieve symptoms, **chew** 1 tablet before swallowing • to prevent symptoms, **chew** 1 tablet before swallowing at any time from **15 to 60 minutes** before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor. **Other information** • Phenylketonurics: Contains Phenylalanine 1.4 mg per chewable tablet

Dist. by: **Johnson & Johnson** • MERCK
CONSUMER PHARMACEUTICALS CO., FORT WASHINGTON, PA 19034 USA
©Registered trademark of Merck & Co., Inc.


Labeling Format Information:

Fonts: Universe Cond regular, bold and bold italic.

Drug Facts:	NA	Leading: 4.4-4.5 pt
Header:	NA	Bullets: 4.5 pt
Subheader:	4.5 pt	Barlines: NA
Body Text:	4.5 pt	Hairlines: NA
Drug Facts (continued):	NA	


Horizontal Scale: 73-75% Average Kerning: -15

LABEL PEPCID CCT 60 (CLASS LABELING) 3004927 C-3
OCTOBER 1, 2004



Do not use if printed foil seal under bottle cap is open or torn.

WHAT YOU SHOULD KNOW ABOUT:



PACKAGE INSERT


Chewable

A non-prescription stomach medicine

PEPCID AC Chewables contain a medicine, different from antacids, that doctors have prescribed for years to treat acid-related problems in millions of people.

Product Benefits:



- 1 chewable tablet relieves heartburn due to acid indigestion.
- Pepcid AC prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods or beverages.
- It contains famotidine, a prescription-proven medicine.



Shown actual size

How to use PEPCID AC Chewables


Adults and children 12 years and over:
Use PEPCID AC Chewables to relieve or prevent heartburn due to acid indigestion and sour stomach.

	TO RELIEVE SYMPTOMS Chew 1 tablet completely before swallowing.	PEPCID AC can be used up to twice daily (up to 2 tablets in 24 hours).
	TO PREVENT SYMPTOMS Chew 1 tablet completely before swallowing any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn.	

Excess acid: a burning problem

It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the wrong place (the esophagus), or too much acid, can cause burning pain and discomfort that interfere with everyday activities.

Heartburn— Caused by acid in the esophagus



A valve-like muscle called the lower esophageal sphincter (LES) is relaxed in an open position.

Burning pain/discomfort

Excess acid moves up into esophagus

Tips for Managing Heartburn

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

Excellent safety record

The ingredient in PEPCID AC, famotidine, has been prescribed by doctors for years to treat millions of patients safely and effectively. The active ingredient in PEPCID AC has been taken safely with many frequently prescribed medications.

Know when to see your doctor

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools.
- if you have signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- if pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Questions or comments?

Please call 1-800-755-4008 for more information and a guide of how to help your heartburn.

Distributed by: **Johanson-Johnson • MERCK**
CONSUMER PHARMACEUTICALS CO.
FORT WASHINGTON, PA 19034 USA

VALUABLE MAIL-IN OFFER

Save \$3.00 on any Pepcid Complete Same heartburn relief as Pepcid AC Starts to neutralize acid in seconds product with purchase of 25 count or larger

BUY: Any Pepcid Complete Package 25 count or larger.
SEND: This completed certificate, UPC from the Pepcid Complete package, plus dated cash register receipt with purchase price circled.
RECEIVE: \$3.00 Coupon by mail.

SEND TO: Pepcid AC / Pepcid Complete \$3 Offer 780
P.O. Box 6607
Douglas, AZ 85555-6607

NOTE: Offer good only in U.S.A. and APO/FPO addresses. This request form may not be mechanically reproduced. LIMIT TWO OFFERS PER FAMILY AND/OR ADDRESS, PER YEAR. No group or organization requests will be honored. Your offer rights may not be transferred or assigned. Offer valid where prohibited or taxed. You should expect to receive your coupon in 8-9 weeks. Cannot be combined with any other offers.

Yes, I'd like to receive future offers and information from McNeil Consumer & Specialty Pharmaceuticals and McNeil Nutritional, Divisions of McNeil-PPC, Inc. and Johnson & Johnson - Merck Consumer Pharmaceuticals Co.

Please circle one: Dr. Mr. Mrs. Miss.

Name (please print) _____
Address _____
City _____ State _____ Zip _____ Apt. _____
E-mail _____

Age: under 35 35-49 50-54 55-64 65 yrs+

Labeling Format Information:

Fonts: Universe regular and bold		
Drug Facts:	NA	Leading: 6.5 pt
Header:	NA	Bullets: 5 pt
Subheader:	6 pt	Barlines: NA
Body Text:	6 pt	Hairlines: NA
Drug Facts (continued):	NA	
Horizontal Scale: 100%		Average Kerning: 0

INSERT PEPCID CCT BOTTLE CODES (CLASS LABELING) 3004927 D-2
OCTOBER 1, 2004