

NDC XXXXX-XXXX-XX

See new warnings information.

Ibuprofen Capsules, 200 mg

Pain Reliever/Fever Reducer (NSAID)

200 Softgels*

*Liquid filled capsules

NDC XXXXX-XXXX-XX Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Drug Facts

Active ingredient (in each capsule)
Ibuprofen 200 mg (NSAID)

Uses

- temporarily relieves minor aches and pains due to:
 - muscular aches
 - minor pain of arthritis
 - backache
 - the common cold
 - menstrual cramps
 - temporarily reduces fever
- toothache

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- swelling of the face, lips, tongue, or throat
- skin redness and rash
- asthma (wheezing)
- shock
- skin reddening
- ulcers
- blisters

If an allergic reaction occurs, seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery

Ask a doctor before use if you have

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- high blood pressure
- heart or kidney disease
- taken a diuretic
- reached age 60 or older

Ask a doctor or pharmacist before use if you are

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- taking a blood thinner (aspirin, clopidogrel, or ticagrelor)
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

Manufactured by: Banner Pharmaceuticals, Inc.
4425 Premier Drive, High Point, NC 27265

Rev #10/06
Lot #
Expiration Date

SEE THE BACK

Drug Facts (continued)

When using this product:

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if:

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- adults and children 12 years and over:
 - take 1 capsule every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 capsule, 2 capsules may be used
 - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other Information

- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- do not use if tamper-evident seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing
- read all warnings and directions before use.

Inactive ingredients
FD&C Blue #1, gelatin, mannitol, pharmaceutical ink, polyethylene glycol, povidone, purified water, sorbitan, sorbitol, and Vitamin E polyethylene glycol succinate

Questions or comments? call toll free 1-800-XXX-XXXX

Customer: BANNER

Product Number: IBUPROFEN NSAID 200 CT

Revision Level: _____

Size: 3 X 7

Artist: MJ PP# 25340BANN

Proof Date: 10/2/06

Proofread: _____ Artist _____ Q.C. _____

Pharmalabel

7915 INDUSTRIAL VILLAGE ROAD, GREENSBORO, NC 27409
PHONE 336-855-7142 • FAX 336-854-3753
www.pharmalabel.com

1st Proof 2nd Proof 3rd Proof 4th Proof _____ Other _____

Revise and resubmit proof

PHARMALABEL provides this proof to represent the image prepared for you according to the instructions provided. PHARMALABEL is not responsible for errors or omissions unless clearly indicated on this proof. This proof does not accurately represent the colors that will print in the finished product. Any colors that are referenced on this proof are representative only. The finished product will be printed to match the referenced Pantone Color or any approved match standard provided by the customer.



Approved By: _____

Date: _____