


08723820



**Drug Facts**

**Active ingredient (in each geltab)**  
Acetaminophen 650 mg. Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - muscular aches
  - backache
  - headache
  - toothache
  - the common cold
  - menstrual cramps
  - minor pain of arthritis
- temporarily reduces fever

**Warnings**

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.

Do not use

- with any other product containing acetaminophen

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**TYLENOL**  
PAIN RELIEVER - FEVER REDUCER  
CONTAINS Acetaminophen  
Extended Release  
**8 Hour**  
FOR UP TO 8 HOUR RELIEF OF  
MUSCULAR ACHES AND PAINS  
80 GELTABS - 650 mg each  
Geltabs®  
\*Gelatin-Coated Tablets

8723820

EXP. 3 0045-0505-80 4

**NEW!**  
**TYLENOL**  
PAIN RELIEVER - FEVER REDUCER  
CONTAINS Acetaminophen  
Extended Release  
**8 Hour**  
FOR UP TO 8 HOUR RELIEF OF  
MUSCULAR ACHES AND PAINS  
80 GELTABS - 650 mg each  
Geltabs®  
\*Gelatin-Coated Tablets

**TYLENOL**  
**8 Hour**  
Geltabs  
McNeil  
McNeil Consumer & Specialty Pharmaceuticals  
DIVISION OF MCNEIL-PPC, INC.  
FORT WASHINGTON, PA 19034 USA  
© MCN-PPC, Inc. '02  
Visit us at [www.tylenol.com](http://www.tylenol.com) or call toll-free  
1-877-TYLENOL (1-877-895-3665)  
U.S. Patent Nos. 4,820,522 and 5,436,026

**Drug Facts (continued)**

**Directions**

- do not take more than directed (see overdose warning)
- take 2 geltabs every 8 hours with water
- swallow whole - do not crush, chew or dissolve
- do not take more than 6 geltabs in 24 hours
- do not use for more than 10 days unless directed by a doctor
- ask a doctor

children under 12 years

**Other information**

- do not use if carbon is opened or neck wrap or foil inner seal imprinted with "Safety Seal" is broken
- store at 20 - 25°C (68 - 77°F)
- avoid excessive heat 40°C (104°F)
- see end panel for lot number and expiration date

**Inactive ingredients**

benzyl alcohol, butylparaben, castor oil, cellulose, corn starch, edible calcium disodium, FD&C Blue #1, FD&C Blue #2, FD&C Red #40, gelatin, hydroxyethyl cellulose, hypromellose, magnesium stearate, methylparaben, polyethylene glycol, polypropylene glycol, sodium lauryl sulfate, sodium propionate, sodium starch glycolate, titanium dioxide

**Questions or comments?**

call toll-free 1-877-TYLENOL (1-877-895-3665)

**What makes Tylenol® 8 Hour Extended Relief Geltabs different?**

- Uses a unique, patented bi-layer tablet. The first layer dissolves quickly to provide prompt relief while the second layer is time released to provide up to 8 hours of relief.
- For more information or questions, visit our website [www.tylenol.com](http://www.tylenol.com).
- The tablets of Tylenol® do not manufacture store brands.

08723820

**Labeling Format Information:**

Fonts: Helvetica regular, bold and bold italic.	
<b>Drug Facts:</b>	9 pt
<b>Header:</b>	8 pt
<b>Subheader:</b>	6 pt
<b>Body Text:</b>	6 pt
<b>Drug Facts (continued):</b>	8 pt
<b>Leading:</b>	6.5 pt
<b>Bullets:</b>	5 pt
<b>Barlines:</b>	2.5 pt
<b>Hairlines:</b>	0.5 pt

<p><b>Active ingredient (in each geltab)</b> Purpose Acetaminophen 650 mg Pain reliever/fever reducer</p>	<p><b>READ THE LABEL</b></p> <p>NDC 50590-505-90</p>	<p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not take more than directed (see overdose warning)</li> <li>Adults and children 12 years and older:             <ul style="list-style-type: none"> <li>take 2 tablets every 8 hours with water</li> <li>swallow whole - do not crush, chew or dissolve</li> <li>do not take more than 6 tablets in 24 hours</li> <li>do not use for more than 10 days unless directed by a doctor</li> </ul> </li> <li>Children under 12 years:             <ul style="list-style-type: none"> <li>ask a doctor</li> </ul> </li> </ul>	<p>PARKEE-DAVIS &amp; COMPANY PHARMACEUTICALS DIVISION OF MCNEIL-PPC, INC.          1000 North 17th Street, P.O. Box 14147, Kansas City, MO 64114-0147          U.S. Patent Nos. 4,829,829; 4,914,488; 5,000,000          EXP. DATE CONTROL 7834070</p>
<p><b>Uses</b> temporarily relieves minor aches and pains due to:</p> <ul style="list-style-type: none"> <li>muscular aches</li> <li>backache</li> <li>headache</li> <li>toothache</li> <li>the common cold</li> <li>menstrual cramps</li> <li>minor pain of arthritis</li> </ul> <p>temporarily reduces fever</p>	<p>DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN</p> <p><b>TYLENOL</b><sup>®</sup></p> <p>PAIN RELIEVER FEVER REDUCER</p> <p><b>8 Hour</b></p> <p>CONTAINS Acetaminophen Extended Release</p> <p>FOR UP TO 8 HOUR RELIEF OF MUSCULAR ACHES AND PAINS</p> <p>60 GELTABS - 650 mg each      <i>Geltabs</i><sup>®</sup> *Gelatin-Coated Tablets</p>	<p><b>Other information</b></p> <ul style="list-style-type: none"> <li>do not use if neck wrap or foil inner seal imprinted with "Safely Seal" is broken</li> <li>store at 20 - 25°C (68 - 77°F)</li> <li>avoid excessive heat 40°C (104°F)</li> </ul> <p><b>Inactive ingredients</b> benzyl alcohol, butylparaben, castor oil, cellulose, corn starch, edetate calcium disodium, FD&amp;C Blue #1, FD&amp;C Blue #2, FD&amp;C Red #40, gelatin, hydroxyethyl cellulose, hydroxymethylcellulose, magnesium stearate, methylparaben, polydioxane, propylparaben, sodium lauryl sulfate, sodium propionate, sodium starch glycolate, titanium dioxide</p>	
<p><b>Warnings</b></p> <p><b>Alcohol warning:</b> If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.</p> <p><b>Do not use</b> with any other product containing acetaminophen. Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> <li>new symptoms occur</li> <li>redness or swelling is present</li> <li>pain gets worse or lasts for more than 10 days</li> <li>fever gets worse or lasts for more than 3 days</li> </ul> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p> <p><b>Overdose warning:</b> Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.</p>		<p><b>Questions or comments?</b> call toll-free 1-877-TYLENOL (1-877-895-3665)</p>	

<b>Labeling Format Information:</b>			
Fonts: Helvetica regular, bold and bold italic.			
Drug Facts:	NA	Leading:	5 pt
Header:	6 pt	Bullets:	4 pt
Subheader:	5 pt	Barlines:	NA
Body Text:	5 pt	Hairlines:	NA
Drug Facts (continued):	NA		
Horizontal Scale: 77-80%		Average Kerning: -10	

**LBL TY 8 HOUR GT 80      3004037 B-6**  
**DECEMBER 4, 2002**