

**Prevalence Report
for
New Body Weight Measures**

1999

TABLE OF CONTENTS
PREVALENCE REPORT FOR NEW BODY WEIGHT MEASURES

NORMAL (BMI LE 24.9)

| | | | |
|------------|--------|--|---|
| TABLE 01.1 | NORMAL | BASED ON NEW BODY MASS MEASURES..... | 1 |
| 01.2 | NORMAL | BASED ON NEW BODY MASS MEASURES - BY SEX..... | 2 |
| 01.3 | NORMAL | BASED ON NEW BODY MASS MEASURES, CHANGE 1998-1999..... | 3 |

OVERWEIGHT I (BMI GE 25.0 and LE 29.9)

| | | | |
|------------|------------|--|---|
| TABLE 02.1 | OVERWEIGHT | BASED ON NEW BODY MASS MEASURES..... | 4 |
| 02.2 | OVERWEIGHT | BASED ON NEW BODY MASS MEASURES - BY SEX..... | 5 |
| 02.3 | OVERWEIGHT | BASED ON NEW BODY MASS MEASURES, CHANGE 1998-1999..... | 6 |

OVERWEIGHT OR OBESE (BMI GE 25.0 and LE 99.8)

| | | | |
|------------|------------|--|---|
| TABLE 03.1 | OVERWEIGHT | BASED ON NEW BODY MASS MEASURES..... | 7 |
| 03.2 | OVERWEIGHT | BASED ON NEW BODY MASS MEASURES - BY SEX..... | 8 |
| 03.3 | OVERWEIGHT | BASED ON NEW BODY MASS MEASURES, CHANGE 1998-1999..... | 9 |

OBESITY 1 (BMI GE 30.0 and LE 34.9)

| | | | |
|------------|-----------|--|----|
| TABLE 04.1 | OBESITY 1 | BASED ON NEW BODY MASS MEASURES..... | 10 |
| 04.2 | OBESITY 1 | BASED ON NEW BODY MASS MEASURES - BY SEX..... | 11 |
| 04.3 | OBESITY 1 | BASED ON NEW BODY MASS MEASURES, CHANGE 1998-1999..... | 12 |

OBESITY 2 (BMI GE 35.0 and LE 39.9)

| | | | |
|------------|-----------|--|----|
| TABLE 05.1 | OBESITY 2 | BASED ON NEW BODY MASS MEASURES..... | 13 |
| 05.2 | OBESITY 2 | BASED ON NEW BODY MASS MEASURES - BY SEX..... | 14 |
| 05.3 | OBESITY 2 | BASED ON NEW BODY MASS MEASURES, CHANGE 1998-1999..... | 15 |

OBESITY 3 (BMI GE 40.0 and LE 99.8)

| | | | |
|------------|-----------|--|----|
| TABLE 06.1 | OBESITY 3 | BASED ON NEW BODY MASS MEASURES..... | 16 |
| 06.2 | OBESITY 3 | BASED ON NEW BODY MASS MEASURES - BY SEX..... | 17 |
| 06.3 | OBESITY 3 | BASED ON NEW BODY MASS MEASURES, CHANGE 1998-1999..... | 18 |

OBESE: ALL CATEGORIES COMBINED (BMI GE 30.0 and LE 99.8)

| | | | |
|------------|-------|--|----|
| TABLE 07.1 | OBESE | BASED ON NEW BODY MASS MEASURES..... | 19 |
| 07.2 | OBESE | BASED ON NEW BODY MASS MEASURES - BY SEX..... | 20 |
| 07.3 | OBESE | BASED ON NEW BODY MASS MEASURES, CHANGE 1998-1999..... | 21 |

TABLE 1.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NORMAL WEIGHT: BY NEW BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL |
|----------------|-------------|---------|----------------|-------------------------|
| ALABAMA | 2045 | 39.2 | 1.2 | (36.8, 41.5) |
| ALASKA | 2007 | 38.9 | 1.6 | (35.7, 42.1) |
| ARIZONA | 1675 | 48.4 | 1.8 | (44.9, 51.9) |
| ARKANSAS | 2908 | 41.2 | 1.0 | (39.2, 43.2) |
| CALIFORNIA | 4152 | 45.2 | 0.9 | (43.4, 46.9) |
| COLORADO | 2986 | 52.1 | 1.3 | (49.6, 54.7) |
| CONNECTICUT | 3320 | 47.4 | 1.2 | (45.0, 49.9) |
| DELAWARE | 2461 | 44.5 | 1.4 | (41.8, 47.2) |
| DISTRICT OF CO | 1242 | 49.1 | 1.6 | (46.0, 52.3) |
| FLORIDA | 5046 | 43.1 | 0.9 | (41.5, 44.8) |
| GEORGIA | 2204 | 42.4 | 1.2 | (39.9, 44.8) |
| HAWAII | 2111 | 49.9 | 1.5 | (47.0, 52.7) |
| IDAHO | 4809 | 44.8 | 0.9 | (43.0, 46.5) |
| ILLINOIS | 2806 | 43.1 | 1.1 | (41.0, 45.2) |
| INDIANA | 2370 | 43.3 | 1.7 | (39.9, 46.6) |
| IOWA | 3528 | 41.6 | 1.0 | (39.7, 43.6) |
| KANSAS | 3680 | 43.9 | 0.9 | (42.0, 45.7) |
| KENTUCKY | 7315 | 41.5 | 0.8 | (39.9, 43.1) |
| LOUISIANA | 1617 | 41.7 | 1.4 | (39.0, 44.4) |
| MAINE | 1613 | 45.2 | 1.4 | (42.4, 48.0) |
| MARYLAND | 3819 | 44.7 | 1.0 | (42.7, 46.6) |
| MASSACHUSETTS | 4689 | 49.7 | 1.0 | (47.8, 51.6) |
| MICHIGAN | 2445 | 40.4 | 1.1 | (38.3, 42.6) |
| MINNESOTA | 5155 | 44.1 | 0.7 | (42.7, 45.6) |
| MISSISSIPPI | 2078 | 37.7 | 1.2 | (35.3, 40.1) |
| MISSOURI | 4161 | 42.3 | 1.1 | (40.3, 44.4) |
| MONTANA | 1743 | 47.4 | 1.3 | (44.7, 50.0) |
| NEBRASKA | 2700 | 42.2 | 1.1 | (40.1, 44.3) |
| NEVADA | 2131 | 46.3 | 1.5 | (43.3, 49.3) |
| NEW HAMPSHIRE | 1192 | 49.9 | 1.7 | (46.6, 53.2) |
| NEW JERSEY | 2677 | 45.5 | 1.2 | (43.2, 47.8) |
| NEW MEXICO | 3378 | 44.7 | 1.0 | (42.9, 46.6) |
| NEW YORK | 2506 | 46.0 | 1.1 | (43.8, 48.3) |
| NORTH CAROLINA | 2331 | 42.1 | 1.2 | (39.8, 44.5) |
| NORTH DAKOTA | 1927 | 39.9 | 1.2 | (37.5, 42.3) |
| OHIO | 1548 | 43.1 | 1.5 | (40.1, 46.1) |
| OKLAHOMA | 2812 | 42.2 | 1.1 | (40.1, 44.2) |
| OREGON | 1741 | 45.5 | 1.3 | (43.0, 48.1) |
| PENNSYLVANIA | 3356 | 43.8 | 1.0 | (41.9, 45.7) |
| RHODE ISLAND | 3802 | 45.8 | 0.9 | (44.0, 47.6) |
| SOUTH CAROLINA | 3263 | 41.7 | 1.0 | (39.8, 43.7) |
| SOUTH DAKOTA | 4001 | 40.7 | 0.9 | (38.9, 42.4) |
| TENNESSEE | 2866 | 43.4 | 1.1 | (41.3, 45.4) |
| TEXAS | 4736 | 41.7 | 0.9 | (39.9, 43.5) |
| UTAH | 3093 | 49.1 | 1.3 | (46.6, 51.5) |
| VERMONT | 3015 | 47.7 | 1.1 | (45.6, 49.7) |
| VIRGINIA | 3372 | 41.4 | 1.3 | (38.9, 43.8) |
| WASHINGTON | 3453 | 45.6 | 1.1 | (43.5, 47.7) |
| WEST VIRGINIA | 2504 | 38.0 | 1.1 | (35.9, 40.2) |
| WISCONSIN | 2117 | 44.3 | 1.2 | (42.0, 46.7) |
| WYOMING | 2343 | 44.6 | 1.2 | (42.3, 46.9) |
| PUERTO RICO | 2894 | 41.5 | 1.1 | (39.4, 43.7) |

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 43.8 RANGE = 37.7-52.1 YEAR 2000 OBJ. 2.3
 * BODY MASS INDEX LE 24.9 FOR MALES AND FEMALES
 June 22, 2000

TABLE 1.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NORMAL WEIGHT: BY NEW BODY MASS INDEX* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | -----MALES----- | | | | * | -----FEMALES----- | | | |
|----------------|-----------------|---------|----------------|-------------------------|---|-------------------|---------|----------------|-------------------------|
| | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL | | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL |
| ALABAMA | 845 | 32.1 | 1.8 | (28.5, 35.6) | * | 1200 | 45.6 | 1.6 | (42.4, 48.8) |
| ALASKA | 997 | 31.9 | 2.1 | (27.7, 36.1) | * | 1010 | 46.9 | 2.5 | (42.0, 51.8) |
| ARIZONA | 698 | 31.7 | 2.5 | (26.9, 36.6) | * | 977 | 64.9 | 2.1 | (60.7, 69.0) |
| ARKANSAS | 1160 | 33.9 | 1.6 | (30.8, 37.0) | * | 1748 | 48.0 | 1.3 | (45.3, 50.6) |
| CALIFORNIA | 1796 | 37.6 | 1.3 | (35.0, 40.1) | * | 2356 | 52.9 | 1.2 | (50.6, 55.1) |
| COLORADO | 1359 | 43.8 | 1.9 | (40.0, 47.5) | * | 1627 | 60.2 | 1.7 | (56.9, 63.6) |
| CONNECTICUT | 1346 | 36.1 | 1.7 | (32.7, 39.5) | * | 1974 | 58.4 | 1.6 | (55.3, 61.4) |
| DELAWARE | 1018 | 35.8 | 2.1 | (31.7, 40.0) | * | 1443 | 52.9 | 1.8 | (49.3, 56.4) |
| DISTRICT OF CO | 456 | 45.2 | 2.7 | (40.0, 50.5) | * | 786 | 52.5 | 2.0 | (48.6, 56.4) |
| FLORIDA | 2029 | 34.6 | 1.3 | (32.0, 37.2) | * | 3017 | 51.0 | 1.1 | (48.9, 53.1) |
| GEORGIA | 874 | 35.7 | 1.9 | (32.1, 39.4) | * | 1330 | 48.6 | 1.6 | (45.5, 51.7) |
| HAWAII | 983 | 40.4 | 2.1 | (36.3, 44.4) | * | 1128 | 59.8 | 1.9 | (56.0, 63.5) |
| IDAHO | 2079 | 36.0 | 1.4 | (33.3, 38.7) | * | 2730 | 53.5 | 1.1 | (51.3, 55.7) |
| ILLINOIS | 1088 | 33.6 | 1.6 | (30.4, 36.8) | * | 1718 | 52.0 | 1.3 | (49.4, 54.6) |
| INDIANA | 1010 | 33.4 | 2.4 | (28.6, 38.2) | * | 1360 | 52.6 | 2.3 | (48.1, 57.0) |
| IOWA | 1409 | 31.4 | 1.5 | (28.4, 34.4) | * | 2119 | 51.1 | 1.3 | (48.6, 53.6) |
| KANSAS | 1546 | 35.0 | 1.4 | (32.2, 37.8) | * | 2134 | 52.8 | 1.2 | (50.4, 55.1) |
| KENTUCKY | 2737 | 33.6 | 1.3 | (31.1, 36.2) | * | 4578 | 48.9 | 1.0 | (46.9, 50.9) |
| LOUISIANA | 658 | 33.9 | 2.1 | (29.7, 38.0) | * | 959 | 49.0 | 1.8 | (45.4, 52.6) |
| MAINE | 731 | 36.4 | 2.1 | (32.2, 40.5) | * | 882 | 53.8 | 1.9 | (50.2, 57.4) |
| MARYLAND | 1570 | 37.8 | 1.6 | (34.7, 40.8) | * | 2249 | 51.2 | 1.3 | (48.6, 53.7) |
| MASSACHUSETTS | 1948 | 37.5 | 1.5 | (34.6, 40.4) | * | 2741 | 61.5 | 1.2 | (59.2, 63.8) |
| MICHIGAN | 1030 | 31.5 | 1.6 | (28.4, 34.6) | * | 1415 | 48.9 | 1.5 | (46.1, 51.8) |
| MINNESOTA | 2416 | 31.5 | 1.0 | (29.5, 33.5) | * | 2739 | 56.6 | 1.0 | (54.6, 58.6) |
| MISSISSIPPI | 843 | 33.6 | 1.9 | (29.8, 37.4) | * | 1235 | 41.6 | 1.6 | (38.5, 44.6) |
| MISSOURI | 1707 | 32.2 | 1.6 | (29.1, 35.2) | * | 2454 | 51.8 | 1.4 | (49.1, 54.5) |
| MONTANA | 765 | 38.5 | 2.0 | (34.6, 42.3) | * | 978 | 56.2 | 1.8 | (52.8, 59.6) |
| NEBRASKA | 1099 | 33.3 | 1.6 | (30.3, 36.4) | * | 1601 | 50.7 | 1.4 | (47.9, 53.4) |
| NEVADA | 1092 | 38.1 | 2.1 | (33.9, 42.3) | * | 1039 | 54.8 | 2.4 | (50.2, 59.4) |
| NEW HAMPSHIRE | 496 | 37.1 | 2.6 | (32.1, 42.2) | * | 696 | 62.5 | 2.1 | (58.4, 66.6) |
| NEW JERSEY | 1103 | 36.0 | 1.8 | (32.5, 39.4) | * | 1574 | 54.5 | 1.5 | (51.5, 57.4) |
| NEW MEXICO | 1509 | 37.3 | 1.4 | (34.6, 40.0) | * | 1869 | 52.0 | 1.3 | (49.4, 54.5) |
| NEW YORK | 1049 | 38.7 | 1.7 | (35.3, 42.1) | * | 1457 | 53.0 | 1.5 | (50.1, 55.9) |
| NORTH CAROLINA | 951 | 31.9 | 1.8 | (28.4, 35.4) | * | 1380 | 52.0 | 1.6 | (48.9, 55.0) |
| NORTH DAKOTA | 856 | 28.7 | 1.7 | (25.3, 32.0) | * | 1071 | 51.2 | 1.7 | (47.9, 54.5) |
| OHIO | 629 | 34.8 | 2.4 | (30.0, 39.5) | * | 919 | 51.2 | 2.0 | (47.3, 55.0) |
| OKLAHOMA | 1119 | 33.3 | 1.6 | (30.2, 36.4) | * | 1693 | 50.7 | 1.4 | (48.0, 53.3) |
| OREGON | 748 | 36.6 | 1.9 | (32.8, 40.4) | * | 993 | 54.1 | 1.7 | (50.7, 57.4) |
| PENNSYLVANIA | 1393 | 33.4 | 1.4 | (30.6, 36.2) | * | 1963 | 53.7 | 1.3 | (51.2, 56.2) |
| RHODE ISLAND | 1530 | 36.0 | 1.4 | (33.2, 38.8) | * | 2272 | 54.9 | 1.2 | (52.5, 57.2) |
| SOUTH CAROLINA | 1362 | 32.4 | 1.5 | (29.5, 35.3) | * | 1901 | 50.7 | 1.3 | (48.2, 53.2) |
| SOUTH DAKOTA | 1670 | 33.0 | 1.3 | (30.4, 35.6) | * | 2331 | 48.2 | 1.1 | (46.0, 50.4) |
| TENNESSEE | 1019 | 34.9 | 1.6 | (31.8, 38.1) | * | 1847 | 51.4 | 1.3 | (48.8, 53.9) |
| TEXAS | 1951 | 33.4 | 1.4 | (30.6, 36.2) | * | 2785 | 49.9 | 1.2 | (47.5, 52.2) |
| UTAH | 1364 | 41.4 | 2.0 | (37.5, 45.2) | * | 1729 | 56.8 | 1.6 | (53.6, 60.0) |
| VERMONT | 1260 | 37.3 | 1.6 | (34.3, 40.4) | * | 1755 | 57.9 | 1.3 | (55.2, 60.5) |
| VIRGINIA | 1396 | 32.4 | 1.8 | (28.9, 36.0) | * | 1976 | 50.2 | 1.6 | (46.9, 53.4) |
| WASHINGTON | 1580 | 36.6 | 1.7 | (33.3, 39.8) | * | 1873 | 54.8 | 1.4 | (52.0, 57.5) |
| WEST VIRGINIA | 1055 | 30.0 | 1.6 | (26.9, 33.1) | * | 1449 | 45.4 | 1.5 | (42.5, 48.3) |
| WISCONSIN | 915 | 32.7 | 1.9 | (29.0, 36.4) | * | 1202 | 55.6 | 1.6 | (52.5, 58.7) |
| WYOMING | 1059 | 35.6 | 1.6 | (32.4, 38.8) | * | 1284 | 53.7 | 1.6 | (50.5, 56.8) |
| PUERTO RICO | 1094 | 35.9 | 1.7 | (32.6, 39.3) | * | 1800 | 46.7 | 1.4 | (44.0, 49.4) |

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 34.8
 RANGE = 28.7-45.2

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 52.2
 RANGE = 41.6-64.9

* BODY MASS INDEX LE 24.9 FOR MALES AND FEMALES

TABLE 1.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
PREVALENCE DIFFERENCES FROM 1998 TO 1999

NORMAL WEIGHT: BY NEW BODY MASS INDEX*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | 1998 PERCENT | 1999 PERCENT | PERCENT DIFFERENCE |
|----------------|-----------------|-----------------|-----------------------|
| ALABAMA | 40.4 | 39.2 | -1.2 |
| ALASKA | 41.7 | 38.9 | -2.8 |
| ARIZONA | 50.5 | 48.4 | -2.1 |
| ARKANSAS | 45.1 | 41.2 | -3.9 |
| CALIFORNIA | 46.4 | 45.2 | -1.3 |
| COLORADO | 50.7 | 52.1 | 1.4 |
| CONNECTICUT | 50.2 | 47.4 | -2.7 |
| DELAWARE | 46.8 | 44.5 | -2.3 |
| DISTRICT OF CO | 49.3 | 49.1 | -0.2 |
| FLORIDA | 46.5 | 43.1 | -3.4 |
| GEORGIA | 45.4 | 42.4 | -3.1 |
| HAWAII | 52.3 | 49.9 | -2.4 |
| IDAHO | 46.9 | 44.8 | -2.1 |
| ILLINOIS | 44.9 | 43.1 | -1.8 |
| INDIANA | 43.8 | 43.3 | -0.5 |
| IOWA | 43.3 | 41.6 | -1.7 |
| KANSAS | 45.3 | 43.9 | -1.4 |
| KENTUCKY | 41.9 | 41.5 | -0.4 |
| LOUISIANA | 43.6 | 41.7 | -1.9 |
| MAINE | 45.6 | 45.2 | -0.4 |
| MARYLAND | 44.5 | 44.7 | 0.1 |
| MASSACHUSETTS | 50.1 | 49.7 | -0.4 |
| MICHIGAN | 42.6 | 40.4 | -2.1 |
| MINNESOTA | 40.5 | 44.1 | 3.7 |
| MISSISSIPPI | 40.7 | 37.7 | -3.0 |
| MISSOURI | 42.9 | 42.3 | -0.6 |
| MONTANA | 48.5 | 47.4 | -1.2 |
| NEBRASKA | 45.9 | 42.2 | -3.7 |
| NEVADA | 49.9 | 46.3 | -3.5 |
| NEW HAMPSHIRE | 48.1 | 49.9 | 1.8 |
| NEW JERSEY | 45.6 | 45.5 | -0.1 |
| NEW MEXICO | 48.3 | 44.7 | -3.6 |
| NEW YORK | 48.7 | 46.0 | -2.6 |
| NORTH CAROLINA | 43.9 | 42.1 | -1.8 |
| NORTH DAKOTA | 42.9 | 39.9 | -3.0 |
| OHIO | 44.9 | 43.1 | -1.9 |
| OKLAHOMA | 47.0 | 42.2 | -4.8 |
| OREGON | 45.4 | 45.5 | 0.1 |
| PENNSYLVANIA | 41.5 | 43.8 | 2.3 |
| RHODE ISLAND | 47.6 | 45.8 | -1.9 |
| SOUTH CAROLINA | 45.0 | 41.7 | -3.3 |
| SOUTH DAKOTA | 48.5 | 40.7 | -7.9 |
| TENNESSEE | 45.6 | 43.4 | -2.2 |
| TEXAS | 42.7 | 41.7 | -1.0 |
| UTAH | 50.7 | 49.1 | -1.7 |
| VERMONT | 52.1 | 47.7 | -4.4 |
| VIRGINIA | 46.9 | 41.4 | -5.5 |
| WASHINGTON | 45.5 | 45.6 | 0.1 |
| WEST VIRGINIA | 42.8 | 38.0 | -4.8 |
| WISCONSIN | 43.5 | 44.3 | 0.8 |
| WYOMING | 48.5 | 44.6 | -3.9 |
| PUERTO RICO | 43.8 | 41.5 | -2.3 |

* BODY MASS INDEX LE 24.9 FOR MALES AND FEMALES
N/A - NOT AVAILABLE

TABLE 2.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT 1: BY NEW BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL |
|----------------|-------------|---------|----------------|-------------------------|
| ALABAMA | 2045 | 38.4 | 1.2 | (36.0, 40.8) |
| ALASKA | 2007 | 40.7 | 1.7 | (37.3, 44.0) |
| ARIZONA | 1675 | 39.3 | 1.8 | (35.8, 42.8) |
| ARKANSAS | 2908 | 36.2 | 1.0 | (34.2, 38.1) |
| CALIFORNIA | 4152 | 36.1 | 0.9 | (34.4, 37.8) |
| COLORADO | 2986 | 33.0 | 1.2 | (30.6, 35.4) |
| CONNECTICUT | 3320 | 37.4 | 1.2 | (35.1, 39.7) |
| DELAWARE | 2461 | 37.9 | 1.3 | (35.3, 40.6) |
| DISTRICT OF CO | 1242 | 32.4 | 1.5 | (29.4, 35.3) |
| FLORIDA | 5046 | 38.3 | 0.8 | (36.6, 39.9) |
| GEORGIA | 2204 | 36.5 | 1.2 | (34.1, 38.9) |
| HAWAII | 2111 | 34.5 | 1.4 | (31.7, 37.2) |
| IDAHO | 4809 | 35.3 | 0.9 | (33.6, 36.9) |
| ILLINOIS | 2806 | 36.0 | 1.1 | (33.9, 38.1) |
| INDIANA | 2370 | 36.9 | 1.7 | (33.5, 40.2) |
| IOWA | 3528 | 36.9 | 1.0 | (35.0, 38.8) |
| KANSAS | 3680 | 37.2 | 0.9 | (35.5, 39.0) |
| KENTUCKY | 7315 | 36.9 | 0.8 | (35.3, 38.4) |
| LOUISIANA | 1617 | 36.0 | 1.4 | (33.3, 38.7) |
| MAINE | 1613 | 35.4 | 1.3 | (32.8, 37.9) |
| MARYLAND | 3819 | 37.2 | 1.0 | (35.2, 39.1) |
| MASSACHUSETTS | 4689 | 35.6 | 0.9 | (33.8, 37.3) |
| MICHIGAN | 2445 | 36.8 | 1.1 | (34.6, 39.0) |
| MINNESOTA | 5155 | 40.4 | 0.8 | (38.9, 41.9) |
| MISSISSIPPI | 2078 | 39.1 | 1.2 | (36.7, 41.6) |
| MISSOURI | 4161 | 36.0 | 1.0 | (34.0, 38.0) |
| MONTANA | 1743 | 36.8 | 1.3 | (34.3, 39.3) |
| NEBRASKA | 2700 | 36.8 | 1.0 | (34.8, 38.9) |
| NEVADA | 2131 | 37.8 | 1.5 | (34.8, 40.8) |
| NEW HAMPSHIRE | 1192 | 35.5 | 1.6 | (32.4, 38.7) |
| NEW JERSEY | 2677 | 37.5 | 1.1 | (35.3, 39.7) |
| NEW MEXICO | 3378 | 37.6 | 0.9 | (35.7, 39.4) |
| NEW YORK | 2506 | 36.5 | 1.1 | (34.4, 38.7) |
| NORTH CAROLINA | 2331 | 36.4 | 1.2 | (34.1, 38.7) |
| NORTH DAKOTA | 1927 | 38.2 | 1.2 | (35.8, 40.6) |
| OHIO | 1548 | 36.7 | 1.5 | (33.8, 39.5) |
| OKLAHOMA | 2812 | 36.8 | 1.1 | (34.6, 38.9) |
| OREGON | 1741 | 34.6 | 1.3 | (32.1, 37.1) |
| PENNSYLVANIA | 3356 | 35.9 | 0.9 | (34.1, 37.8) |
| RHODE ISLAND | 3802 | 37.4 | 0.9 | (35.6, 39.2) |
| SOUTH CAROLINA | 3263 | 37.6 | 1.0 | (35.7, 39.5) |
| SOUTH DAKOTA | 4001 | 39.7 | 0.9 | (38.0, 41.4) |
| TENNESSEE | 2866 | 36.1 | 1.1 | (34.0, 38.1) |
| TEXAS | 4736 | 36.7 | 0.9 | (34.9, 38.4) |
| UTAH | 3093 | 34.2 | 1.2 | (31.9, 36.5) |
| VERMONT | 3015 | 34.4 | 1.0 | (32.4, 36.3) |
| VIRGINIA | 3372 | 39.4 | 1.3 | (36.9, 41.9) |
| WASHINGTON | 3453 | 36.2 | 1.0 | (34.2, 38.1) |
| WEST VIRGINIA | 2504 | 37.4 | 1.1 | (35.2, 39.6) |
| WISCONSIN | 2117 | 35.7 | 1.1 | (33.5, 38.0) |
| WYOMING | 2343 | 38.5 | 1.2 | (36.2, 40.8) |
| PUERTO RICO | 2894 | 37.2 | 1.1 | (35.1, 39.2) |

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 36.8 RANGE = 32.4-40.7 YEAR 2000 OBJ. 2.3
 * BODY MASS INDEX GE 25.0 AND LE 29.9 FOR MALES AND FEMALES June 22, 2000

TABLE 2.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT I: BY NEW BODY MASS INDEX* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | -----MALES----- | | | | * | -----FEMALES----- | | | | * |
|----------------|-----------------|---------|----------------|-------------------------|---|-------------------|---------|----------------|-------------------------|---|
| | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL | | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL | |
| ALABAMA | 845 | 46.6 | 2.0 | (42.8, 50.4) | * | 1200 | 31.0 | 1.5 | (28.0, 33.9) | * |
| ALASKA | 997 | 47.6 | 2.3 | (43.1, 52.1) | * | 1010 | 32.8 | 2.5 | (27.9, 37.8) | * |
| ARIZONA | 698 | 54.2 | 2.7 | (48.8, 59.5) | * | 977 | 24.7 | 2.0 | (20.9, 28.5) | * |
| ARKANSAS | 1160 | 43.8 | 1.6 | (40.6, 46.9) | * | 1748 | 29.1 | 1.2 | (26.7, 31.5) | * |
| CALIFORNIA | 1796 | 44.7 | 1.3 | (42.1, 47.3) | * | 2356 | 27.5 | 1.1 | (25.4, 29.5) | * |
| COLORADO | 1359 | 40.9 | 1.9 | (37.2, 44.6) | * | 1627 | 25.3 | 1.5 | (22.3, 28.3) | * |
| CONNECTICUT | 1346 | 48.5 | 1.8 | (45.0, 52.0) | * | 1974 | 26.7 | 1.3 | (24.1, 29.3) | * |
| DELAWARE | 1018 | 46.8 | 2.1 | (42.7, 51.0) | * | 1443 | 29.4 | 1.6 | (26.2, 32.6) | * |
| DISTRICT OF CO | 456 | 39.4 | 2.6 | (34.4, 44.4) | * | 786 | 26.2 | 1.7 | (23.0, 29.5) | * |
| FLORIDA | 2029 | 47.0 | 1.4 | (44.3, 49.7) | * | 3017 | 30.1 | 1.0 | (28.2, 32.1) | * |
| GEORGIA | 874 | 41.8 | 2.0 | (38.0, 45.7) | * | 1330 | 31.5 | 1.5 | (28.6, 34.4) | * |
| HAWAII | 983 | 43.3 | 2.1 | (39.1, 47.5) | * | 1128 | 25.2 | 1.7 | (21.8, 28.6) | * |
| IDAHO | 2079 | 42.3 | 1.4 | (39.6, 45.0) | * | 2730 | 28.3 | 1.0 | (26.3, 30.2) | * |
| ILLINOIS | 1088 | 44.9 | 1.7 | (41.5, 48.3) | * | 1718 | 27.7 | 1.2 | (25.3, 30.1) | * |
| INDIANA | 1010 | 47.5 | 2.6 | (42.4, 52.7) | * | 1360 | 26.8 | 2.0 | (23.0, 30.6) | * |
| IOWA | 1409 | 45.5 | 1.6 | (42.5, 48.6) | * | 2119 | 28.9 | 1.2 | (26.6, 31.2) | * |
| KANSAS | 1546 | 45.5 | 1.4 | (42.7, 48.3) | * | 2134 | 29.0 | 1.1 | (26.8, 31.2) | * |
| KENTUCKY | 2737 | 44.2 | 1.3 | (41.5, 46.8) | * | 4578 | 30.0 | 0.9 | (28.2, 31.8) | * |
| LOUISIANA | 658 | 43.6 | 2.2 | (39.3, 47.8) | * | 959 | 28.9 | 1.7 | (25.6, 32.2) | * |
| MAINE | 731 | 43.5 | 2.1 | (39.5, 47.5) | * | 882 | 27.4 | 1.6 | (24.2, 30.6) | * |
| MARYLAND | 1570 | 45.8 | 1.6 | (42.7, 48.9) | * | 2249 | 29.0 | 1.2 | (26.6, 31.4) | * |
| MASSACHUSETTS | 1948 | 45.0 | 1.5 | (42.2, 47.9) | * | 2741 | 26.5 | 1.1 | (24.4, 28.5) | * |
| MICHIGAN | 1030 | 45.1 | 1.8 | (41.6, 48.6) | * | 1415 | 28.9 | 1.3 | (26.3, 31.5) | * |
| MINNESOTA | 2416 | 51.3 | 1.1 | (49.1, 53.5) | * | 2739 | 29.6 | 0.9 | (27.8, 31.5) | * |
| MISSISSIPPI | 843 | 45.3 | 2.0 | (41.4, 49.3) | * | 1235 | 33.3 | 1.5 | (30.4, 36.2) | * |
| MISSOURI | 1707 | 45.7 | 1.7 | (42.5, 48.9) | * | 2454 | 27.0 | 1.2 | (24.7, 29.3) | * |
| MONTANA | 765 | 44.2 | 2.0 | (40.3, 48.1) | * | 978 | 29.5 | 1.6 | (26.4, 32.5) | * |
| NEBRASKA | 1099 | 43.1 | 1.6 | (39.9, 46.3) | * | 1601 | 30.8 | 1.3 | (28.3, 33.3) | * |
| NEVADA | 1092 | 47.8 | 2.2 | (43.4, 52.1) | * | 1039 | 27.5 | 2.3 | (23.0, 32.0) | * |
| NEW HAMPSHIRE | 496 | 46.3 | 2.6 | (41.3, 51.3) | * | 696 | 24.9 | 1.9 | (21.1, 28.6) | * |
| NEW JERSEY | 1103 | 44.5 | 1.8 | (41.1, 48.0) | * | 1574 | 30.8 | 1.4 | (28.1, 33.5) | * |
| NEW MEXICO | 1509 | 45.9 | 1.4 | (43.1, 48.7) | * | 1869 | 29.4 | 1.2 | (27.1, 31.7) | * |
| NEW YORK | 1049 | 43.4 | 1.7 | (40.0, 46.7) | * | 1457 | 30.1 | 1.4 | (27.4, 32.8) | * |
| NORTH CAROLINA | 951 | 47.2 | 1.9 | (43.5, 50.8) | * | 1380 | 26.0 | 1.3 | (23.4, 28.6) | * |
| NORTH DAKOTA | 856 | 47.8 | 1.9 | (44.2, 51.5) | * | 1071 | 28.6 | 1.5 | (25.6, 31.5) | * |
| OHIO | 629 | 46.4 | 2.4 | (41.8, 51.0) | * | 919 | 27.3 | 1.7 | (23.8, 30.7) | * |
| OKLAHOMA | 1119 | 45.1 | 1.7 | (41.7, 48.4) | * | 1693 | 28.9 | 1.3 | (26.3, 31.4) | * |
| OREGON | 748 | 43.3 | 2.0 | (39.4, 47.2) | * | 993 | 26.2 | 1.5 | (23.2, 29.2) | * |
| PENNSYLVANIA | 1393 | 45.7 | 1.5 | (42.8, 48.7) | * | 1963 | 26.6 | 1.1 | (24.4, 28.8) | * |
| RHODE ISLAND | 1530 | 46.4 | 1.5 | (43.5, 49.2) | * | 2272 | 29.1 | 1.1 | (26.9, 31.3) | * |
| SOUTH CAROLINA | 1362 | 46.0 | 1.5 | (43.0, 49.0) | * | 1901 | 29.6 | 1.2 | (27.3, 31.8) | * |
| SOUTH DAKOTA | 1670 | 46.7 | 1.4 | (44.0, 49.4) | * | 2331 | 32.8 | 1.1 | (30.7, 34.8) | * |
| TENNESSEE | 1019 | 45.2 | 1.7 | (41.8, 48.6) | * | 1847 | 27.5 | 1.2 | (25.1, 29.8) | * |
| TEXAS | 1951 | 44.2 | 1.4 | (41.4, 46.9) | * | 2785 | 29.3 | 1.1 | (27.1, 31.5) | * |
| UTAH | 1364 | 41.3 | 1.9 | (37.7, 45.0) | * | 1729 | 27.0 | 1.5 | (24.1, 29.9) | * |
| VERMONT | 1260 | 42.7 | 1.6 | (39.6, 45.8) | * | 1755 | 26.1 | 1.2 | (23.8, 28.4) | * |
| VIRGINIA | 1396 | 48.8 | 2.0 | (44.9, 52.7) | * | 1976 | 30.2 | 1.5 | (27.3, 33.0) | * |
| WASHINGTON | 1580 | 44.5 | 1.6 | (41.4, 47.6) | * | 1873 | 27.7 | 1.3 | (25.2, 30.2) | * |
| WEST VIRGINIA | 1055 | 45.7 | 1.7 | (42.3, 49.1) | * | 1449 | 29.7 | 1.4 | (27.0, 32.3) | * |
| WISCONSIN | 915 | 44.6 | 1.8 | (41.0, 48.2) | * | 1202 | 27.2 | 1.4 | (24.5, 30.0) | * |
| WYOMING | 1059 | 48.3 | 1.7 | (44.9, 51.6) | * | 1284 | 28.7 | 1.5 | (25.8, 31.5) | * |
| PUERTO RICO | 1094 | 42.3 | 1.7 | (39.0, 45.7) | * | 1800 | 32.4 | 1.3 | (29.9, 34.8) | * |

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 45.3
 RANGE = 39.4-54.2

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 28.9
 RANGE = 24.7-33.3

* BODY MASS INDEX GE 25.0 AND LE 29.9 FOR MALES AND FEMALES

TABLE 2.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
PREVALENCE DIFFERENCES FROM 1998 TO 1999

OVERWEIGHT I: BY NEW BODY MASS INDEX*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | 1998 PERCENT | 1999 PERCENT | PERCENT DIFFERENCE |
|----------------|-----------------|-----------------|-----------------------|
| ALABAMA | 38.3 | 38.4 | 0.1 |
| ALASKA | 36.9 | 40.7 | 3.8 |
| ARIZONA | 36.4 | 39.3 | 3.0 |
| ARKANSAS | 35.1 | 36.2 | 1.1 |
| CALIFORNIA | 36.3 | 36.1 | -0.2 |
| COLORADO | 34.9 | 33.0 | -1.9 |
| CONNECTICUT | 34.3 | 37.4 | 3.1 |
| DELAWARE | 36.0 | 37.9 | 1.9 |
| DISTRICT OF CO | 30.5 | 32.4 | 1.9 |
| FLORIDA | 35.5 | 38.3 | 2.7 |
| GEORGIA | 35.4 | 36.5 | 1.1 |
| HAWAII | 32.2 | 34.5 | 2.3 |
| IDAHO | 36.7 | 35.3 | -1.4 |
| ILLINOIS | 36.6 | 36.0 | -0.6 |
| INDIANA | 36.3 | 36.9 | 0.5 |
| IOWA | 36.9 | 36.9 | 0.0 |
| KANSAS | 37.1 | 37.2 | 0.2 |
| KENTUCKY | 37.7 | 36.9 | -0.8 |
| LOUISIANA | 34.6 | 36.0 | 1.4 |
| MAINE | 37.0 | 35.4 | -1.6 |
| MARYLAND | 35.0 | 37.2 | 2.1 |
| MASSACHUSETTS | 35.6 | 35.6 | -0.1 |
| MICHIGAN | 36.3 | 36.8 | 0.5 |
| MINNESOTA | 43.3 | 40.4 | -3.0 |
| MISSISSIPPI | 36.5 | 39.1 | 2.6 |
| MISSOURI | 36.6 | 36.0 | -0.6 |
| MONTANA | 36.4 | 36.8 | 0.4 |
| NEBRASKA | 35.7 | 36.8 | 1.1 |
| NEVADA | 36.2 | 37.8 | 1.7 |
| NEW HAMPSHIRE | 36.3 | 35.5 | -0.8 |
| NEW JERSEY | 38.9 | 37.5 | -1.4 |
| NEW MEXICO | 36.5 | 37.6 | 1.0 |
| NEW YORK | 35.0 | 36.5 | 1.5 |
| NORTH CAROLINA | 36.7 | 36.4 | -0.3 |
| NORTH DAKOTA | 37.9 | 38.2 | 0.3 |
| OHIO | 35.1 | 36.7 | 1.6 |
| OKLAHOMA | 33.5 | 36.8 | 3.2 |
| OREGON | 36.4 | 34.6 | -1.8 |
| PENNSYLVANIA | 39.2 | 35.9 | -3.3 |
| RHODE ISLAND | 35.5 | 37.4 | 1.9 |
| SOUTH CAROLINA | 34.4 | 37.6 | 3.2 |
| SOUTH DAKOTA | 35.7 | 39.7 | 4.0 |
| TENNESSEE | 35.2 | 36.1 | 0.9 |
| TEXAS | 37.1 | 36.7 | -0.4 |
| UTAH | 33.4 | 34.2 | 0.8 |
| VERMONT | 33.1 | 34.4 | 1.2 |
| VIRGINIA | 34.4 | 39.4 | 5.0 |
| WASHINGTON | 36.4 | 36.2 | -0.2 |
| WEST VIRGINIA | 33.3 | 37.4 | 4.1 |
| WISCONSIN | 38.1 | 35.7 | -2.4 |
| WYOMING | 36.4 | 38.5 | 2.1 |
| PUERTO RICO | 36.9 | 37.2 | 0.3 |

* BODY MASS INDEX GE 25.0 AND LE 29.9 FOR MALES AND FEMALES
N/A - NOT AVAILABLE

TABLE 3.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT OR OBESE: BY NEW BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL |
|----------------|-------------|---------|----------------|-------------------------|
| ALABAMA | 2045 | 60.9 | 1.2 | (58.5, 63.2) |
| ALASKA | 2007 | 61.1 | 1.6 | (57.9, 64.3) |
| ARIZONA | 1675 | 51.6 | 1.8 | (48.1, 55.1) |
| ARKANSAS | 2908 | 58.8 | 1.0 | (56.8, 60.8) |
| CALIFORNIA | 4152 | 54.8 | 0.9 | (53.1, 56.6) |
| COLORADO | 2986 | 47.9 | 1.3 | (45.3, 50.4) |
| CONNECTICUT | 3320 | 52.6 | 1.2 | (50.1, 55.0) |
| DELAWARE | 2461 | 55.5 | 1.4 | (52.8, 58.2) |
| DISTRICT OF CO | 1242 | 50.9 | 1.6 | (47.7, 54.0) |
| FLORIDA | 5046 | 56.9 | 0.9 | (55.2, 58.5) |
| GEORGIA | 2204 | 57.6 | 1.2 | (55.2, 60.1) |
| HAWAII | 2111 | 50.2 | 1.5 | (47.3, 53.0) |
| IDAHO | 4809 | 55.2 | 0.9 | (53.5, 57.0) |
| ILLINOIS | 2806 | 56.9 | 1.1 | (54.8, 59.0) |
| INDIANA | 2370 | 56.7 | 1.7 | (53.4, 60.1) |
| IOWA | 3528 | 58.4 | 1.0 | (56.4, 60.3) |
| KANSAS | 3680 | 56.1 | 0.9 | (54.3, 58.0) |
| KENTUCKY | 7315 | 58.5 | 0.8 | (56.9, 60.1) |
| LOUISIANA | 1617 | 58.3 | 1.4 | (55.6, 61.0) |
| MAINE | 1613 | 54.8 | 1.4 | (52.0, 57.6) |
| MARYLAND | 3819 | 55.4 | 1.0 | (53.4, 57.3) |
| MASSACHUSETTS | 4689 | 50.3 | 1.0 | (48.4, 52.2) |
| MICHIGAN | 2445 | 59.6 | 1.1 | (57.4, 61.7) |
| MINNESOTA | 5155 | 55.9 | 0.7 | (54.4, 57.3) |
| MISSISSIPPI | 2078 | 62.3 | 1.2 | (59.9, 64.7) |
| MISSOURI | 4161 | 57.7 | 1.1 | (55.6, 59.7) |
| MONTANA | 1743 | 52.7 | 1.3 | (50.0, 55.3) |
| NEBRASKA | 2700 | 57.8 | 1.1 | (55.7, 59.9) |
| NEVADA | 2131 | 53.7 | 1.5 | (50.7, 56.7) |
| NEW HAMPSHIRE | 1192 | 50.1 | 1.7 | (46.8, 53.4) |
| NEW JERSEY | 2677 | 54.5 | 1.2 | (52.2, 56.8) |
| NEW MEXICO | 3378 | 55.3 | 1.0 | (53.4, 57.1) |
| NEW YORK | 2506 | 54.0 | 1.1 | (51.7, 56.2) |
| NORTH CAROLINA | 2331 | 57.9 | 1.2 | (55.5, 60.2) |
| NORTH DAKOTA | 1927 | 60.1 | 1.2 | (57.7, 62.5) |
| OHIO | 1548 | 56.9 | 1.5 | (53.9, 59.9) |
| OKLAHOMA | 2812 | 57.8 | 1.1 | (55.8, 59.9) |
| OREGON | 1741 | 54.5 | 1.3 | (51.9, 57.0) |
| PENNSYLVANIA | 3356 | 56.2 | 1.0 | (54.3, 58.1) |
| RHODE ISLAND | 3802 | 54.2 | 0.9 | (52.4, 56.0) |
| SOUTH CAROLINA | 3263 | 58.3 | 1.0 | (56.3, 60.2) |
| SOUTH DAKOTA | 4001 | 59.3 | 0.9 | (57.6, 61.1) |
| TENNESSEE | 2866 | 56.6 | 1.1 | (54.6, 58.7) |
| TEXAS | 4736 | 58.3 | 0.9 | (56.5, 60.1) |
| UTAH | 3093 | 50.9 | 1.3 | (48.5, 53.4) |
| VERMONT | 3015 | 52.3 | 1.1 | (50.3, 54.4) |
| VIRGINIA | 3372 | 58.7 | 1.3 | (56.2, 61.1) |
| WASHINGTON | 3453 | 54.4 | 1.1 | (52.3, 56.5) |
| WEST VIRGINIA | 2504 | 62.0 | 1.1 | (59.8, 64.1) |
| WISCONSIN | 2117 | 55.7 | 1.2 | (53.3, 58.0) |
| WYOMING | 2343 | 55.4 | 1.2 | (53.1, 57.7) |
| PUERTO RICO | 2894 | 58.5 | 1.1 | (56.3, 60.6) |

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 56.2 RANGE = 47.9-62.3 YEAR 2000 OBJ. 2.3
 * BODY MASS INDEX GE 25.0 AND LE 99.8 FOR MALES AND FEMALES
 June 22, 2000

TABLE 3.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA 8
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT OR OBESE: BY NEW BODY MASS INDEX* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | -----MALES----- | | | | -----FEMALES----- | | | |
|----------------|-----------------|---------|----------------|-------------------------|-------------------|---------|----------------|-------------------------|
| | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL |
| ALABAMA | 845 | 67.9 | 1.8 | (64.4, 71.5) | 1200 | 54.4 | 1.6 | (51.2, 57.6) |
| ALASKA | 997 | 68.1 | 2.1 | (63.9, 72.3) | 1010 | 53.1 | 2.5 | (48.2, 58.0) |
| ARIZONA | 698 | 68.3 | 2.5 | (63.4, 73.1) | 977 | 35.1 | 2.1 | (31.0, 39.3) |
| ARKANSAS | 1160 | 66.1 | 1.6 | (63.0, 69.2) | 1748 | 52.1 | 1.3 | (49.4, 54.7) |
| CALIFORNIA | 1796 | 62.5 | 1.3 | (59.9, 65.0) | 2356 | 47.2 | 1.2 | (44.9, 49.4) |
| COLORADO | 1359 | 56.2 | 1.9 | (52.5, 60.0) | 1627 | 39.8 | 1.7 | (36.4, 43.1) |
| CONNECTICUT | 1346 | 63.9 | 1.7 | (60.5, 67.3) | 1974 | 41.6 | 1.6 | (38.6, 44.7) |
| DELAWARE | 1018 | 64.2 | 2.1 | (60.0, 68.3) | 1443 | 47.1 | 1.8 | (43.6, 50.7) |
| DISTRICT OF CO | 456 | 54.8 | 2.7 | (49.5, 60.0) | 786 | 47.5 | 2.0 | (43.6, 51.4) |
| FLORIDA | 2029 | 65.4 | 1.3 | (62.8, 68.0) | 3017 | 49.0 | 1.1 | (46.9, 51.1) |
| GEORGIA | 874 | 64.3 | 1.9 | (60.6, 67.9) | 1330 | 51.4 | 1.6 | (48.3, 54.5) |
| HAWAII | 983 | 59.7 | 2.1 | (55.6, 63.7) | 1128 | 40.2 | 1.9 | (36.5, 44.0) |
| IDAHO | 2079 | 64.0 | 1.4 | (61.3, 66.7) | 2730 | 46.5 | 1.1 | (44.3, 48.7) |
| ILLINOIS | 1088 | 66.4 | 1.6 | (63.2, 69.6) | 1718 | 48.0 | 1.3 | (45.4, 50.6) |
| INDIANA | 1010 | 66.6 | 2.4 | (61.8, 71.4) | 1360 | 47.5 | 2.3 | (43.0, 51.9) |
| IOWA | 1409 | 68.6 | 1.5 | (65.6, 71.6) | 2119 | 48.9 | 1.3 | (46.4, 51.4) |
| KANSAS | 1546 | 65.0 | 1.4 | (62.2, 67.8) | 2134 | 47.2 | 1.2 | (44.9, 49.6) |
| KENTUCKY | 2737 | 66.4 | 1.3 | (63.8, 68.9) | 4578 | 51.1 | 1.0 | (49.1, 53.1) |
| LOUISIANA | 658 | 66.1 | 2.1 | (62.0, 70.3) | 959 | 51.0 | 1.8 | (47.4, 54.6) |
| MAINE | 731 | 63.7 | 2.1 | (59.5, 67.8) | 882 | 46.2 | 1.9 | (42.6, 49.8) |
| MARYLAND | 1570 | 62.2 | 1.6 | (59.2, 65.3) | 2249 | 48.9 | 1.3 | (46.3, 51.4) |
| MASSACHUSETTS | 1948 | 62.5 | 1.5 | (59.6, 65.4) | 2741 | 38.5 | 1.2 | (36.2, 40.8) |
| MICHIGAN | 1030 | 68.5 | 1.6 | (65.4, 71.6) | 1415 | 51.1 | 1.5 | (48.2, 53.9) |
| MINNESOTA | 2416 | 68.5 | 1.0 | (66.5, 70.5) | 2739 | 43.4 | 1.0 | (41.4, 45.4) |
| MISSISSIPPI | 843 | 66.4 | 1.9 | (62.6, 70.2) | 1235 | 58.4 | 1.6 | (55.4, 61.5) |
| MISSOURI | 1707 | 67.8 | 1.6 | (64.8, 70.9) | 2454 | 48.2 | 1.4 | (45.5, 50.9) |
| MONTANA | 765 | 61.6 | 2.0 | (57.7, 65.4) | 978 | 43.8 | 1.8 | (40.4, 47.2) |
| NEBRASKA | 1099 | 66.7 | 1.6 | (63.6, 69.7) | 1601 | 49.3 | 1.4 | (46.6, 52.1) |
| NEVADA | 1092 | 61.9 | 2.1 | (57.7, 66.1) | 1039 | 45.2 | 2.4 | (40.6, 49.8) |
| NEW HAMPSHIRE | 496 | 62.9 | 2.6 | (57.8, 67.9) | 696 | 37.5 | 2.1 | (33.4, 41.6) |
| NEW JERSEY | 1103 | 64.1 | 1.8 | (60.6, 67.5) | 1574 | 45.5 | 1.5 | (42.6, 48.5) |
| NEW MEXICO | 1509 | 62.7 | 1.4 | (60.0, 65.4) | 1869 | 48.0 | 1.3 | (45.5, 50.6) |
| NEW YORK | 1049 | 61.3 | 1.7 | (57.9, 64.7) | 1457 | 47.0 | 1.5 | (44.1, 49.9) |
| NORTH CAROLINA | 951 | 68.1 | 1.8 | (64.6, 71.6) | 1380 | 48.1 | 1.6 | (45.0, 51.1) |
| NORTH DAKOTA | 856 | 71.3 | 1.7 | (68.0, 74.7) | 1071 | 48.8 | 1.7 | (45.5, 52.1) |
| OHIO | 629 | 65.3 | 2.4 | (60.5, 70.0) | 919 | 48.9 | 2.0 | (45.0, 52.7) |
| OKLAHOMA | 1119 | 66.7 | 1.6 | (63.6, 69.8) | 1693 | 49.3 | 1.4 | (46.7, 52.0) |
| OREGON | 748 | 63.4 | 1.9 | (59.6, 67.2) | 993 | 45.9 | 1.7 | (42.6, 49.3) |
| PENNSYLVANIA | 1393 | 66.6 | 1.4 | (63.8, 69.4) | 1963 | 46.3 | 1.3 | (43.8, 48.8) |
| RHODE ISLAND | 1530 | 64.0 | 1.4 | (61.2, 66.8) | 2272 | 45.1 | 1.2 | (42.8, 47.5) |
| SOUTH CAROLINA | 1362 | 67.6 | 1.5 | (64.7, 70.5) | 1901 | 49.3 | 1.3 | (46.8, 51.8) |
| SOUTH DAKOTA | 1670 | 67.0 | 1.3 | (64.4, 69.6) | 2331 | 51.8 | 1.1 | (49.6, 54.0) |
| TENNESSEE | 1019 | 65.1 | 1.6 | (61.9, 68.2) | 1847 | 48.6 | 1.3 | (46.1, 51.2) |
| TEXAS | 1951 | 66.6 | 1.4 | (63.8, 69.4) | 2785 | 50.1 | 1.2 | (47.8, 52.5) |
| UTAH | 1364 | 58.7 | 2.0 | (54.8, 62.5) | 1729 | 43.2 | 1.6 | (40.0, 46.4) |
| VERMONT | 1260 | 62.7 | 1.6 | (59.6, 65.7) | 1755 | 42.1 | 1.3 | (39.5, 44.8) |
| VIRGINIA | 1396 | 67.6 | 1.8 | (64.0, 71.1) | 1976 | 49.9 | 1.6 | (46.6, 53.1) |
| WASHINGTON | 1580 | 63.4 | 1.7 | (60.2, 66.7) | 1873 | 45.3 | 1.4 | (42.5, 48.0) |
| WEST VIRGINIA | 1055 | 70.0 | 1.6 | (66.9, 73.1) | 1449 | 54.6 | 1.5 | (51.7, 57.5) |
| WISCONSIN | 915 | 67.3 | 1.9 | (63.6, 71.0) | 1202 | 44.4 | 1.6 | (41.3, 47.5) |
| WYOMING | 1059 | 64.4 | 1.6 | (61.2, 67.6) | 1284 | 46.3 | 1.6 | (43.2, 49.5) |
| PUERTO RICO | 1094 | 64.1 | 1.7 | (60.7, 67.4) | 1800 | 53.3 | 1.4 | (50.6, 56.0) |

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 65.2
 RANGE = 54.8-71.3

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 47.8
 RANGE = 35.1-58.4

* BODY MASS INDEX GE 25.0 AND LE 99.8 FOR MALES AND FEMALES

TABLE 3.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1998 TO 1999
 OVERWEIGHT OR OBESE: BY NEW BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | 1998 PERCENT | 1999 PERCENT | PERCENT DIFFERENCE |
|----------------|-----------------|-----------------|-----------------------|
| ALABAMA | 59.6 | 60.9 | 1.2 |
| ALASKA | 58.3 | 61.1 | 2.8 |
| ARIZONA | 49.5 | 51.6 | 2.1 |
| ARKANSAS | 55.0 | 58.8 | 3.9 |
| CALIFORNIA | 53.6 | 54.8 | 1.3 |
| COLORADO | 49.3 | 47.9 | -1.4 |
| CONNECTICUT | 49.8 | 52.6 | 2.7 |
| DELAWARE | 53.2 | 55.5 | 2.3 |
| DISTRICT OF CO | 50.7 | 50.9 | 0.2 |
| FLORIDA | 53.5 | 56.9 | 3.4 |
| GEORGIA | 54.6 | 57.6 | 3.1 |
| HAWAII | 47.7 | 50.2 | 2.4 |
| IDAHO | 53.1 | 55.2 | 2.1 |
| ILLINOIS | 55.1 | 56.9 | 1.8 |
| INDIANA | 56.2 | 56.7 | 0.5 |
| IOWA | 56.7 | 58.4 | 1.7 |
| KANSAS | 54.8 | 56.1 | 1.4 |
| KENTUCKY | 58.1 | 58.5 | 0.4 |
| LOUISIANA | 56.4 | 58.3 | 1.9 |
| MAINE | 54.4 | 54.8 | 0.4 |
| MARYLAND | 55.5 | 55.4 | -0.1 |
| MASSACHUSETTS | 49.9 | 50.3 | 0.4 |
| MICHIGAN | 57.5 | 59.6 | 2.1 |
| MINNESOTA | 59.5 | 55.9 | -3.7 |
| MISSISSIPPI | 59.3 | 62.3 | 3.0 |
| MISSOURI | 57.1 | 57.7 | 0.6 |
| MONTANA | 51.5 | 52.7 | 1.2 |
| NEBRASKA | 54.1 | 57.8 | 3.7 |
| NEVADA | 50.2 | 53.7 | 3.5 |
| NEW HAMPSHIRE | 51.9 | 50.1 | -1.8 |
| NEW JERSEY | 54.4 | 54.5 | 0.1 |
| NEW MEXICO | 51.7 | 55.3 | 3.6 |
| NEW YORK | 51.3 | 54.0 | 2.6 |
| NORTH CAROLINA | 56.1 | 57.9 | 1.8 |
| NORTH DAKOTA | 57.1 | 60.1 | 3.0 |
| OHIO | 55.1 | 56.9 | 1.9 |
| OKLAHOMA | 53.0 | 57.8 | 4.8 |
| OREGON | 54.6 | 54.5 | -0.1 |
| PENNSYLVANIA | 58.6 | 56.2 | -2.3 |
| RHODE ISLAND | 52.4 | 54.2 | 1.9 |
| SOUTH CAROLINA | 55.0 | 58.3 | 3.3 |
| SOUTH DAKOTA | 51.5 | 59.3 | 7.9 |
| TENNESSEE | 54.4 | 56.6 | 2.2 |
| TEXAS | 57.3 | 58.3 | 1.0 |
| UTAH | 49.3 | 50.9 | 1.7 |
| VERMONT | 47.9 | 52.3 | 4.4 |
| VIRGINIA | 53.2 | 58.7 | 5.5 |
| WASHINGTON | 54.5 | 54.4 | -0.1 |
| WEST VIRGINIA | 57.2 | 62.0 | 4.8 |
| WISCONSIN | 56.5 | 55.7 | -0.8 |
| WYOMING | 51.5 | 55.4 | 3.9 |
| PUERTO RICO | 56.2 | 58.5 | 2.3 |

* BODY MASS INDEX GE 25.0 AND LE 99.8 FOR MALES AND FEMALES
 N/A - NOT AVAILABLE

TABLE 4.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OBESE 1: BY NEW BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL |
|----------------|-------------|---------|----------------|-------------------------|
| ALABAMA | 2045 | 14.3 | 0.9 | (12.5, 16.1) |
| ALASKA | 2007 | 13.9 | 1.1 | (11.7, 16.0) |
| ARIZONA | 1675 | 9.0 | 1.0 | (7.0, 11.0) |
| ARKANSAS | 2908 | 14.8 | 0.8 | (13.4, 16.3) |
| CALIFORNIA | 4152 | 13.1 | 0.6 | (11.9, 14.3) |
| COLORADO | 2986 | 11.0 | 0.8 | (9.3, 12.6) |
| CONNECTICUT | 3320 | 10.8 | 0.7 | (9.4, 12.2) |
| DELAWARE | 2461 | 13.3 | 0.9 | (11.5, 15.1) |
| DISTRICT OF CO | 1242 | 12.0 | 1.0 | (10.0, 14.0) |
| FLORIDA | 5046 | 13.2 | 0.6 | (12.1, 14.4) |
| GEORGIA | 2204 | 13.9 | 0.9 | (12.2, 15.6) |
| HAWAII | 2111 | 9.3 | 0.8 | (7.7, 10.9) |
| IDAHO | 4809 | 15.1 | 0.6 | (13.8, 16.3) |
| ILLINOIS | 2806 | 13.3 | 0.7 | (11.8, 14.7) |
| INDIANA | 2370 | 13.2 | 1.1 | (11.0, 15.3) |
| IOWA | 3528 | 15.0 | 0.7 | (13.6, 16.4) |
| KANSAS | 3680 | 13.4 | 0.6 | (12.1, 14.6) |
| KENTUCKY | 7315 | 14.8 | 0.6 | (13.5, 16.0) |
| LOUISIANA | 1617 | 15.0 | 1.0 | (13.0, 16.9) |
| MAINE | 1613 | 14.0 | 1.0 | (12.1, 15.9) |
| MARYLAND | 3819 | 12.3 | 0.7 | (11.0, 13.6) |
| MASSACHUSETTS | 4689 | 10.8 | 0.6 | (9.6, 11.9) |
| MICHIGAN | 2445 | 16.2 | 0.8 | (14.6, 17.8) |
| MINNESOTA | 5155 | 10.7 | 0.5 | (9.8, 11.6) |
| MISSISSIPPI | 2078 | 14.7 | 0.9 | (12.9, 16.4) |
| MISSOURI | 4161 | 15.2 | 0.8 | (13.7, 16.7) |
| MONTANA | 1743 | 11.7 | 0.9 | (10.0, 13.4) |
| NEBRASKA | 2700 | 15.6 | 0.8 | (14.0, 17.1) |
| NEVADA | 2131 | 11.0 | 1.0 | (9.1, 12.9) |
| NEW HAMPSHIRE | 1192 | 10.2 | 1.0 | (8.2, 12.2) |
| NEW JERSEY | 2677 | 12.3 | 0.8 | (10.7, 13.9) |
| NEW MEXICO | 3378 | 13.1 | 0.7 | (11.8, 14.4) |
| NEW YORK | 2506 | 12.4 | 0.8 | (10.9, 13.8) |
| NORTH CAROLINA | 2331 | 14.7 | 0.8 | (13.0, 16.3) |
| NORTH DAKOTA | 1927 | 16.0 | 0.9 | (14.2, 17.9) |
| OHIO | 1548 | 12.7 | 1.0 | (10.8, 14.6) |
| OKLAHOMA | 2812 | 14.2 | 0.8 | (12.7, 15.7) |
| OREGON | 1741 | 13.8 | 0.9 | (12.0, 15.6) |
| PENNSYLVANIA | 3356 | 14.5 | 0.7 | (13.1, 15.8) |
| RHODE ISLAND | 3802 | 12.4 | 0.6 | (11.2, 13.6) |
| SOUTH CAROLINA | 3263 | 14.6 | 0.7 | (13.2, 16.0) |
| SOUTH DAKOTA | 4001 | 14.1 | 0.6 | (12.9, 15.3) |
| TENNESSEE | 2866 | 14.4 | 0.8 | (13.0, 15.9) |
| TEXAS | 4736 | 14.8 | 0.7 | (13.5, 16.1) |
| UTAH | 3093 | 12.4 | 0.8 | (10.7, 14.0) |
| VERMONT | 3015 | 12.5 | 0.7 | (11.1, 13.9) |
| VIRGINIA | 3372 | 14.1 | 0.9 | (12.4, 15.8) |
| WASHINGTON | 3453 | 12.9 | 0.7 | (11.6, 14.1) |
| WEST VIRGINIA | 2504 | 17.5 | 0.9 | (15.8, 19.1) |
| WISCONSIN | 2117 | 14.6 | 0.8 | (13.0, 16.2) |
| WYOMING | 2343 | 11.7 | 0.7 | (10.3, 13.1) |
| PUERTO RICO | 2894 | 15.6 | 0.8 | (14.0, 17.1) |

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 13.6 RANGE = 9.0-17.5 YEAR 2000 OBJ. 2.3
 * BODY MASS INDEX GE 30.0 AND LE 34.9 FOR MALES AND FEMALES
 June 22, 2000

TABLE 4.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OBESE 1: BY NEW BODY MASS INDEX* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | -----MALES-----* | | | | | *-----FEMALES-----* | | | | |
|----------------|------------------|---------|----------------|-------------------------|---|---------------------|---------|----------------|-------------------------|--|
| | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL | | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL | |
| ALABAMA | 845 | 15.2 | 1.5 | (12.3, 18.1) | * | 1200 | 13.5 | 1.1 | (11.4, 15.7) | |
| ALASKA | 997 | 15.1 | 1.7 | (11.9, 18.4) | * | 1010 | 12.4 | 1.4 | (9.7, 15.0) | |
| ARIZONA | 698 | 11.2 | 1.8 | (7.6, 14.7) | * | 977 | 6.9 | 1.0 | (4.9, 8.8) | |
| ARKANSAS | 1160 | 16.1 | 1.2 | (13.8, 18.4) | * | 1748 | 13.6 | 0.9 | (11.8, 15.4) | |
| CALIFORNIA | 1796 | 13.1 | 0.9 | (11.2, 14.9) | * | 2356 | 13.1 | 0.8 | (11.5, 14.7) | |
| COLORADO | 1359 | 12.0 | 1.3 | (9.5, 14.6) | * | 1627 | 9.9 | 1.0 | (7.9, 11.9) | |
| CONNECTICUT | 1346 | 11.2 | 1.1 | (9.1, 13.4) | * | 1974 | 10.4 | 1.0 | (8.5, 12.3) | |
| DELAWARE | 1018 | 13.5 | 1.4 | (10.7, 16.2) | * | 1443 | 13.2 | 1.2 | (10.8, 15.5) | |
| DISTRICT OF CO | 456 | 11.0 | 1.6 | (7.8, 14.1) | * | 786 | 12.9 | 1.3 | (10.3, 15.5) | |
| FLORIDA | 2029 | 13.9 | 0.9 | (12.1, 15.7) | * | 3017 | 12.6 | 0.7 | (11.3, 14.0) | |
| GEORGIA | 874 | 16.8 | 1.4 | (14.0, 19.6) | * | 1330 | 11.2 | 1.0 | (9.2, 13.2) | |
| HAWAII | 983 | 9.3 | 1.2 | (7.0, 11.5) | * | 1128 | 9.4 | 1.1 | (7.2, 11.5) | |
| IDAHO | 2079 | 17.2 | 1.1 | (15.1, 19.3) | * | 2730 | 12.9 | 0.7 | (11.5, 14.3) | |
| ILLINOIS | 1088 | 14.8 | 1.2 | (12.4, 17.1) | * | 1718 | 11.9 | 0.9 | (10.1, 13.6) | |
| INDIANA | 1010 | 13.6 | 1.7 | (10.4, 16.9) | * | 1360 | 12.7 | 1.5 | (9.7, 15.7) | |
| IOWA | 1409 | 17.3 | 1.2 | (14.9, 19.7) | * | 2119 | 12.8 | 0.8 | (11.3, 14.4) | |
| KANSAS | 1546 | 15.1 | 1.0 | (13.1, 17.1) | * | 2134 | 11.6 | 0.8 | (10.2, 13.1) | |
| KENTUCKY | 2737 | 16.4 | 1.1 | (14.4, 18.5) | * | 4578 | 13.2 | 0.7 | (11.8, 14.5) | |
| LOUISIANA | 658 | 16.0 | 1.6 | (13.0, 19.1) | * | 959 | 14.0 | 1.2 | (11.6, 16.4) | |
| MAINE | 731 | 15.7 | 1.5 | (12.8, 18.7) | * | 882 | 12.4 | 1.2 | (10.0, 14.8) | |
| MARYLAND | 1570 | 12.0 | 1.0 | (10.0, 14.0) | * | 2249 | 12.6 | 0.9 | (10.9, 14.3) | |
| MASSACHUSETTS | 1948 | 13.2 | 1.0 | (11.3, 15.1) | * | 2741 | 8.4 | 0.7 | (7.1, 9.7) | |
| MICHIGAN | 1030 | 17.7 | 1.3 | (15.1, 20.3) | * | 1415 | 14.8 | 1.0 | (12.7, 16.8) | |
| MINNESOTA | 2416 | 12.8 | 0.7 | (11.3, 14.2) | * | 2739 | 8.7 | 0.6 | (7.6, 9.8) | |
| MISSISSIPPI | 843 | 13.5 | 1.3 | (11.0, 16.0) | * | 1235 | 15.8 | 1.2 | (13.4, 18.1) | |
| MISSOURI | 1707 | 16.4 | 1.2 | (14.1, 18.8) | * | 2454 | 14.0 | 1.0 | (12.2, 15.9) | |
| MONTANA | 765 | 14.2 | 1.4 | (11.5, 16.9) | * | 978 | 9.3 | 1.1 | (7.2, 11.3) | |
| NEBRASKA | 1099 | 18.0 | 1.3 | (15.6, 20.5) | * | 1601 | 13.2 | 0.9 | (11.4, 15.0) | |
| NEVADA | 1092 | 10.1 | 1.3 | (7.6, 12.6) | * | 1039 | 11.9 | 1.5 | (9.0, 14.7) | |
| NEW HAMPSHIRE | 496 | 12.2 | 1.7 | (8.7, 15.6) | * | 696 | 8.2 | 1.1 | (6.1, 10.4) | |
| NEW JERSEY | 1103 | 15.1 | 1.4 | (12.3, 17.9) | * | 1574 | 9.6 | 0.9 | (7.9, 11.3) | |
| NEW MEXICO | 1509 | 12.8 | 1.0 | (11.0, 14.7) | * | 1869 | 13.4 | 0.9 | (11.6, 15.1) | |
| NEW YORK | 1049 | 13.4 | 1.2 | (11.0, 15.7) | * | 1457 | 11.4 | 1.0 | (9.5, 13.3) | |
| NORTH CAROLINA | 951 | 16.0 | 1.3 | (13.3, 18.6) | * | 1380 | 13.4 | 1.0 | (11.5, 15.4) | |
| NORTH DAKOTA | 856 | 18.8 | 1.5 | (15.9, 21.7) | * | 1071 | 13.2 | 1.1 | (11.0, 15.5) | |
| OHIO | 629 | 12.5 | 1.5 | (9.6, 15.4) | * | 919 | 12.8 | 1.3 | (10.4, 15.3) | |
| OKLAHOMA | 1119 | 15.1 | 1.2 | (12.8, 17.4) | * | 1693 | 13.3 | 0.9 | (11.5, 15.2) | |
| OREGON | 748 | 14.9 | 1.4 | (12.1, 17.6) | * | 993 | 12.8 | 1.1 | (10.6, 15.0) | |
| PENNSYLVANIA | 1393 | 15.6 | 1.1 | (13.5, 17.7) | * | 1963 | 13.4 | 0.9 | (11.7, 15.1) | |
| RHODE ISLAND | 1530 | 13.9 | 1.0 | (11.9, 15.9) | * | 2272 | 11.0 | 0.8 | (9.5, 12.4) | |
| SOUTH CAROLINA | 1362 | 16.9 | 1.2 | (14.6, 19.1) | * | 1901 | 12.4 | 0.8 | (10.8, 14.1) | |
| SOUTH DAKOTA | 1670 | 15.2 | 0.9 | (13.4, 17.0) | * | 2331 | 13.1 | 0.8 | (11.5, 14.6) | |
| TENNESSEE | 1019 | 15.2 | 1.2 | (12.7, 17.6) | * | 1847 | 13.8 | 0.9 | (12.0, 15.5) | |
| TEXAS | 1951 | 16.3 | 1.0 | (14.3, 18.2) | * | 2785 | 13.4 | 0.9 | (11.7, 15.0) | |
| UTAH | 1364 | 14.0 | 1.3 | (11.4, 16.6) | * | 1729 | 10.7 | 1.0 | (8.8, 12.7) | |
| VERMONT | 1260 | 14.9 | 1.2 | (12.6, 17.2) | * | 1755 | 10.0 | 0.8 | (8.5, 11.6) | |
| VIRGINIA | 1396 | 14.9 | 1.3 | (12.3, 17.5) | * | 1976 | 13.4 | 1.2 | (11.1, 15.7) | |
| WASHINGTON | 1580 | 14.5 | 1.0 | (12.5, 16.5) | * | 1873 | 11.2 | 0.8 | (9.7, 12.8) | |
| WEST VIRGINIA | 1055 | 18.8 | 1.3 | (16.2, 21.4) | * | 1449 | 16.2 | 1.1 | (14.0, 18.4) | |
| WISCONSIN | 915 | 17.2 | 1.3 | (14.6, 19.8) | * | 1202 | 12.1 | 1.0 | (10.2, 14.0) | |
| WYOMING | 1059 | 12.5 | 1.1 | (10.4, 14.7) | * | 1284 | 10.9 | 0.9 | (9.1, 12.7) | |
| PUERTO RICO | 1094 | 16.3 | 1.3 | (13.8, 18.8) | * | 1800 | 14.9 | 0.9 | (13.0, 16.7) | |

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 14.9
 RANGE = 9.3-18.8

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 12.7
 RANGE = 6.9-16.2

* BODY MASS INDEX GE 30.0 AND LE 34.9 FOR MALES AND FEMALES

TABLE 4.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
PREVALENCE DIFFERENCES FROM 1998 TO 1999

OBESE 1: BY NEW BODY MASS INDEX*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | 1998 PERCENT | 1999 PERCENT | PERCENT DIFFERENCE |
|----------------|-----------------|-----------------|-----------------------|
| ALABAMA | 14.2 | 14.3 | 0.2 |
| ALASKA | 15.4 | 13.9 | -1.5 |
| ARIZONA | 9.8 | 9.0 | -0.8 |
| ARKANSAS | 14.2 | 14.8 | 0.6 |
| CALIFORNIA | 11.9 | 13.1 | 1.2 |
| COLORADO | 10.8 | 11.0 | 0.2 |
| CONNECTICUT | 11.4 | 10.8 | -0.6 |
| DELAWARE | 10.6 | 13.3 | 2.7 |
| DISTRICT OF CO | 13.2 | 12.0 | -1.2 |
| FLORIDA | 12.1 | 13.2 | 1.1 |
| GEORGIA | 13.3 | 13.9 | 0.6 |
| HAWAII | 11.2 | 9.3 | -1.9 |
| IDAHO | 11.2 | 15.1 | 3.8 |
| ILLINOIS | 13.4 | 13.3 | -0.1 |
| INDIANA | 12.8 | 13.2 | 0.3 |
| IOWA | 13.2 | 15.0 | 1.8 |
| KANSAS | 12.4 | 13.4 | 0.9 |
| KENTUCKY | 14.5 | 14.8 | 0.3 |
| LOUISIANA | 14.5 | 15.0 | 0.5 |
| MAINE | 12.2 | 14.0 | 1.8 |
| MARYLAND | 15.0 | 12.3 | -2.7 |
| MASSACHUSETTS | 10.3 | 10.8 | 0.5 |
| MICHIGAN | 14.3 | 16.2 | 1.9 |
| MINNESOTA | 11.9 | 10.7 | -1.2 |
| MISSISSIPPI | 14.6 | 14.7 | 0.1 |
| MISSOURI | 14.5 | 15.2 | 0.7 |
| MONTANA | 10.6 | 11.7 | 1.1 |
| NEBRASKA | 12.6 | 15.6 | 2.9 |
| NEVADA | 9.3 | 11.0 | 1.7 |
| NEW HAMPSHIRE | 11.8 | 10.2 | -1.6 |
| NEW JERSEY | 11.4 | 12.3 | 0.9 |
| NEW MEXICO | 10.5 | 13.1 | 2.6 |
| NEW YORK | 11.2 | 12.4 | 1.2 |
| NORTH CAROLINA | 13.0 | 14.7 | 1.6 |
| NORTH DAKOTA | 15.0 | 16.0 | 1.0 |
| OHIO | 13.7 | 12.7 | -1.0 |
| OKLAHOMA | 12.9 | 14.2 | 1.3 |
| OREGON | 12.6 | 13.8 | 1.2 |
| PENNSYLVANIA | 13.0 | 14.5 | 1.5 |
| RHODE ISLAND | 12.1 | 12.4 | 0.3 |
| SOUTH CAROLINA | 13.8 | 14.6 | 0.8 |
| SOUTH DAKOTA | 11.9 | 14.1 | 2.3 |
| TENNESSEE | 13.3 | 14.4 | 1.2 |
| TEXAS | 13.9 | 14.8 | 0.9 |
| UTAH | 11.4 | 12.4 | 1.0 |
| VERMONT | 10.5 | 12.5 | 2.0 |
| VIRGINIA | 12.8 | 14.1 | 1.3 |
| WASHINGTON | 11.8 | 12.9 | 1.1 |
| WEST VIRGINIA | 15.4 | 17.5 | 2.0 |
| WISCONSIN | 12.6 | 14.6 | 2.0 |
| WYOMING | 10.3 | 11.7 | 1.4 |
| PUERTO RICO | 14.0 | 15.6 | 1.6 |

* BODY MASS INDEX GE 30.0 AND LE 34.9 FOR MALES AND FEMALES
N/A - NOT AVAILABLE

TABLE 5.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OBESE 2: BY NEW BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL |
|----------------|-------------|---------|----------------|-------------------------|
| ALABAMA | 2045 | 4.9 | 0.6 | (3.7, 6.0) |
| ALASKA | 2007 | 4.5 | 0.6 | (3.3, 5.7) |
| ARIZONA | 1675 | 2.4 | 0.5 | (1.5, 3.4) |
| ARKANSAS | 2908 | 5.4 | 0.5 | (4.5, 6.4) |
| CALIFORNIA | 4152 | 3.6 | 0.3 | (2.9, 4.2) |
| COLORADO | 2986 | 3.0 | 0.5 | (2.1, 3.9) |
| CONNECTICUT | 3320 | 2.8 | 0.4 | (2.1, 3.6) |
| DELAWARE | 2461 | 2.7 | 0.4 | (1.9, 3.4) |
| DISTRICT OF CO | 1242 | 3.8 | 0.6 | (2.7, 5.0) |
| FLORIDA | 5046 | 3.8 | 0.3 | (3.2, 4.5) |
| GEORGIA | 2204 | 5.0 | 0.5 | (4.0, 6.1) |
| HAWAII | 2111 | 4.2 | 0.6 | (3.1, 5.4) |
| IDAHO | 4809 | 3.6 | 0.3 | (3.0, 4.3) |
| ILLINOIS | 2806 | 4.8 | 0.5 | (3.9, 5.7) |
| INDIANA | 2370 | 5.1 | 0.7 | (3.7, 6.6) |
| IOWA | 3528 | 4.3 | 0.4 | (3.5, 5.0) |
| KANSAS | 3680 | 3.9 | 0.4 | (3.2, 4.6) |
| KENTUCKY | 7315 | 4.6 | 0.4 | (3.8, 5.3) |
| LOUISIANA | 1617 | 4.7 | 0.6 | (3.5, 5.9) |
| MAINE | 1613 | 3.4 | 0.5 | (2.4, 4.3) |
| MARYLAND | 3819 | 3.9 | 0.4 | (3.2, 4.7) |
| MASSACHUSETTS | 4689 | 2.7 | 0.3 | (2.1, 3.3) |
| MICHIGAN | 2445 | 4.7 | 0.5 | (3.7, 5.6) |
| MINNESOTA | 5155 | 3.7 | 0.3 | (3.1, 4.2) |
| MISSISSIPPI | 2078 | 5.4 | 0.6 | (4.2, 6.6) |
| MISSOURI | 4161 | 4.1 | 0.4 | (3.3, 4.9) |
| MONTANA | 1743 | 2.9 | 0.4 | (2.1, 3.8) |
| NEBRASKA | 2700 | 3.9 | 0.4 | (3.1, 4.7) |
| NEVADA | 2131 | 3.1 | 0.6 | (2.0, 4.2) |
| NEW HAMPSHIRE | 1192 | 3.0 | 0.6 | (1.9, 4.2) |
| NEW JERSEY | 2677 | 3.4 | 0.4 | (2.6, 4.2) |
| NEW MEXICO | 3378 | 3.4 | 0.4 | (2.8, 4.1) |
| NEW YORK | 2506 | 3.4 | 0.4 | (2.5, 4.2) |
| NORTH CAROLINA | 2331 | 4.6 | 0.5 | (3.6, 5.5) |
| NORTH DAKOTA | 1927 | 3.5 | 0.5 | (2.6, 4.3) |
| OHIO | 1548 | 5.1 | 0.7 | (3.8, 6.4) |
| OKLAHOMA | 2812 | 4.1 | 0.4 | (3.3, 4.9) |
| OREGON | 1741 | 4.0 | 0.5 | (3.0, 5.0) |
| PENNSYLVANIA | 3356 | 4.0 | 0.4 | (3.3, 4.7) |
| RHODE ISLAND | 3802 | 3.1 | 0.3 | (2.4, 3.7) |
| SOUTH CAROLINA | 3263 | 4.0 | 0.4 | (3.2, 4.7) |
| SOUTH DAKOTA | 4001 | 3.7 | 0.4 | (3.0, 4.4) |
| TENNESSEE | 2866 | 4.3 | 0.4 | (3.4, 5.1) |
| TEXAS | 4736 | 4.8 | 0.4 | (4.0, 5.5) |
| UTAH | 3093 | 2.9 | 0.4 | (2.1, 3.7) |
| VERMONT | 3015 | 3.7 | 0.4 | (2.9, 4.5) |
| VIRGINIA | 3372 | 3.8 | 0.4 | (3.0, 4.5) |
| WASHINGTON | 3453 | 3.9 | 0.4 | (3.2, 4.7) |
| WEST VIRGINIA | 2504 | 4.4 | 0.4 | (3.6, 5.3) |
| WISCONSIN | 2117 | 3.8 | 0.5 | (2.8, 4.8) |
| WYOMING | 2343 | 3.2 | 0.4 | (2.5, 4.0) |
| PUERTO RICO | 2894 | 4.1 | 0.4 | (3.2, 4.9) |

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 3.9 RANGE = 2.4- 5.4 YEAR 2000 OBJ. 2.3
 * BODY MASS INDEX GE 35.0 AND LE 39.9 FOR MALES AND FEMALES
 June 22, 2000

TABLE 5.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OBESE 2: BY NEW BODY MASS INDEX* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | -----MALES----- | | | | | -----FEMALES----- | | | | |
|----------------|-----------------|---------|----------------|-------------------------|---|-------------------|---------|----------------|-------------------------|---|
| | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL | | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL | |
| ALABAMA | 845 | 3.7 | 0.7 | (2.3, 5.1) | * | 1200 | 6.0 | 1.0 | (4.1, 7.8) | * |
| ALASKA | 997 | 4.0 | 0.8 | (2.5, 5.6) | * | 1010 | 5.1 | 0.9 | (3.3, 6.9) | * |
| ARIZONA | 698 | 2.5 | 0.8 | (1.0, 4.0) | * | 977 | 2.4 | 0.6 | (1.2, 3.5) | * |
| ARKANSAS | 1160 | 5.0 | 0.7 | (3.6, 6.4) | * | 1748 | 5.9 | 0.6 | (4.7, 7.1) | * |
| CALIFORNIA | 1796 | 3.1 | 0.5 | (2.2, 4.0) | * | 2356 | 4.1 | 0.5 | (3.2, 5.0) | * |
| COLORADO | 1359 | 2.7 | 0.6 | (1.6, 3.8) | * | 1627 | 3.4 | 0.7 | (2.0, 4.7) | * |
| CONNECTICUT | 1346 | 2.9 | 0.6 | (1.8, 4.1) | * | 1974 | 2.8 | 0.5 | (1.8, 3.7) | * |
| DELAWARE | 1018 | 2.7 | 0.6 | (1.5, 3.9) | * | 1443 | 2.7 | 0.5 | (1.7, 3.6) | * |
| DISTRICT OF CO | 456 | 3.5 | 0.9 | (1.6, 5.3) | * | 786 | 4.2 | 0.8 | (2.6, 5.7) | * |
| FLORIDA | 2029 | 3.3 | 0.5 | (2.4, 4.2) | * | 3017 | 4.3 | 0.5 | (3.4, 5.2) | * |
| GEORGIA | 874 | 4.7 | 0.9 | (3.1, 6.4) | * | 1330 | 5.3 | 0.7 | (4.0, 6.6) | * |
| HAWAII | 983 | 4.4 | 0.9 | (2.7, 6.2) | * | 1128 | 4.0 | 0.7 | (2.6, 5.4) | * |
| IDAHO | 2079 | 3.6 | 0.5 | (2.7, 4.6) | * | 2730 | 3.7 | 0.5 | (2.7, 4.6) | * |
| ILLINOIS | 1088 | 3.9 | 0.7 | (2.6, 5.2) | * | 1718 | 5.7 | 0.6 | (4.5, 6.9) | * |
| INDIANA | 1010 | 4.3 | 1.0 | (2.3, 6.2) | * | 1360 | 6.0 | 1.1 | (3.8, 8.1) | * |
| IOWA | 1409 | 4.0 | 0.6 | (2.9, 5.2) | * | 2119 | 4.5 | 0.6 | (3.4, 5.6) | * |
| KANSAS | 1546 | 3.2 | 0.5 | (2.3, 4.2) | * | 2134 | 4.5 | 0.5 | (3.5, 5.5) | * |
| KENTUCKY | 2737 | 4.4 | 0.6 | (3.3, 5.5) | * | 4578 | 4.7 | 0.4 | (3.9, 5.5) | * |
| LOUISIANA | 658 | 4.8 | 1.0 | (2.9, 6.8) | * | 959 | 4.5 | 0.7 | (3.1, 5.9) | * |
| MAINE | 731 | 2.2 | 0.6 | (1.1, 3.3) | * | 882 | 4.5 | 0.7 | (3.0, 5.9) | * |
| MARYLAND | 1570 | 3.6 | 0.5 | (2.5, 4.6) | * | 2249 | 4.3 | 0.5 | (3.3, 5.3) | * |
| MASSACHUSETTS | 1948 | 3.1 | 0.5 | (2.1, 4.2) | * | 2741 | 2.3 | 0.3 | (1.7, 2.9) | * |
| MICHIGAN | 1030 | 4.5 | 0.7 | (3.1, 5.9) | * | 1415 | 4.9 | 0.7 | (3.6, 6.1) | * |
| MINNESOTA | 2416 | 3.7 | 0.4 | (2.9, 4.6) | * | 2739 | 3.6 | 0.4 | (2.9, 4.3) | * |
| MISSISSIPPI | 843 | 5.1 | 1.0 | (3.1, 7.1) | * | 1235 | 5.7 | 0.7 | (4.3, 7.1) | * |
| MISSOURI | 1707 | 4.1 | 0.6 | (2.9, 5.3) | * | 2454 | 4.2 | 0.5 | (3.1, 5.2) | * |
| MONTANA | 765 | 2.1 | 0.6 | (1.0, 3.2) | * | 978 | 3.7 | 0.7 | (2.4, 5.0) | * |
| NEBRASKA | 1099 | 4.1 | 0.7 | (2.8, 5.3) | * | 1601 | 3.7 | 0.5 | (2.7, 4.7) | * |
| NEVADA | 1092 | 2.3 | 0.6 | (1.2, 3.4) | * | 1039 | 4.0 | 1.0 | (2.0, 6.0) | * |
| NEW HAMPSHIRE | 496 | 3.0 | 0.9 | (1.3, 4.7) | * | 696 | 3.1 | 0.8 | (1.6, 4.6) | * |
| NEW JERSEY | 1103 | 3.2 | 0.6 | (2.1, 4.3) | * | 1574 | 3.5 | 0.6 | (2.4, 4.7) | * |
| NEW MEXICO | 1509 | 3.4 | 0.5 | (2.4, 4.5) | * | 1869 | 3.5 | 0.5 | (2.6, 4.3) | * |
| NEW YORK | 1049 | 3.1 | 0.6 | (1.9, 4.2) | * | 1457 | 3.6 | 0.6 | (2.5, 4.8) | * |
| NORTH CAROLINA | 951 | 3.2 | 0.7 | (1.9, 4.4) | * | 1380 | 6.0 | 0.7 | (4.6, 7.3) | * |
| NORTH DAKOTA | 856 | 3.1 | 0.6 | (1.9, 4.3) | * | 1071 | 3.8 | 0.7 | (2.5, 5.1) | * |
| OHIO | 629 | 5.1 | 1.0 | (3.2, 7.0) | * | 919 | 5.1 | 0.9 | (3.4, 6.8) | * |
| OKLAHOMA | 1119 | 4.4 | 0.7 | (3.1, 5.7) | * | 1693 | 3.9 | 0.5 | (3.0, 4.8) | * |
| OREGON | 748 | 3.8 | 0.7 | (2.4, 5.3) | * | 993 | 4.2 | 0.7 | (2.9, 5.5) | * |
| PENNSYLVANIA | 1393 | 3.9 | 0.6 | (2.7, 5.0) | * | 1963 | 4.1 | 0.5 | (3.2, 5.1) | * |
| RHODE ISLAND | 1530 | 3.0 | 0.5 | (2.0, 4.1) | * | 2272 | 3.1 | 0.4 | (2.3, 3.9) | * |
| SOUTH CAROLINA | 1362 | 3.6 | 0.6 | (2.5, 4.7) | * | 1901 | 4.4 | 0.5 | (3.4, 5.3) | * |
| SOUTH DAKOTA | 1670 | 3.9 | 0.6 | (2.8, 5.0) | * | 2331 | 3.6 | 0.4 | (2.8, 4.3) | * |
| TENNESSEE | 1019 | 4.1 | 0.6 | (2.8, 5.3) | * | 1847 | 4.4 | 0.5 | (3.4, 5.5) | * |
| TEXAS | 1951 | 4.3 | 0.5 | (3.3, 5.3) | * | 2785 | 5.2 | 0.6 | (4.0, 6.4) | * |
| UTAH | 1364 | 2.7 | 0.6 | (1.6, 3.8) | * | 1729 | 3.1 | 0.6 | (2.0, 4.3) | * |
| VERMONT | 1260 | 3.6 | 0.6 | (2.4, 4.8) | * | 1755 | 3.8 | 0.5 | (2.8, 4.9) | * |
| VIRGINIA | 1396 | 3.1 | 0.6 | (2.0, 4.2) | * | 1976 | 4.4 | 0.6 | (3.3, 5.5) | * |
| WASHINGTON | 1580 | 3.5 | 0.5 | (2.6, 4.5) | * | 1873 | 4.4 | 0.6 | (3.2, 5.5) | * |
| WEST VIRGINIA | 1055 | 3.6 | 0.6 | (2.4, 4.8) | * | 1449 | 5.2 | 0.6 | (4.0, 6.4) | * |
| WISCONSIN | 915 | 4.6 | 0.9 | (2.9, 6.4) | * | 1202 | 2.9 | 0.5 | (1.9, 3.9) | * |
| WYOMING | 1059 | 2.0 | 0.5 | (1.1, 2.9) | * | 1284 | 4.5 | 0.6 | (3.3, 5.6) | * |
| PUERTO RICO | 1094 | 4.0 | 0.7 | (2.7, 5.3) | * | 1800 | 4.2 | 0.5 | (3.1, 5.2) | * |

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 3.6
 RANGE = 2.0- 5.1

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 4.2
 RANGE = 2.3- 6.0

* BODY MASS INDEX GE 35.0 AND LE 39.9 FOR MALES AND FEMALES

TABLE 5.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1998 TO 1999
 OBESE 2: BY NEW BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | 1998 PERCENT | 1999 PERCENT | PERCENT DIFFERENCE |
|----------------|-----------------|-----------------|-----------------------|
| ALABAMA | 5.1 | 4.9 | -0.3 |
| ALASKA | 3.5 | 4.5 | 1.1 |
| ARIZONA | 2.2 | 2.4 | 0.2 |
| ARKANSAS | 4.2 | 5.4 | 1.3 |
| CALIFORNIA | 3.7 | 3.6 | -0.1 |
| COLORADO | 2.1 | 3.0 | 0.9 |
| CONNECTICUT | 3.0 | 2.8 | -0.2 |
| DELAWARE | 4.1 | 2.7 | -1.4 |
| DISTRICT OF CO | 4.6 | 3.8 | -0.8 |
| FLORIDA | 4.3 | 3.8 | -0.5 |
| GEORGIA | 3.6 | 5.0 | 1.4 |
| HAWAII | 2.8 | 4.2 | 1.4 |
| IDAHO | 3.3 | 3.6 | 0.3 |
| ILLINOIS | 3.6 | 4.8 | 1.2 |
| INDIANA | 4.3 | 5.1 | 0.8 |
| IOWA | 4.8 | 4.3 | -0.5 |
| KANSAS | 3.6 | 3.9 | 0.3 |
| KENTUCKY | 3.7 | 4.6 | 0.8 |
| LOUISIANA | 5.1 | 4.7 | -0.5 |
| MAINE | 3.7 | 3.4 | -0.4 |
| MARYLAND | 3.6 | 3.9 | 0.3 |
| MASSACHUSETTS | 2.7 | 2.7 | 0.0 |
| MICHIGAN | 4.1 | 4.7 | 0.6 |
| MINNESOTA | 2.8 | 3.7 | 0.9 |
| MISSISSIPPI | 5.3 | 5.4 | 0.1 |
| MISSOURI | 3.7 | 4.1 | 0.5 |
| MONTANA | 2.6 | 2.9 | 0.3 |
| NEBRASKA | 3.7 | 3.9 | 0.2 |
| NEVADA | 3.2 | 3.1 | -0.1 |
| NEW HAMPSHIRE | 2.8 | 3.0 | 0.2 |
| NEW JERSEY | 2.9 | 3.4 | 0.5 |
| NEW MEXICO | 3.2 | 3.4 | 0.3 |
| NEW YORK | 3.3 | 3.4 | 0.0 |
| NORTH CAROLINA | 3.8 | 4.6 | 0.8 |
| NORTH DAKOTA | 2.7 | 3.5 | 0.8 |
| OHIO | 4.4 | 5.1 | 0.7 |
| OKLAHOMA | 4.2 | 4.1 | -0.0 |
| OREGON | 4.1 | 4.0 | -0.1 |
| PENNSYLVANIA | 3.9 | 4.0 | 0.1 |
| RHODE ISLAND | 3.7 | 3.1 | -0.7 |
| SOUTH CAROLINA | 4.8 | 4.0 | -0.8 |
| SOUTH DAKOTA | 2.9 | 3.7 | 0.9 |
| TENNESSEE | 4.1 | 4.3 | 0.1 |
| TEXAS | 4.2 | 4.8 | 0.5 |
| UTAH | 2.8 | 2.9 | 0.1 |
| VERMONT | 3.2 | 3.7 | 0.5 |
| VIRGINIA | 3.9 | 3.8 | -0.2 |
| WASHINGTON | 4.3 | 3.9 | -0.3 |
| WEST VIRGINIA | 5.3 | 4.4 | -0.8 |
| WISCONSIN | 4.3 | 3.8 | -0.5 |
| WYOMING | 3.1 | 3.2 | 0.2 |
| PUERTO RICO | 3.9 | 4.1 | 0.2 |

* BODY MASS INDEX GE 35.0 AND LE 39.9 FOR MALES AND FEMALES
 N/A - NOT AVAILABLE

TABLE 6.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OBESE 3: BY NEW BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL |
|----------------|-------------|---------|----------------|-------------------------|
| ALABAMA | 2045 | 3.3 | 0.4 | (2.4, 4.1) |
| ALASKA | 2007 | 2.0 | 0.4 | (1.3, 2.7) |
| ARIZONA | 1675 | 0.9 | 0.3 | (0.4, 1.3) |
| ARKANSAS | 2908 | 2.4 | 0.3 | (1.7, 3.0) |
| CALIFORNIA | 4152 | 2.0 | 0.3 | (1.5, 2.5) |
| COLORADO | 2986 | 0.9 | 0.2 | (0.5, 1.3) |
| CONNECTICUT | 3320 | 1.5 | 0.3 | (1.0, 2.1) |
| DELAWARE | 2461 | 1.6 | 0.3 | (1.0, 2.2) |
| DISTRICT OF CO | 1242 | 2.7 | 0.5 | (1.8, 3.6) |
| FLORIDA | 5046 | 1.5 | 0.2 | (1.1, 1.9) |
| GEORGIA | 2204 | 2.2 | 0.4 | (1.5, 2.9) |
| HAWAII | 2111 | 2.2 | 0.5 | (1.1, 3.2) |
| IDAHO | 4809 | 1.3 | 0.2 | (1.0, 1.6) |
| ILLINOIS | 2806 | 2.8 | 0.5 | (1.8, 3.9) |
| INDIANA | 2370 | 1.6 | 0.4 | (0.9, 2.3) |
| IOWA | 3528 | 2.2 | 0.3 | (1.7, 2.7) |
| KANSAS | 3680 | 1.6 | 0.2 | (1.2, 2.1) |
| KENTUCKY | 7315 | 2.4 | 0.3 | (1.9, 2.9) |
| LOUISIANA | 1617 | 2.7 | 0.4 | (1.9, 3.5) |
| MAINE | 1613 | 2.1 | 0.4 | (1.2, 2.9) |
| MARYLAND | 3819 | 2.0 | 0.3 | (1.5, 2.5) |
| MASSACHUSETTS | 4689 | 1.3 | 0.2 | (0.9, 1.7) |
| MICHIGAN | 2445 | 1.9 | 0.3 | (1.4, 2.5) |
| MINNESOTA | 5155 | 1.1 | 0.1 | (0.8, 1.3) |
| MISSISSIPPI | 2078 | 3.1 | 0.6 | (2.0, 4.2) |
| MISSOURI | 4161 | 2.3 | 0.3 | (1.7, 2.9) |
| MONTANA | 1743 | 1.2 | 0.3 | (0.7, 1.7) |
| NEBRASKA | 2700 | 1.6 | 0.3 | (1.1, 2.0) |
| NEVADA | 2131 | 1.7 | 0.4 | (0.9, 2.6) |
| NEW HAMPSHIRE | 1192 | 1.4 | 0.4 | (0.7, 2.1) |
| NEW JERSEY | 2677 | 1.4 | 0.3 | (0.8, 1.9) |
| NEW MEXICO | 3378 | 1.2 | 0.2 | (0.8, 1.5) |
| NEW YORK | 2506 | 1.7 | 0.3 | (1.2, 2.3) |
| NORTH CAROLINA | 2331 | 2.3 | 0.4 | (1.6, 2.9) |
| NORTH DAKOTA | 1927 | 2.4 | 0.4 | (1.7, 3.1) |
| OHIO | 1548 | 2.5 | 0.4 | (1.6, 3.3) |
| OKLAHOMA | 2812 | 2.7 | 0.3 | (2.0, 3.4) |
| OREGON | 1741 | 2.1 | 0.4 | (1.4, 2.8) |
| PENNSYLVANIA | 3356 | 1.8 | 0.3 | (1.3, 2.3) |
| RHODE ISLAND | 3802 | 1.3 | 0.2 | (0.9, 1.7) |
| SOUTH CAROLINA | 3263 | 2.1 | 0.3 | (1.6, 2.6) |
| SOUTH DAKOTA | 4001 | 1.8 | 0.2 | (1.3, 2.3) |
| TENNESSEE | 2866 | 1.9 | 0.3 | (1.3, 2.4) |
| TEXAS | 4736 | 2.1 | 0.3 | (1.6, 2.6) |
| UTAH | 3093 | 1.5 | 0.3 | (0.9, 2.0) |
| VERMONT | 3015 | 1.8 | 0.3 | (1.3, 2.3) |
| VIRGINIA | 3372 | 1.4 | 0.2 | (0.9, 1.8) |
| WASHINGTON | 3453 | 1.4 | 0.2 | (1.0, 1.9) |
| WEST VIRGINIA | 2504 | 2.7 | 0.3 | (2.1, 3.4) |
| WISCONSIN | 2117 | 1.6 | 0.3 | (1.0, 2.1) |
| WYOMING | 2343 | 1.9 | 0.3 | (1.3, 2.6) |
| PUERTO RICO | 2894 | 1.7 | 0.3 | (1.1, 2.2) |

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 1.8 RANGE = 0.9- 3.3 YEAR 2000 OBJ. 2.3
 * BODY MASS INDEX GE 40.0 AND LE 99.8 FOR MALES AND FEMALES
 June 22, 2000

TABLE 6.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OBESE 3: BY NEW BODY MASS INDEX* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | -----MALES----- | | | | | -----FEMALES----- | | | | |
|----------------|-----------------|---------|----------------|-------------------------|---|-------------------|---------|----------------|-------------------------|---|
| | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL | | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL | |
| ALABAMA | 845 | 2.5 | 0.6 | (1.4, 3.6) | * | 1200 | 4.0 | 0.6 | (2.8, 5.1) | * |
| ALASKA | 997 | 1.4 | 0.5 | (0.4, 2.3) | * | 1010 | 2.8 | 0.6 | (1.7, 3.9) | * |
| ARIZONA | 698 | 0.5 | 0.2 | (0.0, 0.9) | * | 977 | 1.2 | 0.5 | (0.3, 2.1) | * |
| ARKANSAS | 1160 | 1.2 | 0.3 | (0.6, 1.8) | * | 1748 | 3.5 | 0.6 | (2.4, 4.6) | * |
| CALIFORNIA | 1796 | 1.6 | 0.4 | (0.9, 2.3) | * | 2356 | 2.5 | 0.4 | (1.7, 3.3) | * |
| COLORADO | 1359 | 0.6 | 0.3 | (0.0, 1.1) | * | 1627 | 1.3 | 0.3 | (0.6, 1.9) | * |
| CONNECTICUT | 1346 | 1.2 | 0.5 | (0.3, 2.1) | * | 1974 | 1.8 | 0.3 | (1.2, 2.4) | * |
| DELAWARE | 1018 | 1.2 | 0.4 | (0.4, 2.0) | * | 1443 | 1.9 | 0.4 | (1.0, 2.7) | * |
| DISTRICT OF CO | 456 | 1.0 | 0.5 | (0.0, 1.9) | * | 786 | 4.3 | 0.8 | (2.8, 5.7) | * |
| FLORIDA | 2029 | 1.2 | 0.3 | (0.7, 1.6) | * | 3017 | 1.9 | 0.3 | (1.3, 2.5) | * |
| GEORGIA | 874 | 0.9 | 0.5 | (0.0, 1.9) | * | 1330 | 3.3 | 0.6 | (2.3, 4.4) | * |
| HAWAII | 983 | 2.7 | 0.9 | (0.9, 4.5) | * | 1128 | 1.7 | 0.5 | (0.6, 2.7) | * |
| IDAHO | 2079 | 0.9 | 0.2 | (0.4, 1.3) | * | 2730 | 1.7 | 0.3 | (1.2, 2.2) | * |
| ILLINOIS | 1088 | 2.9 | 1.0 | (0.9, 4.9) | * | 1718 | 2.7 | 0.4 | (1.9, 3.6) | * |
| INDIANA | 1010 | 1.2 | 0.4 | (0.3, 2.0) | * | 1360 | 2.0 | 0.6 | (0.9, 3.1) | * |
| IOWA | 1409 | 1.7 | 0.4 | (1.0, 2.4) | * | 2119 | 2.7 | 0.4 | (1.9, 3.4) | * |
| KANSAS | 1546 | 1.2 | 0.3 | (0.6, 1.7) | * | 2134 | 2.1 | 0.3 | (1.4, 2.8) | * |
| KENTUCKY | 2737 | 1.4 | 0.3 | (0.9, 1.9) | * | 4578 | 3.3 | 0.4 | (2.4, 4.1) | * |
| LOUISIANA | 658 | 1.7 | 0.5 | (0.7, 2.7) | * | 959 | 3.6 | 0.7 | (2.3, 4.9) | * |
| MAINE | 731 | 2.2 | 0.7 | (0.9, 3.5) | * | 882 | 1.9 | 0.5 | (0.9, 2.9) | * |
| MARYLAND | 1570 | 0.9 | 0.3 | (0.4, 1.4) | * | 2249 | 3.0 | 0.4 | (2.1, 3.9) | * |
| MASSACHUSETTS | 1948 | 1.2 | 0.3 | (0.6, 1.7) | * | 2741 | 1.4 | 0.3 | (0.8, 1.9) | * |
| MICHIGAN | 1030 | 1.2 | 0.4 | (0.5, 1.9) | * | 1415 | 2.6 | 0.4 | (1.7, 3.5) | * |
| MINNESOTA | 2416 | 0.7 | 0.2 | (0.3, 1.0) | * | 2739 | 1.5 | 0.2 | (1.0, 2.0) | * |
| MISSISSIPPI | 843 | 2.5 | 1.0 | (0.4, 4.5) | * | 1235 | 3.7 | 0.6 | (2.6, 4.8) | * |
| MISSOURI | 1707 | 1.6 | 0.4 | (0.9, 2.3) | * | 2454 | 3.0 | 0.5 | (2.0, 4.0) | * |
| MONTANA | 765 | 1.0 | 0.4 | (0.3, 1.7) | * | 978 | 1.4 | 0.4 | (0.6, 2.1) | * |
| NEBRASKA | 1099 | 1.5 | 0.4 | (0.8, 2.2) | * | 1601 | 1.6 | 0.4 | (0.9, 2.3) | * |
| NEVADA | 1092 | 1.6 | 0.6 | (0.4, 2.9) | * | 1039 | 1.8 | 0.6 | (0.7, 3.0) | * |
| NEW HAMPSHIRE | 496 | 1.4 | 0.5 | (0.4, 2.4) | * | 696 | 1.4 | 0.5 | (0.4, 2.3) | * |
| NEW JERSEY | 1103 | 1.2 | 0.5 | (0.3, 2.1) | * | 1574 | 1.6 | 0.4 | (0.9, 2.3) | * |
| NEW MEXICO | 1509 | 0.6 | 0.2 | (0.2, 0.9) | * | 1869 | 1.8 | 0.3 | (1.2, 2.4) | * |
| NEW YORK | 1049 | 1.5 | 0.4 | (0.7, 2.3) | * | 1457 | 1.9 | 0.4 | (1.2, 2.6) | * |
| NORTH CAROLINA | 951 | 1.9 | 0.5 | (0.9, 2.9) | * | 1380 | 2.7 | 0.5 | (1.7, 3.6) | * |
| NORTH DAKOTA | 856 | 1.6 | 0.5 | (0.7, 2.6) | * | 1071 | 3.2 | 0.6 | (2.0, 4.3) | * |
| OHIO | 629 | 1.3 | 0.5 | (0.4, 2.2) | * | 919 | 3.7 | 0.7 | (2.2, 5.1) | * |
| OKLAHOMA | 1119 | 2.1 | 0.5 | (1.2, 3.1) | * | 1693 | 3.3 | 0.5 | (2.3, 4.2) | * |
| OREGON | 748 | 1.4 | 0.5 | (0.5, 2.3) | * | 993 | 2.8 | 0.6 | (1.7, 3.9) | * |
| PENNSYLVANIA | 1393 | 1.4 | 0.4 | (0.7, 2.1) | * | 1963 | 2.2 | 0.4 | (1.5, 2.9) | * |
| RHODE ISLAND | 1530 | 0.7 | 0.2 | (0.3, 1.1) | * | 2272 | 1.9 | 0.3 | (1.3, 2.6) | * |
| SOUTH CAROLINA | 1362 | 1.1 | 0.3 | (0.6, 1.7) | * | 1901 | 3.0 | 0.4 | (2.1, 3.9) | * |
| SOUTH DAKOTA | 1670 | 1.2 | 0.3 | (0.6, 1.8) | * | 2331 | 2.4 | 0.3 | (1.7, 3.1) | * |
| TENNESSEE | 1019 | 0.7 | 0.3 | (0.2, 1.2) | * | 1847 | 3.0 | 0.5 | (2.1, 3.9) | * |
| TEXAS | 1951 | 1.9 | 0.4 | (1.1, 2.6) | * | 2785 | 2.3 | 0.3 | (1.7, 2.9) | * |
| UTAH | 1364 | 0.7 | 0.2 | (0.2, 1.1) | * | 1729 | 2.3 | 0.5 | (1.2, 3.3) | * |
| VERMONT | 1260 | 1.4 | 0.4 | (0.7, 2.2) | * | 1755 | 2.2 | 0.4 | (1.4, 2.9) | * |
| VIRGINIA | 1396 | 0.8 | 0.2 | (0.4, 1.3) | * | 1976 | 1.9 | 0.3 | (1.2, 2.6) | * |
| WASHINGTON | 1580 | 0.9 | 0.3 | (0.4, 1.5) | * | 1873 | 2.0 | 0.3 | (1.3, 2.6) | * |
| WEST VIRGINIA | 1055 | 1.9 | 0.5 | (1.0, 2.8) | * | 1449 | 3.5 | 0.5 | (2.5, 4.5) | * |
| WISCONSIN | 915 | 0.9 | 0.3 | (0.3, 1.6) | * | 1202 | 2.2 | 0.4 | (1.3, 3.0) | * |
| WYOMING | 1059 | 1.5 | 0.4 | (0.7, 2.4) | * | 1284 | 2.3 | 0.5 | (1.4, 3.3) | * |
| PUERTO RICO | 1094 | 1.4 | 0.5 | (0.5, 2.3) | * | 1800 | 1.9 | 0.4 | (1.2, 2.6) | * |

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 1.3
 RANGE = 0.5- 2.9

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 2.2
 RANGE = 1.2- 4.3

* BODY MASS INDEX GE 40.0 AND LE 99.8 FOR MALES AND FEMALES

TABLE 6.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1998 TO 1999
 OBESE 3: BY NEW BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | 1998 PERCENT | 1999 PERCENT | PERCENT DIFFERENCE |
|----------------|-----------------|-----------------|-----------------------|
| ALABAMA | 2.0 | 3.3 | 1.3 |
| ALASKA | 2.6 | 2.0 | -0.6 |
| ARIZONA | 1.1 | 0.9 | -0.3 |
| ARKANSAS | 1.5 | 2.4 | 0.9 |
| CALIFORNIA | 1.8 | 2.0 | 0.3 |
| COLORADO | 1.5 | 0.9 | -0.6 |
| CONNECTICUT | 1.1 | 1.5 | 0.4 |
| DELAWARE | 2.6 | 1.6 | -1.0 |
| DISTRICT OF CO | 2.4 | 2.7 | 0.4 |
| FLORIDA | 1.5 | 1.5 | 0.0 |
| GEORGIA | 2.3 | 2.2 | -0.1 |
| HAWAII | 1.5 | 2.2 | 0.6 |
| IDAHO | 1.9 | 1.3 | -0.6 |
| ILLINOIS | 1.6 | 2.8 | 1.2 |
| INDIANA | 2.8 | 1.6 | -1.2 |
| IOWA | 1.8 | 2.2 | 0.4 |
| KANSAS | 1.6 | 1.6 | 0.0 |
| KENTUCKY | 2.2 | 2.4 | 0.1 |
| LOUISIANA | 2.2 | 2.7 | 0.5 |
| MAINE | 1.5 | 2.1 | 0.6 |
| MARYLAND | 1.9 | 2.0 | 0.1 |
| MASSACHUSETTS | 1.3 | 1.3 | -0.0 |
| MICHIGAN | 2.8 | 1.9 | -0.9 |
| MINNESOTA | 1.5 | 1.1 | -0.5 |
| MISSISSIPPI | 2.9 | 3.1 | 0.2 |
| MISSOURI | 2.3 | 2.3 | 0.0 |
| MONTANA | 1.9 | 1.2 | -0.7 |
| NEBRASKA | 2.0 | 1.6 | -0.5 |
| NEVADA | 1.5 | 1.7 | 0.3 |
| NEW HAMPSHIRE | 1.0 | 1.4 | 0.3 |
| NEW JERSEY | 1.3 | 1.4 | 0.1 |
| NEW MEXICO | 1.5 | 1.2 | -0.3 |
| NEW YORK | 1.8 | 1.7 | -0.1 |
| NORTH CAROLINA | 2.6 | 2.3 | -0.3 |
| NORTH DAKOTA | 1.5 | 2.4 | 0.9 |
| OHIO | 2.0 | 2.5 | 0.5 |
| OKLAHOMA | 2.4 | 2.7 | 0.3 |
| OREGON | 1.6 | 2.1 | 0.6 |
| PENNSYLVANIA | 2.5 | 1.8 | -0.7 |
| RHODE ISLAND | 1.1 | 1.3 | 0.3 |
| SOUTH CAROLINA | 2.0 | 2.1 | 0.0 |
| SOUTH DAKOTA | 1.1 | 1.8 | 0.7 |
| TENNESSEE | 1.8 | 1.9 | 0.1 |
| TEXAS | 2.1 | 2.1 | -0.0 |
| UTAH | 1.7 | 1.5 | -0.2 |
| VERMONT | 1.1 | 1.8 | 0.7 |
| VIRGINIA | 2.0 | 1.4 | -0.6 |
| WASHINGTON | 2.1 | 1.4 | -0.6 |
| WEST VIRGINIA | 3.3 | 2.7 | -0.5 |
| WISCONSIN | 1.5 | 1.6 | 0.1 |
| WYOMING | 1.7 | 1.9 | 0.2 |
| PUERTO RICO | 1.5 | 1.7 | 0.2 |

* BODY MASS INDEX GE 40.0 AND LE 99.8 FOR MALES AND FEMALES
 N/A - NOT AVAILABLE

TABLE 7.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OBESE : BY NEW BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL |
|----------------|-------------|---------|----------------|-------------------------|
| ALABAMA | 2045 | 22.4 | 1.1 | (20.3, 24.6) |
| ALASKA | 2007 | 20.4 | 1.2 | (18.0, 22.8) |
| ARIZONA | 1675 | 12.3 | 1.1 | (10.1, 14.5) |
| ARKANSAS | 2908 | 22.7 | 0.9 | (20.9, 24.4) |
| CALIFORNIA | 4152 | 18.7 | 0.7 | (17.3, 20.1) |
| COLORADO | 2986 | 14.9 | 0.9 | (13.0, 16.7) |
| CONNECTICUT | 3320 | 15.1 | 0.8 | (13.5, 16.8) |
| DELAWARE | 2461 | 17.6 | 1.0 | (15.6, 19.5) |
| DISTRICT OF CO | 1242 | 18.5 | 1.2 | (16.1, 20.9) |
| FLORIDA | 5046 | 18.6 | 0.7 | (17.3, 19.9) |
| GEORGIA | 2204 | 21.1 | 1.0 | (19.1, 23.1) |
| HAWAII | 2111 | 15.7 | 1.1 | (13.6, 17.8) |
| IDAHO | 4809 | 20.0 | 0.7 | (18.6, 21.4) |
| ILLINOIS | 2806 | 20.9 | 0.9 | (19.1, 22.8) |
| INDIANA | 2370 | 19.9 | 1.3 | (17.3, 22.5) |
| IOWA | 3528 | 21.5 | 0.8 | (19.9, 23.1) |
| KANSAS | 3680 | 18.9 | 0.7 | (17.5, 20.3) |
| KENTUCKY | 7315 | 21.7 | 0.7 | (20.3, 23.0) |
| LOUISIANA | 1617 | 22.3 | 1.2 | (20.1, 24.6) |
| MAINE | 1613 | 19.4 | 1.1 | (17.3, 21.6) |
| MARYLAND | 3819 | 18.2 | 0.8 | (16.7, 19.7) |
| MASSACHUSETTS | 4689 | 14.7 | 0.7 | (13.4, 16.1) |
| MICHIGAN | 2445 | 22.8 | 1.0 | (20.9, 24.6) |
| MINNESOTA | 5155 | 15.5 | 0.6 | (14.4, 16.5) |
| MISSISSIPPI | 2078 | 23.2 | 1.1 | (21.0, 25.4) |
| MISSOURI | 4161 | 21.7 | 0.9 | (19.9, 23.4) |
| MONTANA | 1743 | 15.8 | 1.0 | (13.9, 17.8) |
| NEBRASKA | 2700 | 21.0 | 0.9 | (19.3, 22.7) |
| NEVADA | 2131 | 15.9 | 1.2 | (13.6, 18.1) |
| NEW HAMPSHIRE | 1192 | 14.6 | 1.2 | (12.2, 16.9) |
| NEW JERSEY | 2677 | 17.1 | 0.9 | (15.2, 18.9) |
| NEW MEXICO | 3378 | 17.7 | 0.7 | (16.3, 19.1) |
| NEW YORK | 2506 | 17.4 | 0.9 | (15.7, 19.1) |
| NORTH CAROLINA | 2331 | 21.5 | 1.0 | (19.6, 23.4) |
| NORTH DAKOTA | 1927 | 21.9 | 1.0 | (19.8, 23.9) |
| OHIO | 1548 | 20.3 | 1.2 | (17.9, 22.6) |
| OKLAHOMA | 2812 | 21.1 | 0.9 | (19.3, 22.8) |
| OREGON | 1741 | 19.9 | 1.0 | (17.9, 22.0) |
| PENNSYLVANIA | 3356 | 20.3 | 0.8 | (18.8, 21.8) |
| RHODE ISLAND | 3802 | 16.8 | 0.7 | (15.4, 18.2) |
| SOUTH CAROLINA | 3263 | 20.7 | 0.8 | (19.1, 22.2) |
| SOUTH DAKOTA | 4001 | 19.6 | 0.7 | (18.2, 21.0) |
| TENNESSEE | 2866 | 20.6 | 0.9 | (18.8, 22.3) |
| TEXAS | 4736 | 21.7 | 0.8 | (20.2, 23.1) |
| UTAH | 3093 | 16.7 | 0.9 | (14.9, 18.6) |
| VERMONT | 3015 | 18.0 | 0.8 | (16.4, 19.6) |
| VIRGINIA | 3372 | 19.3 | 1.0 | (17.4, 21.1) |
| WASHINGTON | 3453 | 18.2 | 0.8 | (16.8, 19.7) |
| WEST VIRGINIA | 2504 | 24.6 | 1.0 | (22.7, 26.5) |
| WISCONSIN | 2117 | 19.9 | 1.0 | (18.0, 21.8) |
| WYOMING | 2343 | 16.9 | 0.8 | (15.3, 18.5) |
| PUERTO RICO | 2894 | 21.3 | 0.9 | (19.6, 23.0) |

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 19.8 RANGE = 12.3-24.6 YEAR 2000 OBJ. 2.3
 * BODY MASS INDEX GE 30.0 AND LE 99.8 FOR MALES AND FEMALES
 June 22, 2000

TABLE 7.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OBESE : BY NEW BODY MASS INDEX* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | -----MALES----- | | | | -----FEMALES----- | | | |
|----------------|-----------------|---------|----------------|-------------------------|-------------------|---------|----------------|-------------------------|
| | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL | SAMPLE SIZE | PERCENT | STANDARD ERROR | 95% CONFIDENCE INTERVAL |
| ALABAMA | 845 | 21.3 | 1.6 | (18.1, 24.5) | 1200 | 23.5 | 1.4 | (20.7, 26.2) |
| ALASKA | 997 | 20.5 | 1.8 | (17.0, 24.1) | 1010 | 20.2 | 1.7 | (17.0, 23.5) |
| ARIZONA | 698 | 14.1 | 1.9 | (10.4, 17.9) | 977 | 10.4 | 1.2 | (8.1, 12.8) |
| ARKANSAS | 1160 | 22.3 | 1.4 | (19.7, 25.0) | 1748 | 23.0 | 1.1 | (20.8, 25.2) |
| CALIFORNIA | 1796 | 17.8 | 1.1 | (15.7, 19.8) | 2356 | 19.7 | 1.0 | (17.8, 21.6) |
| COLORADO | 1359 | 15.3 | 1.4 | (12.5, 18.1) | 1627 | 14.5 | 1.2 | (12.1, 16.9) |
| CONNECTICUT | 1346 | 15.4 | 1.3 | (12.9, 17.9) | 1974 | 14.9 | 1.1 | (12.8, 17.0) |
| DELAWARE | 1018 | 17.4 | 1.6 | (14.3, 20.4) | 1443 | 17.7 | 1.3 | (15.1, 20.3) |
| DISTRICT OF CO | 456 | 15.4 | 1.8 | (11.7, 19.0) | 786 | 21.3 | 1.6 | (18.1, 24.4) |
| FLORIDA | 2029 | 18.4 | 1.0 | (16.4, 20.3) | 3017 | 18.8 | 0.9 | (17.2, 20.5) |
| GEORGIA | 874 | 22.5 | 1.6 | (19.3, 25.6) | 1330 | 19.9 | 1.3 | (17.4, 22.3) |
| HAWAII | 983 | 16.4 | 1.6 | (13.2, 19.5) | 1128 | 15.0 | 1.4 | (12.3, 17.7) |
| IDAHO | 2079 | 21.7 | 1.2 | (19.5, 24.0) | 2730 | 18.2 | 0.9 | (16.5, 19.9) |
| ILLINOIS | 1088 | 21.5 | 1.6 | (18.5, 24.6) | 1718 | 20.3 | 1.1 | (18.2, 22.5) |
| INDIANA | 1010 | 19.1 | 1.9 | (15.3, 22.8) | 1360 | 20.7 | 1.8 | (17.0, 24.3) |
| IOWA | 1409 | 23.1 | 1.3 | (20.5, 25.7) | 2119 | 20.0 | 1.0 | (18.0, 21.9) |
| KANSAS | 1546 | 19.5 | 1.1 | (17.3, 21.7) | 2134 | 18.3 | 0.9 | (16.5, 20.1) |
| KENTUCKY | 2737 | 22.2 | 1.2 | (19.9, 24.5) | 4578 | 21.1 | 0.9 | (19.4, 22.8) |
| LOUISIANA | 658 | 22.6 | 1.8 | (19.0, 26.1) | 959 | 22.1 | 1.5 | (19.2, 24.9) |
| MAINE | 731 | 20.1 | 1.7 | (16.8, 23.4) | 882 | 18.8 | 1.4 | (16.0, 21.6) |
| MARYLAND | 1570 | 16.5 | 1.1 | (14.2, 18.7) | 2249 | 19.9 | 1.0 | (17.8, 21.9) |
| MASSACHUSETTS | 1948 | 17.5 | 1.1 | (15.3, 19.7) | 2741 | 12.1 | 0.8 | (10.6, 13.6) |
| MICHIGAN | 1030 | 23.4 | 1.5 | (20.5, 26.3) | 1415 | 22.2 | 1.2 | (19.8, 24.6) |
| MINNESOTA | 2416 | 17.2 | 0.8 | (15.5, 18.8) | 2739 | 13.8 | 0.7 | (12.4, 15.2) |
| MISSISSIPPI | 843 | 21.1 | 1.8 | (17.6, 24.5) | 1235 | 25.2 | 1.4 | (22.4, 27.9) |
| MISSOURI | 1707 | 22.1 | 1.3 | (19.5, 24.7) | 2454 | 21.2 | 1.1 | (19.0, 23.4) |
| MONTANA | 765 | 17.3 | 1.5 | (14.4, 20.3) | 978 | 14.3 | 1.3 | (11.9, 16.8) |
| NEBRASKA | 1099 | 23.6 | 1.4 | (20.8, 26.3) | 1601 | 18.6 | 1.1 | (16.5, 20.6) |
| NEVADA | 1092 | 14.1 | 1.5 | (11.2, 17.0) | 1039 | 17.7 | 1.8 | (14.2, 21.2) |
| NEW HAMPSHIRE | 496 | 16.5 | 2.0 | (12.7, 20.4) | 696 | 12.6 | 1.4 | (9.9, 15.3) |
| NEW JERSEY | 1103 | 19.5 | 1.5 | (16.5, 22.5) | 1574 | 14.7 | 1.1 | (12.6, 16.8) |
| NEW MEXICO | 1509 | 16.8 | 1.1 | (14.7, 18.9) | 1869 | 18.6 | 1.0 | (16.6, 20.6) |
| NEW YORK | 1049 | 18.0 | 1.3 | (15.3, 20.6) | 1457 | 16.9 | 1.1 | (14.7, 19.1) |
| NORTH CAROLINA | 951 | 21.0 | 1.5 | (18.0, 23.9) | 1380 | 22.0 | 1.2 | (19.6, 24.5) |
| NORTH DAKOTA | 856 | 23.5 | 1.6 | (20.4, 26.6) | 1071 | 20.2 | 1.4 | (17.6, 22.9) |
| OHIO | 629 | 18.9 | 1.7 | (15.5, 22.3) | 919 | 21.6 | 1.6 | (18.5, 24.7) |
| OKLAHOMA | 1119 | 21.6 | 1.4 | (19.0, 24.3) | 1693 | 20.5 | 1.1 | (18.4, 22.6) |
| OREGON | 748 | 20.1 | 1.6 | (17.0, 23.2) | 993 | 19.8 | 1.4 | (17.1, 22.4) |
| PENNSYLVANIA | 1393 | 20.9 | 1.2 | (18.5, 23.2) | 1963 | 19.8 | 1.0 | (17.8, 21.7) |
| RHODE ISLAND | 1530 | 17.6 | 1.1 | (15.4, 19.8) | 2272 | 16.0 | 0.9 | (14.3, 17.7) |
| SOUTH CAROLINA | 1362 | 21.6 | 1.3 | (19.1, 24.1) | 1901 | 19.8 | 1.0 | (17.8, 21.7) |
| SOUTH DAKOTA | 1670 | 20.3 | 1.1 | (18.1, 22.4) | 2331 | 19.0 | 0.9 | (17.3, 20.8) |
| TENNESSEE | 1019 | 19.9 | 1.4 | (17.2, 22.6) | 1847 | 21.2 | 1.1 | (19.1, 23.3) |
| TEXAS | 1951 | 22.4 | 1.1 | (20.2, 24.7) | 2785 | 20.9 | 1.0 | (18.9, 22.9) |
| UTAH | 1364 | 17.3 | 1.4 | (14.5, 20.1) | 1729 | 16.2 | 1.2 | (13.8, 18.6) |
| VERMONT | 1260 | 19.9 | 1.3 | (17.4, 22.5) | 1755 | 16.0 | 1.0 | (14.1, 17.9) |
| VIRGINIA | 1396 | 18.8 | 1.4 | (16.0, 21.6) | 1976 | 19.7 | 1.3 | (17.1, 22.2) |
| WASHINGTON | 1580 | 18.9 | 1.1 | (16.7, 21.1) | 1873 | 17.6 | 1.0 | (15.6, 19.5) |
| WEST VIRGINIA | 1055 | 24.3 | 1.4 | (21.5, 27.1) | 1449 | 24.9 | 1.3 | (22.4, 27.4) |
| WISCONSIN | 915 | 22.8 | 1.6 | (19.7, 25.8) | 1202 | 17.2 | 1.2 | (14.9, 19.4) |
| WYOMING | 1059 | 16.1 | 1.2 | (13.7, 18.5) | 1284 | 17.7 | 1.1 | (15.5, 19.9) |
| PUERTO RICO | 1094 | 21.7 | 1.4 | (18.9, 24.6) | 1800 | 20.9 | 1.1 | (18.8, 23.0) |

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 19.9
 RANGE = 14.1-24.3

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 19.7
 RANGE = 10.4-25.2

* BODY MASS INDEX GE 30.0 AND LE 99.8 FOR MALES AND FEMALES

TABLE 7.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
PREVALENCE DIFFERENCES FROM 1998 TO 1999

OBESSE : BY NEW BODY MASS INDEX*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

| PARTICIPANT | 1998 PERCENT | 1999 PERCENT | PERCENT DIFFERENCE |
|----------------|-----------------|-----------------|-----------------------|
| ALABAMA | 21.3 | 22.4 | 1.2 |
| ALASKA | 21.5 | 20.4 | -1.1 |
| ARIZONA | 13.1 | 12.3 | -0.9 |
| ARKANSAS | 19.8 | 22.7 | 2.8 |
| CALIFORNIA | 17.3 | 18.7 | 1.4 |
| COLORADO | 14.4 | 14.9 | 0.5 |
| CONNECTICUT | 15.5 | 15.1 | -0.4 |
| DELAWARE | 17.2 | 17.6 | 0.3 |
| DISTRICT OF CO | 20.2 | 18.5 | -1.7 |
| FLORIDA | 18.0 | 18.6 | 0.7 |
| GEORGIA | 19.2 | 21.1 | 1.9 |
| HAWAII | 15.5 | 15.7 | 0.2 |
| IDAHO | 16.4 | 20.0 | 3.5 |
| ILLINOIS | 18.5 | 20.9 | 2.4 |
| INDIANA | 19.9 | 19.9 | -0.0 |
| IOWA | 19.8 | 21.5 | 1.7 |
| KANSAS | 17.7 | 18.9 | 1.2 |
| KENTUCKY | 20.4 | 21.7 | 1.2 |
| LOUISIANA | 21.8 | 22.3 | 0.5 |
| MAINE | 17.4 | 19.4 | 2.0 |
| MARYLAND | 20.5 | 18.2 | -2.3 |
| MASSACHUSETTS | 14.3 | 14.7 | 0.4 |
| MICHIGAN | 21.2 | 22.8 | 1.6 |
| MINNESOTA | 16.2 | 15.5 | -0.7 |
| MISSISSIPPI | 22.8 | 23.2 | 0.4 |
| MISSOURI | 20.5 | 21.7 | 1.2 |
| MONTANA | 15.0 | 15.8 | 0.8 |
| NEBRASKA | 18.3 | 21.0 | 2.7 |
| NEVADA | 14.0 | 15.9 | 1.9 |
| NEW HAMPSHIRE | 15.6 | 14.6 | -1.0 |
| NEW JERSEY | 15.5 | 17.1 | 1.5 |
| NEW MEXICO | 15.2 | 17.7 | 2.5 |
| NEW YORK | 16.3 | 17.4 | 1.1 |
| NORTH CAROLINA | 19.4 | 21.5 | 2.1 |
| NORTH DAKOTA | 19.2 | 21.9 | 2.7 |
| OHIO | 20.0 | 20.3 | 0.3 |
| OKLAHOMA | 19.5 | 21.1 | 1.6 |
| OREGON | 18.3 | 19.9 | 1.7 |
| PENNSYLVANIA | 19.4 | 20.3 | 0.9 |
| RHODE ISLAND | 16.8 | 16.8 | -0.0 |
| SOUTH CAROLINA | 20.6 | 20.7 | 0.0 |
| SOUTH DAKOTA | 15.8 | 19.6 | 3.8 |
| TENNESSEE | 19.2 | 20.6 | 1.4 |
| TEXAS | 20.3 | 21.7 | 1.4 |
| UTAH | 15.9 | 16.7 | 0.8 |
| VERMONT | 14.8 | 18.0 | 3.2 |
| VIRGINIA | 18.7 | 19.3 | 0.6 |
| WASHINGTON | 18.1 | 18.2 | 0.1 |
| WEST VIRGINIA | 23.9 | 24.6 | 0.7 |
| WISCONSIN | 18.3 | 19.9 | 1.6 |
| WYOMING | 15.1 | 16.9 | 1.8 |
| PUERTO RICO | 19.3 | 21.3 | 2.0 |

* BODY MASS INDEX GE 30.0 AND LE 99.8 FOR MALES AND FEMALES
N/A - NOT AVAILABLE