

product appear to be better or of greater value than it is. Therefore:

(1) Paprika or oleoresin paprika may not be used in or on fresh meat, such as steaks, or comminuted fresh meat, such as chopped and formed steaks or patties; or in any other meat consisting of fresh meat (with or without seasoning).

(2) Paprika or oleoresin paprika may be used in or on chorizo sausage and other meat in which paprika or oleoresin paprika is permitted as an ingredient in a standard of identity or composition in part 319 of this subchapter.

(3) Sorbic acid, calcium sorbate, sodium sorbate, and other salts of sorbic acid shall not be used in cooked sausages or any other meat; sulfurous acid and salts of sulfurous acid shall not be used in or on any meat; and niacin or nicotinamide shall not be used in or on fresh meat product; except that potassium sorbate, propylparaben (propyl p-hydroxybenzoate), calcium propionate, sodium propionate, benzoic acid, and sodium benzoate may be used in or on any product, only as provided in 9 CFR Chapter III.

(b) *Nitrates*. Nitrates shall not be used in curing bacon.

PART 430—REQUIREMENTS FOR SPECIFIC CLASSES OF PRODUCT

Sec.

430.1 Definitions.

430.4 Control of *Listeria monocytogenes* in post-lethality exposed ready-to-eat products.

AUTHORITY: 7 U.S.C. 450; 7 U.S.C. 1901–1906; 21 U.S.C. 451–470, 601–695; 7 CFR 2.18, 2.53.

SOURCE: 68 FR 34224, June 6, 2003, unless otherwise noted.

§ 430.1 Definitions.

Antimicrobial agent. A substance in or added to an RTE product that has the effect of reducing or eliminating a microorganism, including a pathogen such as *L. monocytogenes*, or that has the effect of suppressing or limiting growth of *L. monocytogenes* in the product throughout the shelf life of the product. Examples of antimicrobial agents added to RTE products are potassium lactate and sodium diacetate.

Antimicrobial process. An operation, such as freezing, applied to an RTE

product that has the effect of suppressing or limiting the growth of a microorganism, such as *L. monocytogenes*, in the product throughout the shelf life of the product.

Deli product. A ready-to-eat meat or poultry product that typically is sliced, either in an official establishment or after distribution from an official establishment, and typically is assembled in a sandwich for consumption.

Hotdog product. A ready-to-eat meat or poultry frank, frankfurter, or wiener, such as a product defined in 9 CFR 319.180 and 319.181.

Lethality treatment. A process, including the application of an antimicrobial agent, that eliminates or reduces the number of pathogenic microorganisms on or in a product to make the product safe for human consumption. Examples of lethality treatments are cooking or the application of an antimicrobial agent or process that eliminates or reduces pathogenic microorganisms.

Post-lethality exposed product. Ready-to-eat product that comes into direct contact with a food contact surface after the lethality treatment in a post-lethality processing environment.

Post-lethality processing environment. The area of an establishment into which product is routed after having been subjected to an initial lethality treatment. The product may be exposed to the environment in this area as a result of slicing, peeling, re-bagging, cooling semi-permeable encased product with a brine solution, or other procedures.

Post-lethality treatment. A lethality treatment that is applied or is effective after post-lethality exposure. It is applied to the final product or sealed package of product in order to reduce or eliminate the level of pathogens resulting from contamination from post-lethality exposure.

Prerequisite program. A procedure or set of procedures that is designed to provide basic environmental or operating conditions necessary for the production of safe, wholesome food. It is called “prerequisite” because it is considered by scientific experts to be prerequisite to a HACCP plan.

Ready-to-eat (RTE) product. A meat or poultry product that is in a form that

is edible without additional preparation to achieve food safety and may receive additional preparation for palatability or aesthetic, epicurean, gastronomic, or culinary purposes. RTE product is not required to bear a safe-handling instruction (as required for non-RTE products by 9 CFR 317.2(l) and 381.125(b)) or other labeling that directs that the product must be cooked or otherwise treated for safety, and can include frozen meat and poultry products.

§ 430.4 Control of *Listeria monocytogenes* in post-lethality exposed ready-to-eat products.

(a) *Listeria monocytogenes* can contaminate RTE products that are exposed to the environment after they have undergone a lethality treatment. *L. monocytogenes* is a hazard that an establishment producing post-lethality exposed RTE products must control through its HACCP plan or prevent in the processing environment through a Sanitation SOP or other prerequisite program. RTE product is adulterated if it contains *L. monocytogenes* or if it comes into direct contact with a food contact surface which is contaminated with *L. monocytogenes*.

(b) In order to maintain the sanitary conditions necessary to meet this requirement, an establishment producing post-lethality exposed RTE product must comply with the requirements included in one of the three following alternatives:

(1) *Alternative 1.* Use of a post-lethality treatment (which may be an antimicrobial agent) that reduces or eliminates microorganisms on the product *and* an antimicrobial agent or process that suppresses or limits the growth of *L. monocytogenes*. If an establishment chooses this alternative:

(i) The post-lethality treatment must be included in the establishment's HACCP plan. The antimicrobial agent or process used to suppress or limit the growth of the pathogen must be included in either the establishment's HACCP plan or its Sanitation SOP or other prerequisite program.

(ii) The establishment must validate the effectiveness of the post-lethality treatment incorporated in its HACCP plan in accordance with § 417.4. The es-

tablishment must document, either in its HACCP plan or in its Sanitation SOP or other prerequisite program, that the antimicrobial agent or process, as used, is effective in suppressing or limiting growth of *L. monocytogenes*.

(2) *Alternative 2.* Use of either a post-lethality treatment (which may be an antimicrobial agent) that reduces or eliminates microorganisms on the product *or* an antimicrobial agent or process that suppresses or limits growth of *L. monocytogenes*. If an establishment chooses this alternative:

(i) The post-lethality treatment must be included in the establishment's HACCP plan. The antimicrobial agent or process used to suppress or limit growth of the pathogen must be included in either the establishment's HACCP plan or its Sanitation SOP or other prerequisite program.

(ii) The establishment must validate the effectiveness of a post-lethality treatment incorporated in its HACCP plan in accordance with § 417.4. The establishment must document in its HACCP plan or in its Sanitation SOP or other prerequisite program that the antimicrobial agent or process, as used, is effective in suppressing or limiting growth of *L. monocytogenes*.

(iii) If an establishment chooses this alternative and chooses to use only an antimicrobial agent or process that suppresses or limits the growth of *L. monocytogenes*, its sanitation program must:

(A) Provide for testing of food contact surfaces in the post-lethality processing environment to ensure that the surfaces are sanitary and free of *L. monocytogenes* or of an indicator organism;

(B) Identify the conditions under which the establishment will implement hold-and-test procedures following a positive test of a food-contact surface for *L. monocytogenes* or an indicator organism;

(C) State the frequency with which testing will be done;

(D) Identify the size and location of the sites that will be sampled; and

(E) Include an explanation of why the testing frequency is sufficient to ensure that effective control of *L. monocytogenes* or of indicator organisms is maintained.