

§ 381.92

§ 381.92 **Overscald.**

Carcasses of poultry which have been overscalded, resulting in a cooked appearance of the flesh, shall be condemned.

§ 381.93 **Decomposition.**

Carcasses of poultry deleteriously affected by post mortem changes shall be disposed of as follows:

(a) Carcasses which have reached a state of putrefaction or stinking fermentation shall be condemned.

(b) Any part of a carcass which is green struck shall be condemned and, if the carcass is so extensively affected that removal of affected parts is impracticable, the whole carcass shall be condemned.

(c) Carcasses affected by types of post mortem change which are superficial in nature may be passed for human food after removal and condemnation of the affected parts.

§ 381.94 **Contamination with Microorganisms; process control verification criteria and testing; pathogen reduction standards.**

(a) Criteria for verifying process control; *E. coli* testing.

(1) Each official establishment that slaughters poultry shall test for *Escherichia coli* Biotype I (*E. coli*). Establishments that slaughter more than one type of poultry and/or poultry and livestock, shall test the type of poultry or livestock slaughtered in the greatest number. The establishment shall:

(i) Collect samples in accordance with the sampling techniques, methodology, and frequency requirements in paragraph (a)(2) of this section;

(ii) Obtain analytic results in accordance with paragraph (a)(3) of this section; and

(iii) Maintain records of such analytic results in accordance with paragraph (a)(4) of this section.

(2) Sampling requirements.

(i) *Written procedures.* Each establishment shall prepare written specimen collection procedures which shall identify employees designated to collect samples, and shall address location(s) of sampling, how sampling randomness is achieved, and handling of the sample to ensure sample integrity. The writ-

ten procedure shall be made available to FSIS upon request.

(ii) *Sample collection.* A whole bird must be taken from the end of the chilling process. If this is impracticable, the whole bird can be taken from the end of the slaughter line. Samples must be collected by rinsing the whole carcass in an amount of buffer appropriate for that type of bird. Samples from turkeys or ratites also may be collected by sponging the carcass on the back and thigh.¹

(iii) *Sampling frequency.* Slaughter establishments, except very low volume establishments as defined in paragraph (a)(2)(v) of this section, must take samples at a frequency proportional to the establishment's volume of production at the following rates:

(A) Chickens: 1 sample per 22,000 carcasses, but a minimum of one sample during each week of operation.

(B) Turkeys, Ducks, Geese, Guineas, Squabs, and Ratites: 1 sample per 3,000 carcasses, but at a minimum one sample each week of operation.

(iv) *Sampling frequency alternatives.* An establishment operating under a validated HACCP plan in accordance with § 417.2(b) of this chapter may substitute an alternative frequency for the frequency of sampling required under paragraph (a)(2)(iii) of this section if,

(A) The alternative is an integral part of the establishment's verification procedures for its HACCP plan and,

(B) FSIS does not determine, and notify the establishment in writing, that the alternative frequency is inadequate to verify the effectiveness of the establishment's processing controls.

(v) *Sampling in very low volume establishments.* (A) Very low volume establishments annually slaughter no more than 440,000 chickens, 60,000 turkeys, 60,000 ducks, 60,000 geese, 60,000 guineas, 60,000 squabs, 6,000 ratites, or a combination of all types of poultry not exceeding 60,000 turkeys and 440,000 birds total. Very low volume establishments that slaughter turkeys, ducks,

¹A copy of FSIS's "Guidelines for *Escherichia coli* Testing for Process Control Verification in Poultry Slaughter Establishments," and "FSIS Turkey Microbiological Procedures for Sponge Sample Collection and Methods of Analysis" are available for inspection in the FSIS Docket Room.