

**Environmental Protection Agency**

**§ 180.368**

Finding Aids section of the printed volume and on GPO Access.

**§ 180.367 n-Octyl bicycloheptenedicarboximide; tolerances for residues.**

(a) *General.* (1) Tolerances are established for residues of the insecticide *n*-octyl bicycloheptene-dicarboximide, resulting from dermal application, in food commodities as follows:

Commodity	Parts per million
Cattle, fat	0.3
Goat, fat	0.3
Hog, fat	0.3
Horse, fat	0.3
Milk, fat	0.3
Sheep, fat	0.3

(2) *N*-octylbicycloheptene dicarboximide may be safely used in accordance with the following prescribed conditions:

(i) It is used in combination with piperonyl butoxide and pyrethrins for insect control in food-processing and food-storage areas, provided that the food is removed or covered prior to such use.

(ii) Residues in food resulting from the use described in paragraph (a)(2)(i) of this section shall not exceed 10 parts per million of *N*-octylbicycloheptene dicarboximide, 10 parts per million of piperonylbutoxide, and 1 part per million of pyrethrins.

(iii) To assure safe use of the pesticide, its label and labeling shall conform to that registered with the U.S. Environmental Protection Agency and it shall be used in accordance with such label and labeling.

(b) *Section 18 emergency exemptions.* [Reserved]

(c) *Tolerances with regional registrations.* [Reserved]

(d) *Indirect or inadvertent residues.* [Reserved]

[65 FR 33713, May 24, 2000]

**§ 180.368 Metolachlor; tolerances for residues.**

(a) *General.* (1) Tolerances are established for the combined residues (free and bound) of the herbicide metolachlor [2-chloro-*N*-(2-ethyl-6-methylphenyl)-*N*-(2-methoxy-1-methylethyl)acetamide] and its metabolites, determined as the derivatives,

2-[(2-ethyl-6-methylphenyl)amino]-1-propanol and 4-(2-ethyl-6-methylphenyl)-2-hydroxy-5-methyl-3-morpholinone, each expressed as the parent compound, in or on the raw agricultural commodities.

Commodity	Parts per million
Almond, hulls	0.3
Barley, fodder	0.5
Barley, grain	0.1
Buckwheat, grain	0.1
Cabbage	1.0
Cattle, fat	0.02
Cattle, kidney	0.2
Cattle, liver	0.05
Cattle, meat	0.02
Cattle, meat byproducts, except kidney and liver	0.02
Celery	0.1
Corn, fresh, kernel plus cob with husks removed	0.1
Corn, forage and fodder	8.0
Corn, grain	0.1
Cotton, undelinted seed	0.1
Egg	0.02
Fruit, stone, group 12	0.1
Goat, fat	0.02
Goat, kidney	0.2
Goat, liver	0.05
Goat, meat	0.02
Goat, meat byproducts, except kidney and liver	0.02
Horse, fat	0.02
Horse, kidney	0.2
Horse, liver	0.05
Horse, meat	0.02
Horse, meat byproducts, except kidney and liver	0.02
Milk	0.02
Millet, fodder	0.5
Millet, forage	0.5
Millet, grain	0.1
Milo, fodder	0.5
Milo, forage	0.5
Milo, grain	0.1
Nongrass animal feed (forage, fodder, straw, and hay) group	3.0
Nut, tree, group 14	0.1
Oat, forage	0.5
Oat, grain	0.1
Oat, straw	0.5
Peanut	0.5
Peanut, forage and hay	30.0
Pepper, bell	0.1
Potato	0.2
Poultry, fat	0.02
Poultry, liver	0.05
Poultry, meat	0.02
Poultry, meat byproducts, except liver	0.02
Rice, forage	0.5
Rice, straw	0.5
Rice, grain	0.1
Rye, forage	0.5
Rye, grain	0.1
Rye, straw	0.5
Safflower, seed	0.1
Seed and pod vegetables (except soybean)	0.3
Sheep, fat	0.02
Sheep, kidney	0.2
Sheep, liver	0.05
Sheep, meat	0.02
Sheep, meat byproducts, except kidney and liver	0.02
Sorghum, forage and fodder	2.0
Sorghum, grain	0.3
Soybean	0.2