

Polychlorinated Biphenyls (PCBs)

CAS Number: 1336-36-3

What are PCBs?

Polychlorinated biphenyls (PCBs) were developed in the 1940's and were used extensively in the manufacture of transformers, capacitors, and other heat transfer devices through the late 1970's. PCBs are a group of chemicals that have extremely high boiling points and are practically nonflammable. Because of this, they were used extensively as heat transfer fluids in transformers and capacitors. In 1979 their manufacture and importation was banned in the United States, based on mounting evidence that they were toxic to humans and wildlife. Today they are classified as probable human carcinogens and are listed in the top 10% of EPA's most toxic chemicals. At older federal facilities today, there are still many PCB-containing transformers and capacitors. Many of these transformers are at or close to the end of their useful life and proper disposal of them is required to prevent the unnecessary (and unlawful) release of PCBs into the environment.

How might I be exposed to PCBs?

Fish consumption appears to be the major pathway of exposure. PCBs do not easily break down, and can bioaccumulate in the fatty tissues of fish and mammals. A significant trend of increasing body burden is associated with increased fish consumption. People who

eat sport-caught fish consumed 2-3 times more fish than the overall U.S. population.

Concentrations of PCBs in subsurface soil at a Superfund site have been as high as 750 ppm. People who live near hazardous waste sites may be exposed to PCBs by consuming PCB-contaminated sportfish and game animals, by breathing PCBs in air, or by drinking PCB-contaminated well water.

Although PCBs are no longer made in the United States, people can still be exposed to them. Many older transformers and capacitors may still contain PCBs, and this equipment can be used for 30 years or more. Old fluorescent lighting fixtures and old electrical devices and appliances, such as television sets and refrigerators, may contain PCBs if they were made before PCB use was stopped. When these electric devices get hot during operation, small amounts of PCBs may get into the air and raise the level of PCBs in indoor air. Because devices that contain PCBs can leak with age, they could also be a source of skin exposure to PCBs.

Workplace exposure to PCBs can occur during the repair and maintenance of PCB transformers, accidents, fires, or spills involving PCB transformers and older computers and instruments, and disposal of PCB materials. In addition to older electrical instruments and fluorescent lights that contain

PCB-filled capacitors, caulking materials, elastic sealants, and heat insulation have also been known to contain PCBs. Contact with PCBs at hazardous waste sites can happen when workers breathe air and touch soil containing PCBs. Exposure in the contaminated workplace occurs mostly by breathing air containing PCBs and by touching substances that contain PCBs.

How can PCBs enter and leave my body?

If you breathe air that contains PCBs, they can enter your body through your lungs and pass into the bloodstream. We do not know how fast or how much of the PCBs that are breathed will pass into the blood. A common way for PCBs to enter your body is by eating meat or fish products or other foods that contain PCBs. Exposure from drinking water is less than from food. It is also possible that PCBs can enter your body by breathing indoor air or by skin contact in buildings that have the kinds of old electrical devices that contain and can leak PCBs. For people living near waste sites or processing or storage facilities, and for people who work with or around PCBs, the most likely ways that PCBs will enter their bodies are from skin contact with contaminated soil and from breathing PCB vapors. Once PCBs are in your body, some may be changed by your body into other related chemicals called metabolites. Some metabolites of PCBs may have the potential to be as harmful as some unchanged PCBs. Some of the metabolites may leave your body in the feces in a few days, but others may remain in your body fat for months. Unchanged PCBs may also remain in your body and be stored for years mainly in the fat and liver, but smaller amounts can be found in other organs as well. PCBs collect in milk fat and can enter the bodies of infants through breast-feeding.

How can PCBs affect my health?

Many studies have looked at how PCBs can affect human health. Some of these studies investigated people exposed in the workplace, and others have examined members of the general population. Skin conditions, such as acne and rashes, may occur in people exposed to high levels of PCBs. These effects on the skin are well documented, but are not likely to result from exposures in the general population. Most of the human studies have many shortcomings, which make it difficult for scientists to establish a clear association between PCB exposure levels and health effects. Some studies in workers suggest that exposure to PCBs may also cause irritation of the nose and lungs, gastrointestinal discomfort, changes in the blood and liver, and depression and fatigue. Workplace concentrations of PCBs, such as those in areas where PCB transformers are repaired and maintained, are higher than levels in other places, such as air in buildings that have electrical devices containing PCBs or in outdoor air, including air at hazardous waste sites. Most of the studies of health effects of PCBs in the general population examined children of mothers who were exposed to PCBs.

Is there a medical test to determine if I have been exposed to PCBs?

Levels of PCBs in the environment were zero before PCBs were manufactured. Now, all people in industrial countries have some PCBs in their bodies. There are tests to determine whether PCBs are in the blood, body fat, and breast milk. These are not regular or routine clinical tests, such as the one for cholesterol, but could be ordered by a doctor to detect PCBs in people exposed to them in the environment and at work. If your PCB levels are higher than the background levels, this will show that you have been exposed to high

levels of PCBs. However, these measurements cannot determine the exact amount or type of PCBs that you have been exposed to, or how long you have been exposed. Although these tests can indicate whether you have been exposed to PCBs to a greater extent than the general population, they do not predict whether you will develop harmful health effects. Blood tests are the easiest, safest, and probably the best method for detecting recent exposures to large amounts of PCBs. Results of such tests should be reviewed and carefully interpreted by physicians with a background in environmental and occupational medicine. Nearly everyone has been exposed to PCBs because they are found throughout the environment, and people are likely to have detectable amounts of PCBs in their blood, fat, and breast milk. Recent studies have shown that PCB levels in tissues from United States population are now declining.

What levels of exposure have resulted in harmful health effects?

Human health studies indicate that: 1) reproductive function may be disrupted by high levels of exposure to PCBs; 2) neurobehavioral and developmental deficits occur in newborns and continue through school-aged children who had in-utero exposure to PCBs; 3) other systemic effects (e.g., self-reported liver disease and diabetes, and effects on the thyroid and immune systems) are associated with elevated serum levels of PCBs; and 4) increased cancer risks, e.g., non-Hodgkin's lymphoma, are associated with PCB exposures.

What recommendations has the federal government made to protect human health?

The federal government has developed regulations, guidelines, and standards to protect people from the possible health effects of exposure to PCBs.

The federal government develops regulations and recommendations to protect public health. Regulations can be enforced by law. Federal agencies that develop regulations for toxic substances include the Environmental Protection Agency (EPA), the Occupational Safety and Health Administration (OSHA), and the Food and Drug Administration (FDA). Recommendations provide valuable guidelines to protect public health but cannot be enforced by law. Federal organizations that develop recommendations for toxic substances include the Agency for Toxic Substances and Disease Registry (ATSDR) and the National Institute for Occupational Safety and Health (NIOSH).

Regulations and recommendations can be expressed in not-to-exceed levels in air, water, soil, or food that are usually based on levels that affect animals; then they are adjusted to help protect people. Sometimes these not-to-exceed levels differ among federal organizations because of different exposure times (an 8-hour workday or a 24-hour day), the use of different animal studies, or other factors.

Recommendations and regulations are periodically updated as more information becomes available. For the most current information, check with the federal agency or organization that provides it. Some regulations and recommendations for PCBs include the following:

The EPA standard for PCBs in drinking water is 0.5 parts of PCBs per billion parts (ppb) of water. For the protection of human health from the possible effects of drinking the water or eating the fish or shellfish from lakes and streams that are contaminated with PCBs, the EPA regulates that the level of PCBs in these waters be no greater than 0.17 parts of PCBs per trillion parts (ppt) of water.

The FDA has set residue limits for PCBs in

various foods to protect from harmful health effects. FDA required limits include 0.2 parts of PCBs per million parts (ppm) in infant and junior foods, 0.3 ppm in eggs, 1.5 ppm in milk and other dairy products (fat basis), 2 ppm in fish and shellfish (edible portions), and 3 ppm in poultry and red meat (fat basis).

OSHA regulates that workers not be exposed by inhalation over a period of 8 hours for 5 days per week to more than 1 milligram per cubic meter of air (mg/m³) for 42% chlorine PCBs, or to 0.5 mg/m³ for 54% chlorine PCBs.

NIOSH recommends that workers not breathe air containing 42 or 54% chlorine PCB levels higher than 1 microgram per cubic meter of air (µg/m³) for a 10-hour workday, 40-hour workweek.

EPA requires that companies that transport, store, or dispose of PCBs follow the rules and regulations of the federal hazardous waste management program. EPA also limits the amount of PCBs put into publicly owned waste water treatment plants. To minimize exposure of people to PCBs, EPA requires that industry tell the National Response Center each time one pound or more of PCBs has been released to the environment.

Where can I get more information?

If you have more questions or concerns, please contact your state health or environmental department or:

Agency for Toxic Substances and Disease
Registry
Division of Toxicology
1600 Clifton Road, E-29
Atlanta, Georgia 30333

References

1. Agency for Toxic Substances and Disease Registry (ATSDR). *Public Health Statement for*

PCBs.

Atlanta, GA: U.S. Department of Health and Human Services, 1990.

2. Birmingham et al. 1989
3. Newhook 1988
4. Fitzgerald et al. 1996
5. Courval et al. 1996