

2006 National Youth Tobacco Survey and Key Prevalence Indicators

TABLE 2. Percentage of students in high school (grades 9-12) who were current users* of any tobacco product, by product type, sex, and race/ethnicity — United States, 2004 and 2006

Characteristic	Any tobacco†		Cigarettes		Cigars		Smokeless tobacco		Pipes		Bidis		Kreteks	
	%	(95% CI§)	%	(95% CI§)	%	(95% CI§)	%	(95% CI§)	%	(95% CI§)	%	(95% CI§)	%	(95% CI§)
High school, 2006														
Sex														
Male	30.2	(28.1-32.3)	21.2	(19.4-23.0)	16.6	(15.3-17.9)	11.0	(9.1-12.9)	5.6	(4.8-6.4)	3.3	(2.8-3.8)	3.6	(2.9-4.3)
Female	21.3	(19.2-23.4)	18.4¶	(16.4-20.4)	7.2	(6.3-8.1)	1.5	(1.1-1.9)	1.8	(1.3-2.3)	2.4¶	(2.0-2.8)	2.0	(1.5-2.5)
Race/Ethnicity														
White, non-Hispanic	28.4	(26.0-30.8)	22.3	(20.2-24.4)	12.5	(11.4-13.6)	7.5	(6.1-8.9)	3.6	(3.0-4.2)	2.6	(2.2-3.0)	2.9	(2.3-3.5)
Black, non-Hispanic	15.7	(13.5-17.9)	9.9	(8.1-11.7)	8.9	(7.2-10.6)	1.8	(1.2-2.4)	2.2	(1.5-2.9)	2.7	(2.0-3.4)	1.7	(1.1-2.3)
Hispanic	24.7	(22.2-27.2)	18.8	(16.8-20.8)	12.4	(10.8-14.0)	4.6	(3.3-5.9)	5.3	(4.2-6.4)	4.6	(3.8-5.4)	3.2	(2.4-4.0)
Asian	9.2	(6.6-11.8)	7.3¶	(5.1-9.5)	4.8	(2.7-6.9)	1.5	(0.3-2.7)	1.5	(0.5-2.5)	1.2	(0.1-2.3)	1.4	(0.4-2.4)
Total	25.6	(23.8-27.4)	19.7	(18.0-21.4)	11.8	(11.0-12.6)	6.1	(5.1-7.1)	3.7	(3.2-4.2)	2.9	(2.5-3.3)	2.8	(2.3-3.3)
High school, 2004														
Sex														
Male	30.7	(28.0-33.4)	21.6	(19.2-24.0)	18.4	(16.6-20.2)	9.9	(8.0-11.8)	4.8	(4.0-5.6)	3.7	(3.0-4.4)	3.4	(2.7-4.1)
Female	24.1	(21.6-26.6)	21.8	(19.3-24.3)	7.6	(6.5-8.7)	1.2	(0.8-1.6)	1.5	(1.0-2.0)	1.6	(1.2-2.0)	1.6	(1.1-2.1)
Race/Ethnicity														
White, non-Hispanic	30.8	(27.6-34.0)	24.8	(21.8-27.8)	13.7	(12.0-15.4)	6.9	(5.5-8.3)	3.1	(2.4-3.8)	2.4	(1.9-2.9)	2.5	(1.9-3.1)
Black, non-Hispanic	16.8	(14.3-19.3)	10.9	(8.4-13.4)	10.0	(8.4-11.6)	1.4	(0.7-2.1)	1.7	(1.0-2.4)	2.1	(1.4-2.8)	1.3	(0.7-1.9)
Hispanic	25.7	(23.4-28.0)	20.5	(18.2-22.8)	13.5	(11.9-15.1)	3.2	(2.4-4.0)	4.8	(3.9-5.7)	4.8	(4.0-5.6)	3.6	(2.8-4.4)
Asian	13.3	(10.0-16.6)	11.3	(8.7-13.9)	6.2	(3.7-8.7)	2.0	(0.7-3.3)	2.1	(1.0-3.2)	2.1	(1.0-3.2)	1.3	(0.4-2.2)
Total	27.4	(25.0-29.8)	21.7	(19.5-23.9)	12.9	(11.6-14.2)	5.5	(4.4-6.6)	3.2	(2.7-3.7)	2.7	(2.3-3.1)	2.5	(2.1-2.9)

* Used tobacco on on ≥ 1 of the 30 preceding days.

† Cigarettes, cigars, smokeless tobacco, pipes, bidis (leaf-wrapped, flavored cigarettes from India), or kreteks (clove cigarettes).

§ Confidence interval.

¶ Significant difference ($p < 0.05$), 2006 versus 2004.