









U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES

National Institutes of Health

National Heart, Lung, and Blood

Women's Healt

## The Heart Truth Professional Education Campaign Lecture Component: References

www.womenshealth.gov/hearttruth

ACOG Task Force for Hormone Therapy. Summary of Balancing Risks and Benefits. Obstet Gynecol 2004; 104 (4 Suppl): 1S-129S.

Ades PA, Savage PD, Tischler MD, et al. Determinants of disability in older coronary patients. Am Heart J 2002; 143:151-156.

American Diabetes Association. Screening for Diabetes. Diabetes Care 2001; 24 Suppl 1:521-4.

American Heart Association. Heart Disease and Stroke Statistics 2004.

Anderson GL, Limacher M, Assaf AR, et al. Effects of conjugated equine estrogen in postmenopausal women with hysterectomy: the Women's Health Initiative randomized controlled trial. JAMA 2004; 291:1701-1712.

Antiplatelet Trialists' Collaboration. Collaborative overview of randomized trials of antiplatelet therapies in various categories of patients. BMJ 1994; 308:81-106.

Braunwald E, Antman EM, Beasley JW, et al. ACC/AHA 2002 guideline update for the management of patients with unstable angina and non-ST-segment elevation myocardial infarction: summary article; a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on the Management of Patients with Unstable Angina). J Am Coll Cardiol 2002; 40:1366-1374.

Burt V, Whelton P, Roccella EJ, et al. Prevalence of hypertension in the US adult population: results from the Third National Health and Nutrition Examination Survey, 1988-1991. Hypertension 1995; 25:305-313.

Carr MC, Kim KH, Zambon A, et al. Changes in LDL density across the menopausal transition. J Investig Med 2000; 48: 245-250.

Centers for Disease Control and Prevention, Department of Health and Human Services. National Diabetes Fact Sheet, 2003.

Chandra NC, Ziegelstein RC, Rogers WJ, et al. Observations of the treatment of women in the United States with myocardial infarction; a report from the National Registry of Myocardial Infarction-I. Arch Intern Med 1998; 158:981-988.

Charney P. Presenting symptoms and diagnosis of coronary heart disease in women. J Cardiovasc Risk 2002; 9:303-307.

Cheitlin MD, Armstrong WF, Aurigemma GP, et al. ACC/AHA/ASE 2003 guideline update for the clinical application of echocardiography: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/ASE Committee to Update the 1997 Guidelines for the Clinical Application of Echocardiography). J Am Soc Echocardiogr 2003; 16:1091-1110

Coronary heart disease incidence, by sex—United States, 1971-1987. MMWR 1992; 41:526-529.

DePuey EG, M.D., Image: Breast Attenuation.

Devries S, Wokfkiel C, Fusman B, et al. Influence of age and gender on the presence of coronary calcium detected by ultrafast computed tomography. J Am Coll Cardiol 1995; 25:76-82..

Doherty TM, Tang W, Detrano RC. Racial differences in the significance of coronary calcium in asymptomatic black and white subjects with coronary risk factors. J Am Coll Cardiol 1999; 35:787-794.

Downs JR, Clearfield M, Weis S, et al. Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels: results of AFCAPS/TexCAPS. JAMA 1998; 279:1615-1622.

Duvernoy C, M.D., personal communication.

Elhendy A, Geleijnse ML, van Domburg RT, et al. Gender differences in the accuracy of dobutamine stress echocardiography for the diagnosis of coronary artery disease. Am J Cardiol 1997; 80: 1414-1418.

Eriksson M, Egbert Nils, Wamala S, et al. Relationship between plasma fibrinogen and coronary heart disease in women. Arterioscler Thromb Vasc Biol 1999.;19:67-72.

Espeland MA, Hogan PE, Fineberg SE, et al. Effect of post-menopausal therapy on glucose and insulin concentrations. Diabetes Care 1998; 21:1589-1595.

Facts about the DASH eating plan. Bethesda, Md: National Heart, Lung, and Blood Institute 1998. NIH publication no. 03-4082.

FDA approves new labels for estrogen and estrogen plus progestin therapies for postmenopausal women following review of Women's Health Initiative data. Rockville, Md: Food and Drug Administration, January 8, 2003, available at: <a href="http://www.fda.gov/bbs/topics/NEWS/2003/NEW00863.html">http://www.fda.gov/bbs/topics/NEWS/2003/NEW00863.html</a>, accessed July 26, 2005.

Fields LE, Burt VL, Cutler J, et al. The burden of adult hypertension in the United States 1999-2000: a rising tide. Hypertension 2004; 44:398-404.

Fiore MC, Bailey WC, Cohen SJ, et al. Treating tobacco use and dependence. Clinical Practice Guideline. Rockville, MD: U.S. Department of Health and Human Services. Public Health Service. June 2000.

Garber AM, Browner WS, Hulley SB. Cholesterol screening in asymptomatic adults, revisited. Ann Int Med 1996; 124:518-531.

Gibbons RJ, Balady GJ, Bricker JT, et al. ACC/AHA 2002 guidelines update for exercise testing: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1997 Exercise Testing Guidelines). J Am Coll Cardiol 2002;40:1531-1540.

Goldberg RJ, O'Donnell C, Yarzebski J, et al. Sex differences in symptoms presentation associated with acute myocardial infarction: a population-based perspective. Am Heart J 1998. 136; 189-195.

Grady D, Rubin SM, Petitti DB, et al. Hormone therapy to prevent disease and prolong life in postmenopausal women. Ann Intern Med 1992; 117: 1016-1037.

Grady D. Diagnosis and treatment of coronary heart disease in women: systematic reviews of evidence on selected topics. U.S. Department of Health and Human Services, Public Health Service, Agency for Healthcare Research and Quality, Evidence Report/Technology Assessment Number 81, 2003.

Grundy SM, Cleeman JI, Bairey Merz N, et al. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. Circulation 2004; 110:227-239.

Grundy, SM, Cleeman JI, Daniels SR et al. Diagnosis and management of metabolic syndrome: an American Heart Association/National Heart, Lung and Blood Institute scientific statement. Accessed at http://www.circulationaha.org on October 25, 2005

Hansen CL, Crabbe D, Rubin S. Lower diagnostic accuracy of thallium-201 SPECT myocardial perfusion imaging in women: an effect of smaller chamber size. J Am Coll Cardiol 1996; 28: 1214-1219.

Harpaz D, Benderly M, Goldbourt U, et al. Effect of aspirin on mortality in women with symptomatic or silent myocardial ischemia. Am J Cardiol 1996; 78:1215-1219.

Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of cholesterol lowering with simvastatin in 20,536 high-risk individuals: a randomised placebocontrolled trial. Lancet 2002; . 360:7-22.

Hendrix KH, Riehle JE, Egan BM. Ethnic, gender, and age-related differences in treatment and control of dyslipidemia in hypertensive patients. Ethn Dis 2005; 15:11-16.

Hokanson JE, Austin MA. Plasma triglyceride level is a risk factor for cardiovascular disease independent of high-density lipoprotein cholesterol level: a meta-analysis of population-based prospective studies. J Cardiovasc Risk 1996; 3: 213-219.

Hulley S, Grady D, Bush T, et al. Randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women. Heart and Estrogen/progestin Replacement Study (HERS) Research Group. JAMA 1998; 280: 605-613.

ISIS-2 Collaborative Group. Randomised trial of intravenous streptokinase, oral aspirin, both, or neither. Lancet 1988; 8607:349-360.

Jacobs AK. Coronary revascularization in women in 2003: sex revisited. Circulation 2003; 107:375-377.

Jacobs AK, Kelsey SF, Yeh W, et al. Documentation of decline in morbidity in women undergoing coronary angioplasty (a report from the 1993-94 NHLBI percutaneous transluminal coronary angioplasty registry). Am J Cardiol 1997; 80:979-984.

Keil U. Coronary artery disease: the role of lipids, hypertension, and smoking. Basic Res Cardiol 2000; 95: I/52-I/58.

Klag MJ, Whelton PK, Randall BL et al. End-stage renal disease in African American and white men. 16-year MRFIT findings. JAMA 1997; 277:1293-1298.

Kris-Etherton PM, Harris WS, Appel LJ, et al. Omega-3 fatty acids and cardiovascular disease. Attrioscler Thromb Vasc Biol 2003; 23:151-152.

Krolewski AS, Warram JH, Christlieb AR, et al. Evolving natural history of coronary artery disease in diabetes mellitus. Am J Med 1991; 90: 56S-61S.

Kwok Y, Kim C, Grady D, et al. Meta-analysis of exercise testing to detect coronary artery disease in women. Am J Cardiol 1999; 83:660-666.

Lange H, Suryapranata H, De Luca G, et al. Folate therapy and in-stent restenosis after coronary stenting. N Engl J Med 2004; 350:2673-2681.

Lauzon C, Dupuis R, Beck CA, et al. Depression and prognosis following hospital admission because of acute myocardial infarction. CMAJ 2003; 168: 547-552.

Lee IM, Cook NR, Gaziano JM, et al. Vitamin E in the primary prevention of cardiovascular disease and cancer. The Women's Health Study: a randomized controlled trial. JAMA 2005; 294:56-65.

Leim A, Reynierse-Buitenwerf GH, Zwinderman AH, et al. Secondary prevention with folic acid: effects on clinical outcomes. J Am Coll Cardiol 2003; 41:2105-2113.

Limacher MC. Hormones and heart disease: what we thought, what we have learned, and what we still need to know. Trans Am Clin Climatol Assoc 2002. 113: 31-41.

Lonn E, Bosch J, Yusuf S, et al. Effects of long-term vitamin E supplementation on cardiovascular events and cancer; a randomized controlled trial. JAMA 2005; 293:1338-1347.

Manson JE, Hu FB, Rich-Edwards JW, et al. A prospective study of walking as compared with vigorous exercise in the prevention of coronary heart disease in women. N Engl J Med 1999; 341:650-658.

Manson JE, Willett WC, Stampfer MJ, Colditz GA, Hunter DJ, Hankinson SE, Hennekens CH, Speizer FE. Body weight and mortality among women. N Engl J Med 1995. 333:677-685.

Maron DJ. The epidemiology of low levels of high-density lipoprotein cholesterol in patients with and without coronary artery disease. Am J Cardiol 2000; 86:11L-14L.

Marwick TH, Anderson T, Williams MJ. Exercise echocardiogarphy is an accurate and cost-effectcient technique for detection of coronary artery disease in women. J Am Coll Cardiol 1995; 26: 335-341.

Mendelsohn ME, Karas RH. The protective effects of estrogen on the cardiovascular system. N Engl J Med 1999; 340:1801-1811.

Mosca L, Appel LJ, Benjamin EJ, et al. Evidence-based guidelines for cardiovascular disease prevention in women. Circulation 2004; 109:672-693.

Mosca L, Ferris A, Fabunmi R, et al. Tracking women's awareness of heart disease. Circulation. 2004;109:573-579.

Narayan KM, Boyle JP, Thompson TJ, et al. Lifetime risk for diabetes mellitus in the United States JAMA 2003; 290:1884-1890.

National Vital Statistics System, Health, United States, 1996-97.

Nishimura RA, Gibbons RJ, Blockner JF, Tajik AJ.
Noninvasive cardiac imaging: echocardiography, nuclear cardiology, and MRI/CT imaging in DL Kasper, E Braunwald, AS Fauci, SL Hauser, DL Longo, JL Jameson, eds. Harrison's Principles of Internal Medicine, 16th ed. New York; McGraw Hill, 2005.

Nohria A, Vaccarino V, Krumholz HM. Gender differences in coronary artery disease in women: gender differences in mortality after myocardial infarction: why women fare worse than men. Cardiol Clin 1998; 16:45-57.

O'Meara JG, Kardia SL, Armon JJ, et al. Ethnic and sex differences in the prevalence, treatment, and control of dyslipidemia among hypertensive adults in the GENOA study. Arch Intern Med 2004;164:1313-1318.

O'Rourke RA, Bundate BH, Froelicher VF, et al. American College of Cardiology/American Heart Association expert consensus document on electron-beam computed tomography for the diagnosis and prognosis of coronary artery disease. J Am Coll Cardiol 2000; 36:326-340.

Orth-Gomer K, Wamala SP, Horsten M, et al. Marital stress worsens prognosis in women with coronary heart disease: The Stockholm Female Coronary Risk Study. JAMA. 2000; 283: 3008-14.

Petro KR, Dullum KM, Garcia JM, et al. Minimally invasive coronary revascularization in women: a safe approach in a high risk group. Heart Surg Forum 2000; 3:41-46.

Pires LA, Sethuraman B, Guduguntla VD, et al. Outcome of women versus men with ventricular tachyarrythmias treated with the implantable cardioverter defibrillator. J Cardiovasc Electrophysiol 2002; 13:563-568.

Pregler J, M.D., personal communication, March, 2005

Prescott E, Hippe M, Schnohr P, et al. Smoking and risk of myocardial infarction in women and men: longitudinal population study. BMJ 1998; 316:1043-1047.

Prochaska JO, Diclemente CC, Norcross JC. In search of how people change: applications to addictive behavior. American Psychologist 1992; 47:1102-1114.

Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S) Lancet. 1994. 344: 1383-1389.

Ridker PM, Cook NR, Lee I, et al. A randomized trial of low-dose aspirin in the primary prevention of cardiovascular disease in women. N Engl J Med 2005; 352:1293-1304.

Ridker PM, Hennekens CH, Buring JE, et al. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. N Engl J Med 2000. 342:836-843.

Ritchie JL, Bateman TM, Bonow RO, et al. Special report: guidelines for clinical use of cardiac radionuclide imaging: a report of the AHA/ACC task force on assessment of diagnostic and therapeutic cardiovascular procedures, committee on radionuclide imaging, developed in collaboration with the American Society of Nuclear Cardiology. Circulation 1995; 91:1278-1303.

Rossouw JE, Anderson GL, Prentice RL, et al. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results from the Women's Health Initiative randomized controlled trial. JAMA 2002; 288: 321-333.

Sabatine MS, Cannon CP, Gibson CM, et al. Addition of clopidogrel to aspirin and fibrinolytic therapy for myocardial infarction with ST-segment elevation. N Engl J Med 2005; 352:1179-1189.

Sacks FM, Svetky LP, Vollmer WM et al. Effects on blood pressure of reduced dietary sodium and the dietary approaches to stop hypertension (DASH) diet. N Eng J Med 2001 344:3-10.

Scanlon PJ, Faxon DP, Audet A, et al. . ACC/AHA guidelines for coronary angiography: executive summary and recommendations. Circulation 1999; 99: 2345-2357.

Scott LB, Allen JK. Providers' perceptions of factors affecting women's referral to outpatient cardiac rehabilitation programs: an exploratory study. J Cardiopulm Rehab 2004; 24:387-391.

The Seventh Report of the Joint National Committee on Prevention, Evaluation, and Treatment of High Blood Pressure. (JNC 7). U.S. Department of Health and Human Services. National Institutes of Health. National Heart, Lung, and Blood Institute, NIH Publication No. 04-5230, 2004.

Stampfer, MJ, Hu FB, Manson JE, et al. Primary prevention of coronary heart disease in women through diet and lifestyle. N Engl J Med 2000; 343:16-22.

Statement from Elizabeth G. Nabel, M.D., Director of the National Heart, Lung, and Blood Institute of the National Institutes of Health on the Findings of the Women's Health Study, http://www.nhlbi.nih.gov/new/press/05-03-07.htm, accessed April 19, 2005.

Stromberg A, Martensson J. Gender differences in patients with heart failure. Eur J Cardiovasc Nurs, 2003; 2(1):7-18.

Third Report of the National Cholesterol Education Program (NCEP) Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) (ATPIII), NIH, NHLBI, 2002.

Travin MI, Katz MS, Moulton AW, et al. Accuracy of dipyridamole SPECT imaging in identifying individual coronary stenosis and multiivessel disease in women versus men. J Nuc Cardiol 2000; 7:213-220.

U.S. Department of Health and Human Services. Physical activity and health: a Report of the Surgeon General. Atlanta, Georgia: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, 1996.

Vaccarino V, Abramson JL, Veledar E, et al. Sex differences in hospital mortality after coronary artery bypass surgery: evidence for a higher mortality in younger women. Circulation 2002; 105: 1176-1181.

Vaccarino V, Parsons L, Every NR, et al. Sex-based differences in early mortality after myocardial infarction. National registry of myocardial infarction 2 participants. N Engl J Med 1999; 341: 217-225.

Vaccarino V, Zhen QL, Stanislav VK, et al. Sex differences in health status after coronary artery bypass surgery. Circulation 2003; 108:2642-2647.

Wassertheil-Smoller S, Shumaker S, Ockene J, et al. Depression and cardiovascular sequelae in postmenopausal women. The Women's Health Inititiative (WHI). Arch Intern Med. 2004; 164:289-98.

Welch GN and Loscalzo J. Homocysteine and atherothrombosis. N Engl J Med 1998; 338:1042-1050.

Wenger NK. Cardiovascular health and disease in women problems and prospects. Circulation 2004; 109:558-560.

Whitlock EP, Williams SB. The primary prevention of heart disease in women through health behavior change promotion in primary care. Women's Health Issues 2003; 13:122-141.

Yusuf S, Shao F, Mehta S, et al. Effects of clopidogrel in addition to aspirin in patients with acute coronary syndromes without ST-segment elevation. N Engl J Med 2001; 345:494-502.

Zimmerman GL, Olsen CG, Bosworth MF. A 'stages of change' approach to helping patients change behavior. Am Fam Physician 2000; 61:1409-1416.