

CONTENTS

Abstract.....	1
Introduction	2
Purpose and Scope.....	2
Health and Regulatory Concerns Related to Arsenic, Iron, and Manganese in Ground Water.....	4
Previous Investigations.....	5
Acknowledgments	6
Description of the Study Area	7
Stratified-Drift Aquifers	7
Bedrock Aquifers.....	7
Water Use	9
Study Design and Methods.....	10
Lithogeochemical Reclassification of Bedrock Units and Regional-Scale Approach	10
Sources of Ground-Water Data and Well Selection	11
Statistical Analyses.....	14
Relation of Arsenic, Iron, and Manganese in Ground Water to Aquifer Type, Lithogeochemistry, and Land Use.....	17
Occurrence of Arsenic, Iron, and Manganese by Aquifer Type	18
Occurrence of Arsenic, Iron, and Manganese by Lithogeochemical Group	18
Anomalous Concentrations of Arsenic by Bedrock Geologic Unit	25
Relation of Arsenic in Ground Water to Land Use and Lithology	26
Possible Sources of, and Controls on, Arsenic in Ground Water	29
Arsenic in Water from Public-Supply Wells and Future Drinking-Water Standards	31
Summary and Conclusions	32
References Cited.....	33
Appendix 1: Geologic, Land-Use, and Chemical Data for Bedrock Wells in the New England Coastal Basins Study Unit.....	38
Appendix 2: Chemical Data for Surficial Wells in the New England Coastal Basins Study Unit.....	60

FIGURES

1-4. Maps showing:	
1. Location of the New England Coastal Basins study unit.....	3
2. Location of stratified-drift deposits in the New England Coastal Basins study unit.....	8
3. Areal distribution and description of major lithogeochemical groups of bedrock geologic units.....	12
4a. Areal distribution of arsenic concentrations in water from selected bedrock wells	15
4b. Areal distribution of arsenic concentrations in water from selected stratified-drift wells.....	16
5-10. Graphs showing:	
5. Percent detection of arsenic, iron, and manganese concentrations in ground water by aquifer type	18
6. Percent detection of arsenic concentrations in ground water, at or above 0.005 milligrams per liter, by major lithogeochemical group	19
7. Percent detection of arsenic concentrations in ground water, at or above 0.005 milligrams per liter, in selected bedrock geologic units in lithogeochemical group M_c	22
8. Boxplots of iron and manganese concentrations in ground water by major lithogeochemical group.....	23
9. Correlation of arsenic and iron, arsenic and manganese, and iron and manganese concentrations in ground water by major lithogeochemical group.....	24
10. Percent detection of arsenic concentrations, at or above 0.005 milligrams per liter, in ground water, by selected bedrock geologic units and their associated lithogeochemical groups	25

11. Map showing areal distribution of selected bedrock geologic units, and their associated lithogeochemical groups that have anomalous arsenic concentrations in ground water.....	27
12. Map showing areal distribution of major land-use categories.....	28
13. Graph showing percent detection of arsenic concentrations in ground water, at or above 0.005 milligrams per liter, by major land-use category	29
14. Graph showing percent detection of arsenic concentrations in ground water by lithogeochemical group and major land-use category; (a) major lithogeochemical group with and without wells in the agricultural category, and (b) major land-use category with and without wells in the M _c lithogeochemical group	30

TABLES

1. Summary of estimated water use for public water supplies by aquifer type in the New England Coastal Basins study unit	9
2. Summary of public water supplies compiled by State and aquifer type.....	14
3. Percent of wells in stratified-drift and bedrock aquifers yielding water with detectable concentrations of arsenic, iron, and manganese, and Chi-square statistics	18
4. Summary statistics for concentrations of arsenic, iron, and manganese in ground water by major lithogeochemical group.....	20
5. Summary of attained significance levels (p-values) for Kruskal-Wallis tests of the concentrations of water-quality variables compared by lithogeochemical and land use variables	21
6. Summary of Spearman's rho rank-correlation coefficients for concentrations of arsenic and iron, arsenic and manganese, and iron and manganese	24
7. Percent of wells yielding water with arsenic concentrations exceeding 0.005, 0.01, and 0.02 milligrams per liter by lithogeochemical group	31

CONVERSION FACTORS

	Multiply	By	To obtain
	foot (ft)	0.3048	meter
	square mile (mi ²)	2.590	square kilometer
	million gallons per day (Mgal/d)	0.04381	cubic meter per second
	feet per day (ft)	0.3048	meters per day
	pounds per acre (lbs/acre)	1.12	kilograms per hectare
	pounds per acre per year (lbs/acre/yr)	1.12	kilograms per hectare per year
	gallons per day per person (gal/d/person)	0.06309	liters per second per person
	gallons per minute (gal/min)	3.785	liters per minute

OTHER ABBREVIATIONS USED IN THIS REPORT

milligrams per liter	(mg/L)
parts per million	(ppm)