

New Episodes of Volcanism at Kilauea Volcano, Hawaii

Mid-2007 was a time of intense activity at Kilauea Volcano, Hawaii (see Figure 1). In June, the long-lived Pu'u 'Ō'ō-Kupaianaha eruption, a dual-vent system along the east rift zone (ERZ) that has been erupting since 1983 [Heliker *et al.*, 2003], paused due to the outbreak of a new vent farther up the rift (see Figure 2). The Pu'u 'Ō'ō vent collapsed following that activity, and the resulting reorganization of the magma plumbing system led to the formation of a second new eruptive vent 2 kilometers downrift of Pu'u 'Ō'ō.

These events were well documented by geological, geophysical, and geochemical monitoring. This article summarizes results from these monitoring efforts and interprets the changes that occurred at Kilauea during June–October 2007.

Before 17 June 2007: Setting the Stage

Kilauea's summit deflated throughout most of the sustained Pu'u 'Ō'ō-Kupaianaha eruption. In late 2003, however, the summit started to inflate, accumulating 0.55 meters of extension and 0.25 meters of uplift by mid-2007. Sulfur dioxide (SO₂) emissions from the ERZ, which indicate the volume of magma degassing at Pu'u 'Ō'ō, doubled in early 2005 and remained elevated through mid-2007. Both the summit inflation and the doubling of SO₂ emissions suggest increased magma supply to the shallow volcanic system.

From 17 to 30 June: ERZ Activity and Aftermath

A new episode of activity began at 0216 local time (LT) on 17 June 2007, when summit tiltmeters on Kilauea began recording rapid deflation. At the same time, increased earthquake activity and ground tilt were

By M. POLAND, A. MIKLIUS, T. ORR, J. SUTTON, C. THORNER, AND D. WILSON

measured along the ERZ between Pauahi crater and Mauna Ulu (Figure 3b, time T₁), suggesting the intrusion of magma in that region. Deflationary tilt began at Pu'u 'Ō'ō a few minutes later, indicating that magma supply there had been disrupted. Because of this deflation, Pu'u 'Ō'ō began to collapse on the morning of 17 June, and within a few days parts of the crater floor had dropped as much as 80 meters.

Seismic and geodetic data suggest that over approximately the next 48 hours there were three additional pulses of intrusion to the ERZ (Figure 3b, starting at times T₂, T₃, and T₄). Between 18 and 19 June, a small eruption (area, 2211 square meters; volume, about 1500 cubic meters) occurred north of Makaopuhi crater (see Figure 2). The chemistry and temperature of the erupted lava (Figure 3a) suggest that the magma that fed the eruption came directly from Kilauea's summit without incorporating magma stored in the rift zone. At 1030 LT on 19 June (about 57 hours after the onset of intrusion), summit tilt switched from deflation to inflation, and deformation and swarm seismicity in the ERZ were minimal.

Between 20 June and 1 July, active lava was not visible anywhere on the surface of Kilauea Volcano. The combined SO₂ emission rate from Pu'u 'Ō'ō and the 18–19 June eruption site declined through this time period to less than 200 tons per day (Figure 3a), and deflationary tilt at Pu'u 'Ō'ō continued until 27 June.

From 1 to 20 July: Lava Returns to Pu'u 'Ō'ō

Eruptive activity resumed at Pu'u 'Ō'ō on 1 July, and lava had filled the crater to within 30 meters of the rim by 12 July. On 13 July, new vents appeared high on the crater walls, well above the lava lake, and Webcam images recorded piston-like uplift of about 1 meter per day along the interior crater walls. Inflation of Pu'u 'Ō'ō was docu-

mented by ground tilt and Global Positioning System (GPS; see Figures 3a and 3c).

SO₂ emission rates during 1–20 July remained low. The temperature and chemistry of lava lake samples (Figure 3a) are indicative of shallow magma storage and crystallization prior to eruption, as opposed to direct transport from Kilauea's summit as is inferred for the 18–19 June eruption near Makaopuhi crater.

From 21 July to 31 October: Downrift Migration of the Eruption

Just before midnight on 20 July, tilt and GPS measurements at Pu'u 'Ō'ō indicated cone deflation as well as a deformation source to the east (Figure 3c). Webcam imagery revealed draining of the Pu'u 'Ō'ō lava lake, followed by glow that indicated the start of a fissure eruption on the east flank of Pu'u 'Ō'ō a few minutes after midnight. The fissure system consisted of four segments (designated A–D) that extended 2 kilometers east from near the east rim of Pu'u 'Ō'ō crater (Figure 3c). No unusual seismicity accompanied the new outbreak, and deformation associated with fissure opening, as seen from interferometric synthetic aperture radar (InSAR) and GPS data, is highly localized (Figure 3c), suggesting a shallow source.

Immediately following fissure opening, SO₂ emissions from the ERZ increased by an order of magnitude and, following a brief decline, rose to more than 2000 tons each day by the end of August. Lava that erupted from the fissure system was initially similar to that from the Pu'u 'Ō'ō lava lake, although a slight but steady increase in magnesium oxide content through the end of October (Figure 3a) may reflect an increasing component of Kilauea summit-derived magma mixed with rift-stored, crystallizing magma.

The erupted lava initially built a series of ponds between Pu'u 'Ō'ō and Kupaianaha that were perched as much as 15 meters above surrounding terrain. By 10 August, all eruptive activity focused on fissure D, which formed an open lava channel (Figure 1) and fed a series of northeast directed 'a'ā flows (rough-textured blocky lava) that extended




Fig. 1. View to the south of the perched lava channel emanating from fissure D, with Pu'u 'Ō'ō in the background. USGS photograph by Jim Kauahikaua taken on 7 October 2007.

a maximum distance of 6 kilometers from the vent by mid-August. Subsequent lava flows consisted of channel overflows, breakouts from the base of the channel walls, and short-lived flows fed through breaches at the end of the lava channel.

Characteristics of Intrusive and Extrusive Activity During Mid-2007 at Kilauea

ERZ intrusions in 1997 and 1999 grew in single pulses and were accompanied by relatively little seismicity, suggesting passive magma emplacement facilitated by seaward motion of Kilauea's south flank [Cervelli *et al.*, 2002; Owen *et al.*, 2000]. In contrast, the June 2007 intrusion had several distinct pulses, released relatively more seismic energy, and erupted lavas characteristic of summit-derived magma (the eruption that resulted from the 1997 intrusion was clearly derived from rift-stored magma [Thornber *et al.*, 2003]), all of which suggest forcible intrusion in June 2007. The difference in styles between the 2007 activity and past intrusions (both eruptive and noneruptive) is probably related to the inflated state of the summit magma system in 2007.

Interferograms formed from radar images acquired by the European Space Agency's Environmental Satellite (Envisat) that span mid-2007 indicate subsidence of Kilauea's summit and a butterfly-shaped pattern of uplift centered on the ERZ that is indicative of dike intrusion (Figure 2). A simple model of the deformation using a point source of volume loss at Kilauea's summit and an opening dislocation source between Pauahi and Makaopuhi craters suggests 1–2 million cubic meters of volume loss from a shallow magma reservoir near Halema'uma'u crater during 17–19 June and an ERZ intrusion volume of about 15 million cubic meters. The discrepancy in volumes introduces the possibility that magma from other sources (for example, Pu'u 'Ō'ō, where the volume of collapse was of the order of 5 million cubic meters) contributed to the ERZ intrusion.

When the magma supply to Pu'u 'Ō'ō was cut off on 17 June, the resulting collapse disrupted the local magma plumbing system. As the connection between Kilauea's summit and Pu'u 'Ō'ō was reestablished, high rates of uplift and the formation of vents on the crater walls indicated increasing pressure




Fig. 2. Envisat interferogram spanning 12 April to 21 June 2007. Craters and other geographic features are outlined by white solid lines. Dashed white line indicates the ERZ axis. Red dot shows the location of the 18–19 June eruption. Dotted white line shows the extent of the 21 July to 30 September lava flow. Tilt and GPS stations discussed in the text are labeled as white dots and white triangles, respectively. Earthquakes are noted by white stars. White box shows the extent of the Figure 3c map.

Kilauea

cont. from page 37

beneath the Pu'u 'Ō'ō cone. The 21 July fissure eruption was a response to the pressure increase. Formation of a highly efficient vent allowed increased magma flow from the summit to the eruption site and resulted in summit deflation (Figure 3a). Field measurements and SO₂ emissions suggest an effusion rate from the 21 July vent of 0.5–0.8 million cubic meters each day through the end of October, compared with about 0.4 million cubic meters each day prior to 17 June.

In addition to changing the character of Kilauea's eruptive activity, the June intrusion and eruption caused local widening of the ERZ, which increased compressive stress on the volcano's south flank. Heightened seismicity during the second half of 2007 and an unusually strong series of earthquakes between Pu'u 'Ō'ō and the coast in mid-August (Figure 2) were probably a result of this stress.

Kilauea's Current Status

Starting on 21 July 2007, lava erupting from fissure D built and fed a 1.4-kilometer-long, northeast-directed open channel (Figure 1). Frequent overflows from the edges of the channel built levees that elevated the lava stream up to 45 meters above the pre-21 July ground level. From 21 November 2007 to early January 2008, however, the channel has been mostly abandoned, with lava building a series of shields to the southeast of the fissure. About three weeks ago, one of the shields sent a lava flow to within a few hundred meters of the Royal Gardens subdivision. Although the subdivision was mostly deserted after lava inundated and isolated the area in the 1980s and 1990s, there are still homes standing and one permanent resident. The lava flow stagnated before reaching the subdivision, but the current shield-building activity continues to threaten the region.

Kilauea's summit has continued to deflate slowly since 21 July. Although gas emissions and observational evidence suggest a sub-

tle decrease in effusion rate since the end of October, the ongoing extrusion from fissure D and the persistence of summit deflation imply a strong connection between the summit magma system and vent, so we expect erupted lavas to show an increasing component of summit-derived magma over time. The new eruption site has replaced Pu'u 'Ō'ō as the locus of lava effusion, and is likely to remain active for months to years.

The lava poses no immediate threat to communities on the island of Hawaii (besides the mostly empty Royal Gardens subdivision), although the situation may change if the character of the eruption allows for long lava flows fed by tubes and directed at populated areas [Kauahikaua, 2007]. Updates, maps, live Webcam views, and images of the most recent activity are available at <http://volcano.wr.usgs.gov/hvostatus.php>.

Acknowledgments

The response to the mid-2007 activity of Kilauea Volcano was a team effort and would not have been successful without the hard work of colleagues at the U.S. Geological Survey's Hawaiian Volcano Observatory. We acknowledge collaborations with Stanford University and the Pacific GPS Facility at the University of Hawai'i on GPS results. Kristine Larson (University of Colorado, Boulder) assisted with kinematic GPS processing. The European Space Agency provided rapid access to Envisat radar data through Cat-1 project 2765. The Pu'u 'Ō'ō GPS network was supported by a NASA New Investigator Program grant and aided by equipment loans from Ben Brooks (University of Hawai'i, Manoa). Dan Dzurisin, John Power, and Gordon Keating provided constructive and timely reviews.

References

- Cervelli, P., P. Segall, F. Amelung, H. Garbeil, C. Merteens, S. Owen, A. Miklius, and M. Lisowski (2002), The 12 September 1999 Upper East Rift Zone dike intrusion at Kilauea Volcano, Hawaii, *J. Geophys. Res.*, **107**(B7), 2150, doi:10.1029/2001JB000602.
- Heliker, C., J. Kauahikaua, D. R. Sherrod, M. Lisowski, and P. Cervelli (2003), The rise and fall of Pu'u 'Ō'ō cone, 1983–2002, in *The Pu'u 'Ō'ō-Kupaianaha Eruption of Kilauea Volcano, Hawaii: The First 20 Years*, edited by C. Heliker et al., *U.S. Geol. Surv. Prof. Pap.*, 1676, 29–51.
- Kauahikaua, J. P. (2007), Lava flow hazard assessment, as of August 2007, for Kilauea East Rift Zone Eruptions, Hawai'i Island, *U.S. Geol. Surv., Open File Rep.*, 2007-1264, 9 pp.
- Owen, S., P. Segall, M. Lisowski, A. Miklius, M. Murray, M. Bevis, and J. Foster (2000), January 30, 1997, eruptive event on Kilauea Volcano, Hawaii, as




Fig. 3. Time series showing the chronology of the mid-2007 events at Kilauea Volcano. GPS and tilt station locations are shown in Figures 2 and 3c, with the same labeling conventions. (a) Daily GPS solutions, SO₂, and petrology data from 1 January to 31 October, 2007. Distance change between AHUP and UWEV is a proxy for summit inflation/deflation (positive/negative). Note dramatic changes in established trends of deformation, lava chemistry and temperature, and gas emissions during and following both the mid-June and mid-July eruptive events (noted by grey bars). (b) Tilt (top, left axis), 4-minute GPS solutions (top, right axis), and seismicity (bottom); circle size and scale in upper rift indicate earthquake magnitude, and color indicates timing) during June 2007 ERZ intrusion and eruption. Times are Hawaii Standard Time (UTC minus 10 hours). Components of tilt are in degrees, with the positive tilt toward that azimuth. The map coverage and symbols are as in Figure 2. Note the downrift progression of seismicity and correlation with changes in deformation over time, both indicating the propagation of a dike from Kilauea's summit to east of Makaopuhi crater during 17–19 June. (c) Tilt and 4-minute GPS solutions (top) from Pu'u 'Ō'ō during a 24-hour period spanning 20–21 July. Map (bottom) shows the interferogram from Figure 2 with 10-meter topographic contours, GPS and tilt station locations (symbols are as in Figure 2), GPS horizontal displacements (white arrows), fissure segments (red lines), and 21 July to 31 October lava flows (red). Note the high rate and magnitude of the deformation that preceded and accompanied the 21 July fissure eruption.

monitored by continuous GPS, *Geophys. Res. Lett.*, **27**(17), 2757–2760.

Thornber, C. R., C. Heliker, D. R. Sherrod, J. P. Kauahikaua, A. Miklius, P. G. Okubo, F. A. Truedell, J. R. Budahn, I. W. Ridley, and G. P. Meeker, (2003), Kilauea East Rift Zone magmatism: An episode 54 perspective, *J. Petrol.*, **44**(9), 1525–1559.

Author Information

Michael Poland, U.S. Geological Survey (USGS) Hawaiian Volcano Observatory (HVO), Hawaii National Park, Hawaii; E-mail: mpoland@usgs.gov; Asta Miklius, Tim Orr, and Jeff Sutton, USGS HVO; Carl Thornber, USGS Cascades Volcano Observatory, Vancouver, Wash.; and David Wilson, USGS HVO.

EOS

TRANSACTIONS
AMERICAN GEOPHYSICAL UNION
The Newspaper of the Earth and Space Sciences

Editors

John W. Geissman: Department of Earth and Planetary Science, University of New Mexico, Albuquerque, USA; jgeiss@unm.edu

Wendy S. Gordon: Texas Commission on Environmental Quality, wsgordon@tceq.state.tx.us