

REFERENCE COPY

Do Not Remove from the Library

U. S. Fish and Wildlife Service


Biological Report 82(11.44)
April 1986

National Wetlands Research Center
700 Cajun Dome Boulevard
Lafayette, Louisiana 70506

TR EL-82-4

**Species Profiles: Life Histories and
Environmental Requirements of Coastal Fishes
and Invertebrates (Pacific Southwest)**

CALIFORNIA HALIBUT



Fish and Wildlife Service

U.S. Department of the Interior

**Coastal Ecology Group
Waterways Experiment Station**

U.S. Army Corps of Engineers

Biological Report 82(11.44)
TR EL-82-4
April 1986

Species Profiles: Life Histories and Environmental Requirements
of Coastal Fishes and Invertebrates (Pacific Southwest)

CALIFORNIA HALIBUT

by

Stephen T. Kucas and Thomas J. Hassler
California Cooperative Fishery Research Unit
Humboldt State University
Arcata, CA 95521

Project Officer
John Parsons
National Coastal Ecosystems Team
U.S. Fish and Wildlife Service
1010 Gause Boulevard
Slide11, LA 70458

Performed For
Coastal Ecology Group
Waterways Experiment Station
U.S. Army Corps of Engineers
Vicksburg, MS 39180

and

National Coastal Ecosystems Team
Division of Biological Services
Research and Development
Fish and Wildlife Service
U.S. Department of the Interior
Washington, DC 20240

This series should be referenced as follows:

U.S. Fish and Wildlife Service. 1983-19___. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates. U.S. Fish Wildl. Serv. Biol. Rep. 82(11). U.S. Army Corps of Engineers, TR EL-82-4.

This profile should be cited as follows:

Kucas, S.T., and T.J. Hassler. 1986. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Southwest)--California halibut. U.S. Fish Wildl. Serv. Biol. Rep. 82(11.44). U.S. Army Corps of Engineers, TR EL-82-4. 8 pp.

PREFACE

This species profile is one of a series on coastal aquatic organisms, principally fish, of sport, commercial, or ecological importance. The profiles are designed to provide coastal managers, engineers, and biologists with a brief comprehensive sketch of the biological characteristics and environmental requirements of the species and to describe how populations of the species may be expected to react to environmental changes caused by coastal development. Each profile has sections on taxonomy, life history, ecological role, environmental requirements, and economic importance, if applicable. A three-ring binder is used for this series so that new profiles can be added as they are prepared. This project is jointly planned and financed by the U.S. Army Corps of Engineers and the U.S. Fish and Wildlife Service.

Suggestions or questions regarding this report should be directed to one of the following addresses.

Information Transfer Specialist
National Coastal Ecosystems Team
U.S. Fish and Wildlife Service
NASA-Slidell Computer Complex
1010 Gause Boulevard
Slidell, LA 70458

or

U.S. Army Engineer Waterways Experiment Station
Attention: WESER-C
Post Office Box 631
Vicksburg, MS 39180

CONVERSION TABLE

Metric to U.S. Customary

| <u>Multiply</u> | <u>By</u> | <u>To Obtain</u> |
|--------------------------------------|--------------|-----------------------|
| millimeters (mm) | 0.03937 | inches |
| centimeters (cm) | 0.3937 | inches |
| meters (m) | 3.281 | feet |
| kilometers (km) | 0.6214 | miles |
| square meters (m ²) | 10.76 | square feet |
| square kilometers (km ²) | 0.3861 | square miles |
| hectares (ha) | 2.471 | acres |
| liters (l) | 0.2642 | gallons |
| cubic meters (m ³) | 35.31 | cubic feet |
| cubic meters | 0.0008110 | acre-feet |
| milligrams (mg) | 0.00003527 | ounces |
| grams (g) | 0.03527 | ounces |
| kilograms (kg) | 2.205 | pounds |
| metric tons (t) | 2205.0 | pounds |
| metric tons | 1.102 | short tons |
| kilocalories (kcal) | 3.968 | British thermal units |
| Celsius degrees | 1.8(°C) + 32 | Fahrenheit degrees |

U.S. Customary to Metric

| | | |
|---------------------------------|-----------------|-------------------|
| inches | 25.40 | millimeters |
| inches | 2.54 | centimeters |
| feet (ft) | 0.3048 | meters |
| fathoms | 1.829 | meters |
| miles (mi) | 1.609 | kilometers |
| nautical miles (nmi) | 1.852 | kilometers |
| square feet (ft ²) | 0.0929 | square meters |
| acres | 0.4047 | hectares |
| square miles (mi ²) | 2.590 | square kilometers |
| gallons (gal) | 3.785 | liters |
| cubic feet (ft ³) | 0.02831 | cubic meters |
| acre-feet | 1233.0 | cubic meters |
| ounces (oz) | 28.35 | grams |
| pounds (lb) | 0.4536 | kilograms |
| short tons (ton) | 0.9072 | metric tons |
| British thermal units (Btu) | 0.2520 | kilocalories |
| Fahrenheit degrees | 0.5556(°F - 32) | Celsius degrees |

CONTENTS

| | <u>Page</u> |
|-------------------------------------|-------------|
| PREFACE..... | iii |
| CONVERSION TABLE..... | iv |
| ACKNOWLEDGMENTS..... | vi |
| NOMENCLATURE/TAXONOMY/RANGE..... | 1 |
| MORPHOLOGY/IDENTIFICATION AIDS..... | 1 |
| REASON FOR INCLUSION IN SERIES..... | 1 |
| LIFE HISTORY..... | 3 |
| Spawning..... | 3 |
| Larval Stage..... | 3 |
| Juveniles..... | 3 |
| Maturity and Life Span..... | 3 |
| GROWTH CHARACTERISTICS..... | 3 |
| COMMERCIAL AND SPORT FISHERIES..... | 4 |
| ECOLOGICAL ROLE..... | 4 |
| ENVIRONMENTAL REQUIREMENTS..... | 5 |
| Temperature..... | 5 |
| Depth..... | 5 |
| Substrate..... | 5 |
| Other Environmental Factors..... | 5 |
| LITERATURE CITED..... | 7 |

ACKNOWLEDGMENTS

We are grateful for the reviews by Peter L. Haaker and Roger N. Lea, California Department of Fish and Game. David Moran served as assistant project officer at NCET.




Figure 1. California halibut.

CALIFORNIA HALIBUT

NOMENCLATURE/TAXONOMY/RANGE

Scientific name Paralichthys californicus (Ayres)
 Preferred common name California halibut (Figure 1)
 Class Osteichthyes
 Order Pleuronectiformes
 Family..... Bothidae

Geographic range: Gulf of California (isolated population), and from Magdalena Bay, California, to Quillayute River, Washington (Miller and Lea 1972; Figure 2). Most occur below San Francisco Bay (Haaker 1975).

MORPHOLOGY/IDENTIFICATION AIDS¹

Fin rays -- dorsal 66-76, anal 49-59, pectoral 10-13; lateral line

¹Largely extracted from Miller and Lea (1972).

scales about 100; gill rakers 7-11 + 18-23; vertebrae 34-36; eyes sinistral or dextral. Body elliptical, slender; head small; mouth large, maxillary extending to or beyond rear edge of lower eye; teeth strong and sharp on both sides of head; eyes small; interorbital space wide, flat; lateral line with high arch over pectoral fin; scales small, smooth; caudal fin indented.

Color in life: eyed side grayish or greenish-brown, sometimes mottled with lighter and darker shades; small white spots; and blind side white to cream. Young fish with bluish-white spots (Bane and Bane 1971); partial pigmentation has been reported on the eyed side (Haaker and Lane 1973).

REASON FOR INCLUSION IN SERIES

The California halibut commonly lives in bays and estuaries along the




Figure 2. Coastal distribution of the California halibut.

central and southern California coast (Haaker 1975), and supports a small but valuable sport and commercial fishery. A sharp reduction in California halibut landings from California in the late 1920's and declining landings in recent years have prompted scientists to determine the cause. To date little is known about its biology and life history.

LIFE HISTORY

Spawning

The major spawning areas for the California halibut are unknown. In spring, halibut emigrate from relatively deep offshore waters to shallow coastal waters (Clark 1930a, 1930b) to spawn at depths of 5 to 18 m (Young 1960). California halibut spawn from February through July, peaking in May (Ginsberg 1952). The eggs are demersal. After spawning the adults return to water about 40 to 100 m deep (Ginsberg 1952).

Larval Stage

The larval and postlarval stages of the California halibut are pelagic for several months before the postlarvae settle to the bottom (Winzler and Kelly Consulting Engineers 1977). In southern California, the planktonic larval stages, about 10 mm standard length (SL), are pelagic and live in water 12 to 45 m deep and 1.9 to 5.4 km out from shore. Larger and older larvae live closer to shore, and most are suspected to live in embayments (Plummer et al. 1983).

Juveniles

Juveniles begin to settle and assume bottom dwelling habits in June (Frey 1971); they remain in shallow water and do not move extensively (Frey 1971; Haaker 1975). Juvenile halibut begin to emigrate to deeper

water when they are about 200 mm total length (TL) (Frey 1971).

Maturity and Life Span

Most halibut females are sexually mature at 430 mm TL or at the end of their third year of life (Frey 1971). Most males mature when about 230 mm TL long or at the end of their second year of life.

Halibut are generally long-lived. Both males and females may live as long as 30 years, though generally the males do not grow as large or live as long as the females (Frey 1971). Halibut are about 1,080 mm TL long when they are 12 years old.

GROWTH CHARACTERISTICS

The age and rate of growth of halibut from California commercial landings were reported by Frey (1971). Table 1 summarizes this information.

The growth rate of female halibut in their first and second years of life (Haaker 1975) was slower

Table 1. Age and growth (mid-year lengths) of female halibut from California commercial landings (from Frey 1971).

| Year | Mid-year length (mm TL) | Weight (kg) |
|------|----------------------------|----------------|
| 1 | 178 | 0.057 |
| 2 | 318 | 0.340 |
| 3 | 439 | 0.907 |
| 4 | 553 | 1.758 |
| 5 | 648 | 2.835 |
| 7 | 813 | 5.783 |
| 9 | 940 | 9.072 |
| 12 | 1054 | 10.745 |

in Anaheim Bay, California, than that reported by Frey (1971) from California commercial landings. In their third year, growth of females from Anaheim Bay was greater than that reported from California landings. Data were not available for other age groups.

COMMERCIAL AND SPORT FISHERIES

The California halibut supports a small but lucrative sport and commercial fishery. The demand for the species is high, but its abundance and availability are relatively low.

Records of the commercial catches of California halibut from 1916 to 1947 were summarized by Holmberg (1949). Frey (1971) summarized the California landings data to 1969. Commercial landings data of California halibut from 1939 to 1984 are presented in Figure 3.




Figure 3. Commercial landings of California halibut from 1939 to 1984 (data from U.S. National Marine Fisheries Service 1942-1984 and California Department of Fish and Game preliminary monthly landings data 1978-1984).

The commercial catch of the California halibut for California, Washington, Oregon, and Mexican waters declined from 2,131 t in 1919 to 330 t in 1941 (Holmberg 1949). From 1941 to 1969 the average annual catch was 419 t. Large catches in 1946 and 1947, 1,134 t and 811 t respectively, may be attributed to reduced fishing during World War II, which allowed the halibut stocks to build up (Frey 1971). After 10 years (1947-1956) of unusually cold water along the California coast, the commercial catch increased from 161 t in 1959 to 578 t in 1964. The catch then declined to 117 t in 1970, the lowest catch on record.

From 1971 to 1983 the catch of California halibut averaged 271 t (range: 123 t to 563 t). The value of the commercial landings of California halibut in 1977 was estimated at \$460,000 (U.S. National Marine Fisheries Service 1942-1984). More recent estimates have not been published.

ECOLOGICAL ROLE

The larvae and young juveniles of the California halibut are planktivorous, but as they grow to adulthood they become piscivorous (Haaker 1975; Plummer et al. 1983). Halibut 230 mm TL long or longer feed heavily on fish (primarily during daylight) and, to a lesser extent, on shrimp (Haaker 1975).

Juvenile and adult California halibut are especially adapted for catching and digesting fish because they are equipped with long heavily toothed gill rakers that aid in holding and swallowing, an intestine with a simple loop, and a brain with large optic lobes (Haaker 1975). This raptorial predator has sharp canine-like teeth and a large mouth.

Haaker (1975) found that the stomachs of halibut 12 to 510 mm long contained 65% fish, 34% crustaceans,

and 1% mollusks, though the diet of these fish changed as the fish grew. For example, halibut less than 55 mm long ate mostly small fish (gobies) and small crustaceans such as amphipods, copepods and mysids. Fish from 55 to 230 m ate larger crustaceans and fishes. Larger halibut ate larger prey (Plummer et al. 1983). The northern anchovy, Engraulis mordax, is probably the most important prey of the California halibut (Frey 1971).

ENVIRONMENTAL REQUIREMENTS

Temperature

Laboratory behavioral studies indicated that California halibut in age groups 0 and I are eurythermal, whereas individuals in age group II and older tended to be stenothermal (Innis 1980). Adult halibut placed in a water temperature gradient chamber preferred water temperatures between 15 °C and 23 °C (Innis 1980).

Depth

California halibut about 100 mm long (SL) were largely absent from nearshore coastal waters 6 to 30 m deep off northern San Diego County and

occurred primarily in embayments (Plummer et al. 1983). Age I+ fish (15-25 cm SL) and II+ (30-45 cm SL) and older fish (Haaker 1975; Innis 1980) segregate by depth, with older juveniles and young adults occurring deeper than younger juveniles. Adults prefer deeper water (Haaker 1975; Innis 1980). The California halibut inhabits offshore waters as deep as 100 m (Miller and Lea 1972), but most are caught commercially on the bottom at water depths of 6 to 40 m (Ginsberg 1952).

Substrate

Adult California halibut occur over sandy bottoms from the surf zone to 100 m (Jones and Stokes, Inc. 1981).

Other Environmental Factors

Shallow bays and estuaries may be critical habitat for the survival and growth of larvae and young juveniles, and the shallow waters of the open coast almost surely serve as nursery grounds for California halibut (Plummer et al. 1983; Haaker 1975). Consequently, major alterations of bays and estuaries along the coast of California could be harmful to halibut habitat and reduce the abundance of halibut.

LITERATURE CITED


- Bane, G. W. and A.W. Bane. 1971. Bay fishes of Northern California. Mariscos Publication, Southampton, N.Y. 143 pp.
- Clark, G.H. 1930a. The California halibut (Paralichthys californicus) and an analysis of the boat catches. Calif. Dep. Fish Game Fish Bull. 32:1-52.
- Clark, G.H. 1930b. California halibut. Calif. Fish Game 16:315-317.
- Frey, H.W., ed. 1971. California's living marine resources and their utilization. California Department Fish Game, Sacramento. 148 pp.
- Ginsberg, I. 1952. Flounders of the genus Paralichthys and related genera in American waters. U.S. Fish Wildl. Serv. Fish. Bull. 71:267-351.
- Haaker, P.L. 1975. The biology of the California halibut, Paralichthys californicus (Ayres), in Anaheim Bay, California. Pages 137-151 in E.D. Lane and C.W. Hills, eds. The marine resources of Anaheim Bay. Calif. Dep. Fish Game Fish. Bull. 165.
- Haaker, P.L., and E.D. Lane. 1973. Frequencies of anomalies in a bothid, Paralichthys californicus, and a pleuronectid, Hypsopsetta guttulata, flatfish. Copeia 1973: 22-25.
- Holmberg, E.K. 1949. California halibut: the commercial fish catch of California for the year 1947 with an historical review 1916-1947. Calif. Fish Game Fish. Bull. 74:75-77.
- Innis, D.B. 1980. Growth characteristics and temperature preference behavior of juvenile California halibut, Paralichthys californicus, and their relationship to the effects of thermal effluent. Master's Thesis. San Diego State University, Calif. 110 pp.
- Jones and Stokes, Inc. 1981. An ecological characterization of the central and northern California coastal region. U.S. Fish Wildl. Serv. FWS/OBS-80/47.1, Vol. III Part I.
- Miller, D.J., and R.N. Lea. 1972. Guide to the coastal marine fishes of California. Calif. Dep. Fish Game Fish. Bull. 157. 249 pp.
- Plummer, K.M., E.E. DeMartini, and D.A. Roberts. 1983. The feeding habits and distribution of juvenile-small adult California halibut (Paralichthys californicus) in coastal waters off northern San Diego County, Calif. Coop. Oceanic Fish. Invest. Rep. 24:194-201.
- U.S. National Marine Fisheries Service. 1942-1984. Fishery Statistics of the United States 1939-1977, Statistical Digests Numbers 1 through 71.
- Winzler and Kelly. 1977. A summary of knowledge of the central and northern California coastal zone and

offshore areas. Volume 2 (Chapters 5-6), Biological Conditions. Prepared for U.S. Bureau of Land Management. Winzler and Kelly Consulting Engineers. Eureka, Califor-

nia. 337 pp.

Young, P.H. 1960. California ocean fisheries resources to the year 1960. Calif. Fish Game Comm. 148 pp.

| | | | | |
|--|--|---|--|-------------------------------------|
| REPORT DOCUMENTATION PAGE | | 1. REPORT NO. Biol. Rep. 82(11.44)* | 2. | 3. Recipient's Accession No. |
| 4. Title and Subtitle Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Pacific Southwest)--California Halibut | | | 5. Report Date April 1986 | |
| 7. Author(s) Stephen T. Kucas and Thomas J. Hassler | | | 8. Performing Organization Rept. No. | |
| 9. Performing Organization Name and Address California Cooperative Fishery Research Unit Humboldt State University Arcata, CA 95521 | | | 10. Project/Task/Work Unit No. | |
| 12. Sponsoring Organization Name and Address National Coastal Ecosystems Team Fish and Wildlife Service U.S. Department of the Interior Washington, DC 20240 | | | 11. Contract(C) or Grant(G) No. (C) (G) | |
| U.S. Army Corps of Engineers Waterways Experiment Station P.O. Box 631 Vicksburg, MS 39180 | | | 13. Type of Report & Period Covered | |
| 14. | | | | |
| 15. Supplementary Notes *U.S. Army Corps of Engineers Report No. TR EL-82-4 | | | | |
| 16. Abstract (Limit: 200 words) Species profiles are literature summaries of the taxonomy, morphology, distribution, life history, and environmental requirements of coastal aquatic species. They are prepared to assist in environmental impact assessment. The California halibut, <i>Paralichthys californicus</i> , usually lives nearshore at depths of 100 m or less. It spawns at depths of 5 to 18 m; spawning is from February to July and peaks in May. Some California halibut may live as long as 30 years. Annual commercial catches (1916-1984) for California ranged from 29 to 779 t. Juvenile halibut feed on zooplankton and small fish; adults are almost totally piscivorous. Adults preferred temperatures of 15 to 23 °C. Shallow bays and estuaries are probably important nursery grounds for the California halibut. Information about the environmental requirements of the California halibut is scarce. | | | | |
| 17. Document Analysis a. Descriptors Fishes Life Cycles Growth Fisheries Feeding habits Depth Estuaries | | | | |
| b. Identifiers/Open-Ended Terms California halibut Spawning <u>Paralichthys californicus</u> | | | | |
| c. COSATI Field/Group | | | | |
| 18. Availability Statement Release unlimited | | 19. Security Class (This Report) Unclassified | | 21. No. of Pages 8 |
| | | 20. Security Class (This Page) Unclassified | | 22. Price |



REGION 1

Regional Director
 U.S. Fish and Wildlife Service
 Lloyd Five Hundred Building, Suite 1692
 500 N.E. Multnomah Street
 Portland, Oregon 97232

REGION 2

Regional Director
 U.S. Fish and Wildlife Service
 P.O. Box 1306
 Albuquerque, New Mexico 87103

REGION 3

Regional Director
 U.S. Fish and Wildlife Service
 Federal Building, Fort Snelling
 Twin Cities, Minnesota 55111

REGION 4

Regional Director
 U.S. Fish and Wildlife Service
 Richard B. Russell Building
 75 Spring Street, S.W.
 Atlanta, Georgia 30303

REGION 5

Regional Director
 U.S. Fish and Wildlife Service
 One Gateway Center
 Newton Corner, Massachusetts 02158

REGION 6

Regional Director
 U.S. Fish and Wildlife Service
 P.O. Box 25486
 Denver Federal Center
 Denver, Colorado 80225

REGION 7

Regional Director
 U.S. Fish and Wildlife Service
 1011 E. Tudor Road
 Anchorage, Alaska 99503



DEPARTMENT OF THE INTERIOR

U.S. FISH AND WILDLIFE SERVICE



As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering the wisest use of our land and water resources, protecting our fish and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to assure that their development is in the best interests of all our people. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.