

2. Screening for High Blood Cholesterol and Other Lipid Abnormalities

RECOMMENDATION

Periodic screening for high blood cholesterol is recommended for all men ages 35–65 and women ages 45–65. There is insufficient evidence to recommend for or against routine screening of asymptomatic persons over age 65, but recommendations to screen healthy men and women ages 65–75 may be made on other grounds (see *Clinical Intervention*). There is also insufficient evidence to recommend for or against routine screening in children, adolescents, or young adults. Recommendations for screening adolescents and young adults with risk factors for coronary disease, and against routine screening in children, may be made on other grounds (see *Clinical Intervention*). There is insufficient evidence to recommend for or against routine screening for other lipid abnormalities. All patients should receive periodic screening and counseling regarding other measures to reduce their risk of coronary disease (see Chapter 3, Screening for Hypertension; Chapter 54, Counseling to Prevent Tobacco Use; Chapter 55, Counseling to Promote Physical Activity; and Chapter 56, Counseling to Promote a Healthy Diet).

Burden of Suffering

Elevated blood cholesterol is one of the major modifiable risk factors for coronary heart disease (CHD),¹ the leading cause of death in the U.S. CHD accounts for approximately 490,000 deaths each year,² and angina and nonfatal myocardial infarction (MI) are a source of substantial morbidity. CHD is projected to cost over \$60 billion in 1995 in the U.S. in medical expenses and lost productivity.³ The incidence of CHD is low in men under age 35 and in premenopausal women (1–2/1,000 annually),⁴ but climbs exponentially during middle age for both men and women. The onset of CHD is delayed approximately 10 years in women compared with men, probably due to effects of estrogen,⁵ but women account for 49% of all CHD deaths in the U.S.² Clinical events are the result of a multifactorial process that begins years before the onset of symptoms. Autopsy studies detected early lesions of atherosclerosis in many adolescents and young

adults.^{6–10} The onset of atherosclerosis and symptomatic CHD is earlier among persons with inherited lipid disorders such as familial hypercholesterolemia (FH)¹¹ and familial combined hyperlipidemia (FCH).¹²

Serum Cholesterol and Risk of Coronary Heart Disease. Epidemiologic, pathologic, animal, genetic, and clinical studies support a causal relationship between blood lipids (usually measured as serum levels) and coronary atherosclerosis.^{1,13–15} Extended follow-up of large cohorts (predominantly middle-aged men)^{16–18} provides evidence that CHD risk increases in a continuous and graded fashion, beginning with cholesterol levels as low as 150–180 mg/dL;^a this association extends to cholesterol levels measured as early as age 20 in men.^{14,19} During middle age, for each 1% increase in total cholesterol, CHD risk increases by an estimated 3%.²⁰ High cholesterol (240 mg/dL) is also a risk factor in middle-aged women, but most coronary events in women occur well after menopause.^{5,17,21–24} Some studies report that cholesterol alone is a weak predictor of CHD mortality in the elderly,^{24a,190} but an overview of 24 cohort studies indicates that high cholesterol remains a risk factor for CHD after age 65,²³ with the strongest associations among healthier elderly populations followed over longer periods.^{25–27} The association is weaker in older women than in men²³ and is not consistent for cholesterol levels measured after age 75.^{28–31}

Expert panels have defined high and “borderline high” (200–239 mg/dL) cholesterol to simplify clinical decisions.¹ Because CHD is a multifactorial process, however, there is no definition of high cholesterol that discriminates well between individuals who will or will not develop CHD.^{32,33} Due to nonlipid risk factors, persons with cholesterol below 240 mg/dL account for the majority of all CHD events.^{34,35} Among middle-aged men, 9–12% of those with cholesterol 240 mg/dL or greater will develop symptomatic CHD over the next 7–9 years,^{34,36} but most of them have multiple other risk factors for CHD.³⁵ The excess (i.e., absolute) risk due to high cholesterol (and the probable benefit of lowering cholesterol) increases with the underlying risk of CHD. In a 12-year study of over 316,000 men aged 35–57, the excess CHD mortality attributable to high cholesterol was greatest in men over age 45, and in those who smoked or had hypertension.¹⁶ The increase in CHD mortality associated with a given increment in serum cholesterol was steepest at very high values (>300 mg/dL).¹⁶ Excess risk from high cholesterol is smaller in women, who have less than half the CHD risk as do men at any given cholesterol level.^{17,23,37} Although the relative risk associated with high serum chole-

^aTo convert values for serum total cholesterol, HDL-C, and LDL-C to mmol/L, multiply by 0.02586. Equivalent values for commonly used thresholds are 280 mg/dL = 7.2 mmol/L, 240 mg/dL = 6.2 mmol/L, 200 mg/dL = 5.2 mmol/L.

terol declines with age,^{17,23,28} the excess risk generally does not, due to the much higher incidence of CHD in older persons.^{31,38,39}

Other Lipid Constituents and Risk of Coronary Disease. The risk associated with high total cholesterol is primarily due to high levels of low-density lipoprotein cholesterol (LDL-C),¹ but there is a strong, independent, and inverse association between high-density lipoprotein cholesterol (HDL-C) levels and CHD risk.^{40–42} Low HDL-C increases risk even when cholesterol is below 200 mg/dL,⁴¹ a pattern present in up to 20% of men with confirmed CHD.⁴³ In many studies, measures of HDL-C or the ratio of total cholesterol to HDL-C are better predictors of CHD risk than is serum cholesterol alone.^{5,22,23,24a,41,44} High total cholesterol in association with high HDL-C (60 mg/dL) is common in older women (especially those taking estrogen) but is not associated with an increased risk for CHD.^{1,41} The importance of triglycerides as an independent risk factor for CHD remains uncertain.^{40,45} Three large studies reported strong associations between triglyceride levels over 200–300 mg/dL (2.26–3.39 mmol/L) and cardiovascular mortality in women,^{21,22,24} but other analyses found no association after controlling for obesity, fasting glucose, or low HDL-C.⁴⁶ The combination of high triglycerides and low HDL-C often occurs in association with other CHD risk factors such as hypertension and diabetes and is associated with a high risk of CHD.^{46a}

Prevalence of High Cholesterol and Low HDL-C. Serum total cholesterol and LDL-C increase 1–2 mg/dL per year in men from ages 20–40, 2 mg/dL per year in women from ages 40–60,⁴⁷ and an average 18% during the perimenopausal period, due in part to age-related increases in weight.⁴⁸ The prevalence of serum cholesterol 240 mg/dL or higher increases from 8–9% in adults under age 35 to nearly 25% for men age 55 and nearly 40% for women over 65.⁴⁹ Approximately 11% of men and 3% of women over age 20 have low HDL-C (<35 mg/dL) with desirable or borderline-high total cholesterol.⁴⁹

Accuracy of Screening Tests

Both total cholesterol and HDL-C can be measured in venipuncture or finger-stick specimens from fasting or nonfasting individuals. Due to normal physiologic variation and measurement error, a single measurement may not reflect the patient's true (or average) cholesterol level. Stress, minor illness, posture, and seasonal fluctuations may cause serum cholesterol to vary 4–11% within an individual.⁵⁰ Laboratory assays are subject to random errors, due to variation in sample collection, handling, and reagents, and to systematic errors (bias), due to methods that consistently overestimate or underestimate cholesterol values.⁵¹ In a survey of 5,000 clinical laboratories, 93% of the measurements were within 9% of a reference standard.⁵²

Desktop analyzers can produce reliable results, but some devices may not meet standards for accuracy.⁵³ Variation in training and operating technique can introduce additional error when instruments are used outside clinical laboratories.⁵⁴ Average bias for measurements based on capillary specimens compared to venous specimens was +4–7%.⁵⁵

As a result of these considerations, a single measure of serum cholesterol could vary as much as 14% from an individual's average value under acceptable laboratory conditions.⁵⁰ For an individual with a "true" cholesterol level of 200 mg/dL, the 95% range of expected values is 172–228 mg/dL.⁵⁶ Some authorities therefore recommend advising patients of their "cholesterol range," rather than a single value.⁵⁶ Where more precise estimates are necessary, an average of at least two measurements on two occasions has been recommended, and a third if the first two values differ by more than 16%.⁵⁰

Screening Children by Family History. Although cholesterol levels in childhood correlate moderately well with levels in adulthood (correlation coefficient 0.4–0.6), many children with elevated serum cholesterol (defined as serum cholesterol 200 mg/dL or LDL-C 130 mg/dL, the 90–95th percentile in U.S. children under 19 years)⁵⁷ do not have high cholesterol as adults.^{58–60} Furthermore, the association between childhood cholesterol levels and CHD in adults has not been studied. Because of the familial aggregation of CHD and hypercholesterolemia,^{57,61,62} some experts recommend screening for family history of either premature cardiovascular disease (age 55 or younger) or parental hypercholesterolemia (240 mg/dL) to identify a subset of children who are more likely to be at risk from hypercholesterolemia as adults.⁵⁷ Under this definition, only 25% of all children would be screened, but the predictive value of family history is limited: 81–90% of children with such histories have normal cholesterol.^{63–66} Even when parental cholesterol has been measured and found to be elevated, most children have normal cholesterol values.^{57,67,68}

Parental and childhood cholesterol levels are highest in heterozygous FH (estimated prevalence 1 in 500), which is strongly associated with premature CHD. Up to 50% of men with FH develop clinical CHD by age 50.^{69,70} Screening based on family history, as defined above, does not appear to be an efficient strategy for detecting FH, however. Many children would be screened, and few of those identified and treated for high cholesterol would have FH.⁷¹ By itself, a parental history of premature CHD is likely to detect less than half of all children with FH.⁷⁰ Tracing and screening families of index cases with FH may be more cost-effective than population screening for FH.⁷²

Screening for Other Lipid Abnormalities. Measurements of HDL-C and triglycerides are less reliable than measurement of total cholesterol due to greater

biologic and analytic variability.^{73,74} The 95% range of expected values for an individual with HDL-C of 37 mg/dL is 29–45 mg/dL.⁷⁵ A survey of 250 laboratories found that one third of all HDL-C measurements varied more than 10% from a reference value.⁷⁶ Triglycerides must be measured on fasting specimens. Even then, intraindividual variation is greater than 20%, and a single measure is inadequate to categorize levels as high or normal.^{73,74} Measurement of apolipoproteins (e.g., apoB) has been evaluated as a screening test for FH, familial coronary disease, and high LDL-C, but these assays are not yet widely available or adequately standardized.⁵⁷

Effectiveness of Early Detection

No long-term study has compared routine cholesterol screening to alternate strategies (selective case-finding or universal dietary advice without screening) with change in cholesterol levels or CHD incidence as an outcome. The increase in cholesterol screening over the past decade in the U.S. has been accompanied by significant improvements in dietary knowledge,⁷⁷ fat consumption,⁷⁸ average cholesterol levels,⁷⁹ and CHD mortality,¹⁴ but it is difficult to isolate the contribution of screening from other factors (e.g., public education, changes in food supply) that may account for these trends. In community- or practice-based trials, patients receiving risk-factor screening and targeted dietary advice had slightly lower average cholesterol levels (1–3%) than did unscreened controls at 1–3-year follow-up, but dietary interventions were limited.^{80–82} Whether screening improves the effectiveness of routine dietary advice has been examined in two short-term studies where all subjects received counseling about diet; cholesterol screening modestly improved mean cholesterol levels in one study but had no effect in the other.^{83,84} In a school-based study in which all children received similar health education, cardiovascular risk-factor screening (including cholesterol measurement) was associated with improved dietary knowledge and self-reported behavior, but changes in lipid levels were not assessed.⁸⁵

The primary evidence to support cholesterol screening is the ability of cholesterol-lowering interventions to reduce the risk of CHD in patients with high cholesterol. These benefits are now well established for persons with preexisting atherosclerotic vascular disease. In individual trials and overviews of studies enrolling persons with angina or prior myocardial infarction (MI), cholesterol-lowering treatments slowed the progression of atherosclerosis,⁸⁶ reduced the incidence of CHD,^{87,88} and reduced overall mortality.⁸⁹ In the first long-term trial of newer cholesterol-lowering drugs, treatment with simvastatin over 5.4 years reduced coronary mortality 42% and all-cause mortality 30% in 4,444 men and women with coronary disease.⁹⁰

The absolute benefit of treating high cholesterol in persons without cardiovascular disease, however, is much smaller due to the much lower risk of death or MI (annual CHD mortality 0.1–0.3% in middle-aged men with asymptomatic high cholesterol vs. 2–10% per year in patients with symptomatic CHD).⁹¹ The risks and benefits of lowering cholesterol in asymptomatic persons—primarily middle-aged men with very high cholesterol—have been examined in trials using medications, modified diets in institutional patients, or outpatient dietary counseling, and in overviews of these trials.

Trials of Cholesterol-Lowering Drugs in Asymptomatic Men. Three large, multicenter, placebo-controlled trials of lipid-lowering medications provide the best evidence that lowering cholesterol can reduce combined CHD incidence (fatal and nonfatal events) in asymptomatic persons. These trials enrolled hypercholesterolemic middle-aged men (age 30–59, mean cholesterol 246–289 mg/dL) and lowered total cholesterol 9–10% (and LDL-C 10–13%) over periods of 5–7 years. In the World Health Organization Cooperative Trial, treatment with clofibrate significantly reduced the incidence of nonfatal MI by 25%,⁹² but this benefit was offset by significant increases in noncardiac and total mortality (40% and 30% respectively, $p = 0.01$).⁹³ The Lipid Research Clinics (LRC) Coronary Primary Prevention Trial reported a significant 19% reduction in cumulative incidence of MI and sudden cardiac death in patients treated with cholestyramine over 7 years (7.0% vs. 8.6%).³⁶ In the Helsinki Heart Study, treatment with gemfibrozil significantly reduced the 5-year cumulative incidence of cardiac events by 34% (2.7% vs. 4.1%).⁹⁴ Most of the benefit of gemfibrozil was confined to men with a high ratio of LDL-C to HDL-C (> 5) and triglycerides > 200 mg/dL.⁹⁵ Effects on CHD mortality were not statistically significant in any of these trials. Two additional drug trials reported 1–3-year results in largely asymptomatic populations.^{96,97} Roughly 30% of subjects had CHD at entry, however, and these patients accounted for most of the coronary events during follow-up.

Trials of Diet in Institutionalized Persons. Demonstrating a clinical benefit of modern cholesterol-lowering diets in asymptomatic persons has proven difficult. In three controlled trials in institutionalized patients, fat-modified diets reduced serum cholesterol 12–14% with generally favorable effects on CHD over periods of up to 8 years.^{98–100} Each of these studies used diets high in polyunsaturated fat, which have been associated with adverse effects,¹⁵ and none excluded patients with CHD. As a result, their findings may not be applicable to currently recommended low-fat diets in asymptomatic persons.

Trials of Dietary Advice in Outpatients. The only trials to examine the clinical benefits of a diet low in total and saturated fat in persons without CHD are

multifactorial intervention trials, which offered dietary counseling, smoking cessation advice, and/or treatment of high blood pressure to middle-aged men.^{101–105} Among Norwegian smokers with very high cholesterol levels (mean 320 mg/dL) and fat consumption (44% calories), dietary advice lowered cholesterol 13% and, in conjunction with smoking cessation, reduced CHD incidence by 47%.¹⁰³ The remaining trials achieved much smaller (0–5%) reductions in cholesterol and insignificant effects on CHD; the benefits of intervention in some studies may have been limited by ineffective counseling and follow-up,^{101,104} lower cholesterol levels at baseline,¹⁰¹ or adverse effects of other therapies.^{102,105} In the most systematic test of dietary counseling in adults, 10 weekly group sessions and periodic individual counseling were provided over 6 years to over 6,000 men (mean cholesterol 253 mg/dL).¹⁰² Average cholesterol level declined 5% in men receiving counseling, but only 2% compared to controls. Greater changes were observed in men who lost at least 5 pounds and those with higher serum cholesterol at baseline,¹⁰⁶ but there was no significant reduction in CHD mortality or incidence in the intervention group.^{102,107}

Short-term metabolic studies and selected trials in patients with CHD indicate that reducing dietary saturated fat and/or increasing polyunsaturated fat intake can reduce elevated total and LDL-C as much as 10–20%.^{108–110} Due to variable compliance, trials of diet counseling in the primary care setting have achieved much smaller and inconsistent average reductions in serum cholesterol in asymptomatic persons (0–4%).^{80–82,111–116} Although larger changes have been reported in uncontrolled follow-up studies after cholesterol screening,^{117,118} these results may be biased by selective or short-term follow-up and regression to the mean in persons with high cholesterol. Ongoing studies are examining the efficacy of cholesterol screening and intervention in primary care settings in the U.S.¹¹⁹ More stringent diets can produce larger reductions in cholesterol,¹²⁰ but long-term data in asymptomatic persons are limited. Two trials in women at risk for breast cancer lowered total fat intake to 20% of calories and reduced total cholesterol 6–7% over 1–2 years.^{121,122}

Overviews of Cholesterol-Lowering Trials. At least 10 quantitative overviews (meta-analyses) of randomized trials have attempted to resolve uncertainties about the risks and benefits of lowering cholesterol, including effects on mortality.^{18,88,89,91,123–128} Three recent overviews provide the most comprehensive analyses of long-term cholesterol-lowering trials published through 1993; 35 diet and drug trials were included in one analysis,⁹¹ 28 in the second (excluding trials that used estrogens or thyroxine),^{18,89} and 22 in the third (all trials achieving at least a 4% reduction in cholesterol for at least 3 years).¹²⁸ These overviews support a dose-response relationship between change in serum cholesterol and reduction in CHD incidence (fatal and nonfatal

events combined) comparable to that predicted from epidemiologic studies: after 2–5 years of treatment, each 1% reduction in serum cholesterol yields a 2–3% reduction in total CHD, for both diet and drug interventions, and in patients with or without CHD at entry.^{18,89,128}

When only trials enrolling asymptomatic persons were analyzed, however, neither CHD mortality nor total mortality was significantly reduced by cholesterol lowering:¹²⁸ difference in total mortality among treated versus control subjects = +6%, 95% confidence interval (CI) –3% to +17%.⁸⁹ Moreover, noncardiac mortality was increased 20–24% among patients treated with lipid-lowering medications.^{89,91,128} While observing similar effects of treatment, each overview offered distinct interpretations of these findings. Law et al. concluded that the increase in noncoronary mortality was most likely due to chance: the finding was of borderline statistical significance ($p = 0.02$), did not reflect any consistent cause of excess mortality among trials, and was independent of compliance with therapy.⁸⁹ Gordon attributed adverse effects to trials employing hormones or fibrate medications.¹²⁸ Davey Smith et al. concluded that lipid-lowering drugs reduced overall mortality in high-risk persons (i.e., persons with CHD), but were harmful in those at lower risk.⁹¹ When trials were stratified by the observed CHD mortality in the control group, drug treatment was associated with a significant 20% increase in all-cause mortality in 10 trials enrolling low-risk subjects (CHD mortality <1% per year), including the WHO, LRC, and Helsinki studies.⁹¹ A single trial (the WHO clofibrate study) accounts for nearly half of all patient-years of treatment in persons without CHD¹²⁸ and has a strong influence on results of any meta-analysis.

Due to methodologic concerns about combining results from trials employing different cholesterol-lowering drugs and diets, meta-analysis cannot prove or disprove possible harms from lipid-lowering medications.¹²⁹ These analyses, however, illustrate the importance of underlying CHD risk in determining whether expected benefits are likely to justify possible risks of treatment. Even if drugs are safe, the margin of benefit may be small for many persons with asymptomatic hypercholesterolemia. In the LRC and Helsinki trials, preventing one coronary death required treating 300–400 middle-aged men for 5–7 years.^{36,94} The benefits of lipid-lowering medications on nonfatal CHD are more pronounced but must be weighed against the unpleasant and occasionally serious side effects of some drugs (see below).^{92,127,130} The newest class of lipid-lowering drugs, HMG-CoA reductase inhibitors or “statins,” lowers cholesterol more effectively and appears to be well-tolerated in trials lasting up to 6 years.^{90,97} These drugs are more likely to have significant effects on mortality in patients without CHD, but long-term trials of these agents in asymptomatic persons have yet to report results.¹³¹

Cholesterol Reduction in Women. Lipid-lowering medications and diet effectively lower cholesterol in women,¹³² but no trial has specifically examined the benefits of cholesterol reduction in asymptomatic women.¹³³ Trials that included female subjects with CHD observed qualitatively similar benefits of cholesterol reduction on angiographic or clinical endpoints in women and men.^{90,99,133} In the 4S trial, simvastatin significantly reduced CHD incidence, but not mortality, in women with CHD.⁹⁰ Two trials in women without CHD, with a cumulative enrollment of more than 6,000 women, observed no effect of drug or diet treatment on CHD incidence or mortality after 1–3 years;^{96,100} the short duration of follow-up may have limited the power of these studies to detect a difference.¹³³ Long-term data on drug therapy in women are limited, with the exception of estrogen therapy (see Chapter 68).

Cholesterol Reduction in Older Adults. The benefit of lowering cholesterol in older persons has been questioned due to the weak association between serum cholesterol and all-cause mortality after age 60.^{17,28,30} Associations between cholesterol and mortality in unselected elderly populations, however, are likely to be confounded by the increasing prevalence of chronic illnesses which increase mortality and independently lower serum cholesterol.^{26,134,135} Direct evidence that cholesterol reduction is beneficial in asymptomatic older persons is not yet available, but cholesterol-lowering diets and medications reduced overall mortality 26–30% in persons over 60 with clinical CHD.^{90,136,137} In two trials in patients without CHD that included older subjects, however, cholesterol reduction produced significant benefits in younger but not in older patients (over age 60 or 65).^{96,98} Newer cholesterol-lowering agents are efficacious and well-tolerated in older patients.^{90,138} A large multicenter trial is under way to examine the effectiveness of pravastatin and various antihypertensive medications in asymptomatic persons over age 60 with hypertension and high cholesterol.¹³⁸ There are few controlled trials of dietary counseling to lower cholesterol in older patients; no significant change in cholesterol levels was observed among rural Medicare recipients offered diet counseling¹³⁹ or older patients receiving diet counseling and placebo medication.¹³⁸

Cholesterol Reduction in Adolescents and Young Adults. Determining the benefits of lowering cholesterol in children, adolescents, and young adults is difficult, due to their low near-term risk of clinical coronary disease. The assumption that early treatment is more effective than treatment begun later in life⁵⁷ rests on observations that early atherosclerosis is present in many adolescents and young adults, is associated with lipid levels, progresses with age,⁶ and is difficult to reverse in middle age.⁸⁶ New evidence, however, suggests that much of the clinical benefit of lowering cholesterol can

be achieved within 2–5 years of initiating therapy.¹⁸ These benefits have been attributed to stabilizing “lipid-rich” lesions⁸⁷ and improving endothelial function,¹⁴⁰ and they suggest that the additional benefits of early drug therapy for hypercholesterolemia (i.e., before middle age) may not justify the added expense and possible risks of longer treatment. Intensive diet or drug intervention for adolescents and young adults with FH, although never tested in a prospective trial, has become standard treatment due to the very high levels of LDL-C and dramatically increased risk of premature CHD in persons with FH.^{11,71} Even in FH, however, most clinical events occur in middle age (i.e., after age 40), and risk is variable: MI was rare before age 30 in men in one study, and onset of CHD is later in women and nonsmokers with FH.^{69,141}

Modified diets lower cholesterol in young adults, but the contribution of universal screening in motivating risk reduction in young persons is uncertain. Neither a multiple-intervention trial in Australian workers¹⁴² nor a study of risk assessment in a general practice in the U.K.¹¹⁶ demonstrated that screening and dietary advice led to long-term reduction in cholesterol levels in younger men (under age 35–40). The effectiveness of screening and dietary counseling has not been adequately studied in young adults and cannot be predicted reliably from studies in middle-aged men.

Cholesterol Reduction in Children Dietary fat intake in children is associated with total cholesterol and LDL-C levels,^{143,144} but controlled trials have not consistently demonstrated that individual dietary counseling is effective in children.^{145–147} Results from the largest trial reported that children with elevated LDL-C who received intensive family-oriented dietary counseling (30 sessions over 3 years) experienced a significant but modest (3.2 mg/dL) reduction in mean LDL-C compared with controls.¹⁴⁸ Uncontrolled studies of dietician counseling for hyperlipidemic children and adolescents have reported larger short-term reductions in mean cholesterol and LDL-C,^{149–153} but such studies are prone to bias from regression to the mean and selective follow-up. Physical activity and fitness are associated with higher levels of HDL-C in children and adolescents, but controlled and uncontrolled trials^{154–159} have reported inconsistent effects of exercise interventions on lipids. Drug therapy effectively lowers cholesterol in children, but side effects limit compliance with bile-acid resins, the only therapy currently recommended for routine use in children.⁵⁷ In one study of 80 children with FH or FCH, only 13% were still compliant with resin therapy after 3 years.¹⁶⁰ Ongoing studies are examining the safety and efficacy of newer agents in children.

Potential Adverse Effects of Screening and Intervention. Measurement of serum cholesterol is safe and relatively inexpensive, but widespread screening may have some undesirable consequences. In populations in which the potential ben-

efits of early detection may be small (e.g., low-risk young persons), the possibility of harm may influence decisions about universal screening.¹⁶¹ Anecdotal reports have reported decreased well-being in persons diagnosed with high cholesterol (i.e., “labeling”),¹⁶² but a prospective study did not confirm this effect.¹⁶³ Other possible adverse effects of screening include inconvenience and expense of screening and follow-up, opportunity costs to the busy clinician, misinformation due to inaccurate results, and reduced attention to diet in persons with “desirable” cholesterol levels.¹⁶⁴ The importance of possible adverse effects of screening has not been systematically studied.

The safety of cholesterol-lowering interventions is especially important in children and young persons. Dietary restrictions may reduce intake of calories, calcium, vitamins, and iron in children,^{165–167} and failure-to-thrive due to excessively fat-restricted diets has been reported, albeit rarely, in children.^{168,169} In the most comprehensive trial of dietary intervention in children, however, no adverse effects on growth, sexual development, psychological measures, iron status, or blood micronutrients were detected at 3-year follow-up.¹⁴⁸ Other controlled studies also support the safety of properly performed dietary intervention in children.^{147,166,170} The elderly may also be at risk from modified diets if adequate intake of calories, calcium, and essential vitamins is not maintained, but these effects have not been directly examined.

The inappropriate use of drug therapy is of greater concern, especially in young persons in whom the benefit of early drug treatment may not justify the costs and possible risks.^{18,161} According to a national survey of pediatricians and family physicians, one in six regularly prescribed drugs for hypercholesterolemic children, and a substantial number did so based on inappropriate criteria, or used drugs not routinely recommended for children.¹⁷¹ Persons under age 40 accounted for over 1 million prescriptions for lipid-lowering drugs in 1992;¹⁷² gemfibrozil was the second most commonly prescribed lipid-lowering drug in the U.S. in 1992,¹⁷² despite limited indications for its use¹ and important safety concerns. Fibrate medications (e.g., clofibrate and gemfibrozil) have been associated with an increase in gallstone disease,⁹² adverse trends in CHD mortality^{93,173} and cancer mortality in individual trials,^{93,174} and a significant increase in non-coronary mortality in a recent overview of long-term trials.¹²⁸ HMG-CoA reductase inhibitors have not been associated with important adverse effects in trials lasting up to 6 years.⁹⁰ The safety of lifelong therapy with these agents cannot yet be determined; several medications in this class have been reported to cause liver tumors in animal studies.

Early Detection of Other Lipid Abnormalities. The importance of detecting low HDL-C or high triglycerides remains unproven, especially in persons with normal serum cholesterol. Weight loss in obese subjects,^{132,175} smoking

cessation, exercise,^{176,176a} and moderate alcohol consumption¹⁷⁷ can raise HDL-C and/or lower triglyceride levels. Some of these lifestyle interventions have only small effects, however, and most can be recommended independent of lipid levels. Most importantly, no trial has directly examined the benefit of raising HDL-C or lowering triglycerides.^{40,45} Secondary analyses of several trials have attributed varying proportions of the clinical benefit of drug therapy to increases in HDL-C,^{40,94,95} or reductions in triglycerides,¹³⁶ but all of the subjects had high total or LDL cholesterol. The benefit of drug treatment for low HDL-C and normal cholesterol has not been determined but is being studied in men with CHD.⁴³

Recommendations of Other Groups

The National Cholesterol Education Program Adult Treatment Panel II, convened by the National Heart, Lung, and Blood Institute, recommends routine measurement of nonfasting total cholesterol and HDL-C in all adults age 20 or older at least once every 5 years.^{1,178} The Canadian Task Force on the Periodic Health Examination concluded there was insufficient evidence to recommend routine cholesterol screening but endorsed case-finding in men 30–59 years old.¹²⁷ The American Academy of Family Physicians¹⁷⁹ recommends measurement of total cholesterol at least every 5 years in adults age 19 and older; these recommendations are under review. The American College of Obstetricians and Gynecologists recommends periodic screening of cholesterol in all women over age 20, and in selected high-risk adolescents.¹⁸⁰ In guidelines revised in 1995, the American College of Physicians (ACP) concluded that screening serum cholesterol was appropriate but not mandatory for asymptomatic men aged 35–65 and women aged 45–65; screening is not recommended for younger persons unless they are suspected of having a familial lipoprotein disorder or have multiple cardiac risk factors. The ACP concluded that evidence was not sufficient to recommend for or against screening asymptomatic persons between the ages of 65 and 75, but it recommends against screening after age 75.¹⁸¹

Selective screening of children and adolescents is recommended by the National Cholesterol Education Program Expert Panel on Blood Cholesterol Levels in Children and Adolescents,⁵⁷ the American Academy of Pediatrics (AAP),¹⁸² the Bright Futures guidelines,¹⁸³ the American Medical Association Guidelines for Adolescent and Preventive Services (GAPS),¹⁸⁴ and the American Academy of Family Physicians.¹⁷⁹ Screening with nonfasting cholesterol in all children and adolescents who have a parental history of hypercholesterolemia, and with fasting lipid profile in those with a family history of premature cardiovascular disease, is recommended. These organizations recommend that children who have multiple risk fac-

tors for CHD (such as smoking or obesity) and whose family history cannot be ascertained be screened at the discretion of the physician.

Discussion

Elevated serum cholesterol is an important risk factor for CHD in men and women in the U.S., and there is now good evidence that lowering serum cholesterol can reduce the risk of CHD. Whereas measures that lower serum cholesterol and provide other health benefits (e.g., regular physical activity, reducing dietary fat, and maintaining a healthy weight) should be encouraged in all persons, cholesterol screening can identify high-risk individuals who are most likely to benefit from individualized dietary counseling or drug treatment. In addition, screening may help clinicians and patients identify priorities for risk factor modification and reinforce public awareness of the importance of a healthy diet.

Some important questions remain, however, about routine lipid screening in asymptomatic and low-risk persons, including when to begin screening and which constituents to measure. Overall, evidence is strongest for screening for high serum cholesterol in middle-aged men (ages 35–65), based on the reduction in coronary morbidity in trials enrolling asymptomatic men with very high cholesterol (mean 280 mg/dL). The epidemiology and pathophysiology of CHD is similar in men and women, suggesting that reducing high cholesterol levels will also reduce CHD in asymptomatic women. Extrapolations to premenopausal women may not be appropriate, given their low risk of CHD and the apparent protective effects of estrogen on CHD incidence. The optimal age to screen women is not known; the later onset of hypercholesterolemia and CHD suggests that routine screening should begin around age 45.

Direct evidence that screening and intervention is effective in persons over age 65 is not yet available, but epidemiologic studies indicate that the risks of high cholesterol extend up to age 75. Given the high risk of CHD in the elderly, and the benefits of lowering cholesterol in symptomatic older men and women, screening may be reasonable in older persons who do not have major comorbid illnesses. Since individual cholesterol levels usually plateau by age 65 in women (and earlier in men), continued screening is less important in patients who have had desirable cholesterol levels throughout middle age.

There is not yet evidence that routine lipid screening is effective in reducing cholesterol levels or CHD risk in younger populations. Universal screening is an inefficient way to identify the small number of hypercholesterolemic young persons at risk for premature CHD, most of whom have multiple nonlipid risk factors or a history suggestive of familial dyslipidemia. Most “high-risk” young persons (excluding young men with FH)

have a near-term risk of CHD well below that of hypercholesterolemic middle-aged men,¹⁸⁵ and are not appropriate candidates for early drug therapy. Screening young persons can provide information to help stimulate lifestyle changes, but promoting a healthy lifestyle (e.g., healthy diet, regular physical activity, etc.) is important for all young persons, including the majority with “desirable” cholesterol levels. International comparisons suggest that cholesterol levels explain only part of the strong association between diet and heart disease.^{186a} As a result, it is uncertain whether routine cholesterol screening in low-risk younger populations is of sufficient benefit to justify the inconvenience, costs, and possible risks of screening and treatment. In a study modeling benefits of cholesterol screening, a conservative strategy of screening only middle-aged men and others with multiple CHD risk factors produced benefits comparable to screening all adults over age 20; if interventions had adverse effects on quality of life, the more conservative strategy was preferable.¹⁸⁶ Should future studies demonstrate that routine screening and targeted interventions are more effective in the primary care setting than universal dietary advice in young persons, this would provide some additional justification for early screening.

The benefits of screening children are even less certain. Progression of atherosclerosis in childhood is limited, many children with high cholesterol are not hypercholesterolemic as adults, and it is uncertain whether or not reducing cholesterol levels in childhood will significantly alter the risk of CHD many years later. Given the limited effectiveness of dietary counseling, poor compliance with currently recommended drug therapy, and the potential for adverse reactions in children, widespread pediatric screening might result in more harm than good.

The benefit of measuring HDL-C or triglycerides at initial screening is unproven. Measures to lower high triglycerides or raise HDL-C (e.g., weight reduction in obese persons and exercise) have relatively modest effects and should be encouraged regardless of lipid levels. Measures of HDL-C and lipoprotein analysis improve the estimation of coronary risk and should be obtained to guide treatment decisions in patients with high total cholesterol. There is, however, no evidence that they significantly improve the management of patients who do not have high total cholesterol.

While a single cholesterol test is relatively inexpensive, the cumulative costs of screening can be substantial under protocols calling for measurement of HDL-C, periodic screening, and detailed evaluation and treatment of the large population with high cholesterol. To be effective, dietary interventions require regular follow-up and reinforcement. Under optimistic assumptions, tailored dietary therapy in middle-aged men is estimated to cost more than \$20,000 per year of life gained, when costs of screening and follow-up are included.¹⁸⁷ Drug treatment of asymptomatic

middle-aged men (assuming no important adverse effects) has been estimated to cost at least \$50,000–90,000 per year of life saved.^{35,188} HMG-CoA reductase inhibitors are substantially more expensive than earlier medications, but they lower LDL-C more effectively and also raise HDL-C. These agents may improve the cost-effectiveness of drug therapy for asymptomatic hypercholesterolemia, especially in high-risk men,¹⁸⁹ but the long-term safety and effectiveness of these agents in persons without CHD have not yet been established.

CLINICAL INTERVENTION

Periodic screening for high blood cholesterol, using specimens obtained from fasting or nonfasting individuals, is recommended for all men ages 35–65 and women ages 45–65 (“B” recommendation). There is insufficient evidence to recommend for or against routine screening in asymptomatic persons after age 65, but screening may be considered on a case-by-case basis (“C” recommendation). Older persons with major CHD risk factors (smoking, hypertension, diabetes) who are otherwise healthy may be more likely to benefit from screening, based on their high risk of CHD and the proven benefits of lowering cholesterol in older persons with symptomatic CHD. Cholesterol levels are not a reliable predictor of risk after age 75, however. There is insufficient evidence to recommend routine screening in children, adolescents, or young adults (“C” recommendation). For adolescents and young adults who have a family history of very high cholesterol, premature CHD in a first-degree relative (before age 50 in men or age 60 in women), or major risk factors for CHD screening may be recommended on other grounds: the greater absolute risk attributable to high cholesterol in such persons, and the potential long-term benefits of early lifestyle interventions in young persons with high cholesterol. Recommendations against routine screening in children may be made on other grounds, including the costs and inconvenience of screening and follow-up, greater potential for adverse effects of treatment, and the uncertain long-term benefits of small reductions in childhood cholesterol levels.

The appropriate interval for periodic screening is not known. Periodic screening is most important when cholesterol levels are increasing (e.g., middle-aged men, perimenopausal women, and persons who have gained weight). An interval of 5 years has been recommended by experts,¹ but longer intervals may be reasonable in low-risk subjects (including those with previously desirable cholesterol levels).

There is insufficient evidence to recommend for or against routine measurement of HDL-C or triglycerides at initial screening (“C” recommendation). For high-risk persons (middle-aged persons with high cholesterol or multiple nonlipid risk factors for CHD), measurement of HDL-C

or lipoprotein analysis can be recommended to help identify individuals at highest risk of CHD, in whom individual diet or drug therapy may be indicated.

Decisions about interventions for high cholesterol should be based on at least two measures of cholesterol and assessment of the absolute risk of CHD in each individual. This assessment should take into account the age of the patient (higher risk in men over 45 and women over 55), results of lipoprotein analysis (or ratio of total cholesterol to HDL-C), and the presence and severity of other risk factors for CHD (see above).¹⁷⁸ More specific algorithms for risk assessment have been published.¹⁸⁵ Initial therapy for patients with elevated cholesterol is counseling to reduce consumption of fat (especially saturated fat) and promote weight loss in overweight persons. A two-step dietary program effective in lowering serum cholesterol has been described in detail elsewhere.¹ Benefits of drug therapy are likely to justify costs and potential risks only in persons at high risk of CHD (e.g., middle-aged men and postmenopausal women with very high cholesterol or multiple risk factors). The risks and benefits of drug therapy in asymptomatic persons over 65 have not yet been determined. In postmenopausal women with high cholesterol, estrogen therapy can lower LDL-C and raise HDL-C and is associated with lower risk of CHD in epidemiologic studies (see Chapter 68). Patients should receive information on the potential benefits, costs, and risks of long-term therapy before beginning treatment on cholesterol-lowering drugs.

All adults, adolescents, and children over age 2 years, including those with normal cholesterol levels, should receive periodic counseling regarding dietary intake of fat and saturated fat (see Chapter 56) and other measures to reduce the risk of coronary disease (see Chapters 3, 54, and 55).

The draft update of this chapter was prepared for the U.S. Preventive Services Task Force by David Atkins, MD, MPH, and Carolyn DiGiuseppi, MD, MPH.

REFERENCES

1. National Cholesterol Education Program. Second report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel II). Bethesda: National Heart, Lung, and Blood Institute, National Institutes of Health, 1993. (Publication no. 93-3095.)
2. National Center for Health Statistics. Annual summary of births, marriages, divorces, and deaths: United States, 1993. Monthly vital statistics report; vol 42 no 13. Hyattsville, MD: Public Health Service, 1994.
3. American Heart Association. Heart and stroke facts: 1995 statistical supplement. Dallas, TX: American Heart Association, 1994.
4. National Center for Health Statistics. Coronary heart disease incidence by sex—United States, 1971–87. *MMWR* 1992;41:526–529.
5. Bush TL, Fried LP, Barrett-Connor E. Cholesterol, lipoproteins, and coronary heart disease in women. *Clin Chem* 1988;34:B60–B70.

6. Pathological Determinants of Atherosclerosis in Youth (PDAY) Research Group. Relationship of atherosclerosis to serum lipoprotein cholesterol concentrations and smoking. *JAMA* 1990; 264:3018–3024.
7. Newman WP III, Wattigney W, Berenson GS. Autopsy studies in United States children and adolescents. Relationship of risk factors to atherosclerotic lesions. *Ann NY Acad Sci* 1991;623:16–25.
8. PDAY Research Group. Natural history of aortic and coronary atherosclerotic lesions in youth: findings from the PDAY study. *Arterioscler Thromb* 1993;131:291–298.
9. Strong JP, McGill HC Jr. The pediatric aspects of atherosclerosis. *J Atheroscler Res* 1969;9:251–265.
10. Stary HC. The sequence of cell and matrix changes in atherosclerotic lesions of coronary arteries in the first forty years of life. *Eur Heart J* 1990;11 (Suppl E):E3–E19.
11. Bild D, Williams R, Brewer H, et al. Identification and management of heterozygous familial hypercholesterolemia: summary and recommendations from an NHLBI workshop. *Am J Cardiol* 1993;72:1D–5D.
12. Cortner JA, Coates PM, Liacouras CA, et al. Familial combined hyperlipidemia in children: clinical expression, metabolic defects, and management. *J Pediatr* 1993;123:177–184.
13. LaRosa JC, Hunninghake D, Bush D, et al. The cholesterol facts. A joint statement by the American Heart Association and the National Heart, Lung, and Blood Institute. *Circulation* 1990; 81:1721–1733.
14. Stamler J, Stamler R, Brown V, et al. Serum cholesterol. Doing the right thing. *Circulation* 1993;88:1954–1960.
15. Grundy SM. Cholesterol and coronary heart disease. Future directions. *JAMA* 1990;264:3053–3059.
16. Neaton JD, Wentworth D, for the Multiple Risk Factor Intervention Trial Research Group. Serum cholesterol, blood pressure, cigarette smoking, and death from coronary heart disease. Overall findings and differences by age for 316,099 white men. *Arch Intern Med* 1992;152:56–64.
17. Anderson KM, Castelli WP, Levy D. Cholesterol and mortality: 30 years of follow-up from the Framingham Study. *JAMA* 1987;257:2176–2180.
18. Law MR, Wald NJ, Thompson SG. By how much and how quickly does reduction in serum cholesterol lower risk of ischemic heart disease? *BMJ* 1994;308:367–373.
19. Klag MJ, Ford DE, Mead LA, et al. Serum cholesterol in young men and subsequent cardiovascular disease. *N Engl J Med* 1993;328:313–318.
20. Davis CE, Rifkind BM, Brenner H, et al. A single cholesterol measurement underestimates the risk of coronary heart disease. *JAMA* 1991;264:3044–3046.
21. Stensvold I, Tverdal A, Urdal P, Graff-Iversen S. Non-fasting serum triglyceride concentration and mortality from coronary heart disease and any cause in middle aged Norwegian women. *BMJ* 1993;307:1318–1322.
22. Bass KM, Newschaffer CJ, Klag MJ, Bush TL. Plasma lipoprotein levels as predictors of cardiovascular death in women. *Arch Intern Med* 1993;153:2209–2216.
23. Manolio TA, Pearson TA, Wenger NK, et al. Cholesterol and heart disease in older persons and women: review of an NHLBI workshop. *Ann Epidemiol* 1992;2:161–176.
24. Bengtsson C, Bjorkelund C, Lapidus L, Lissner L. Association of serum lipid concentrations and obesity with mortality in women: 20 year follow up of participants in prospective population study in Gothenburg, Sweden. *BMJ* 1993;307:1385–1388.
- 24a. Grover SA, Palmer CS, Coupal L. Serum lipid screening to identify high-risk individuals for coronary death: the results of the Lipids Research Clinics Prevalence Cohort. *Arch Intern Med* 1994; 154:679–684.
25. Harris T, Cook EF, Kannel WB, et al. Proportional hazards analysis of risk factors for coronary heart disease in individuals aged 65 or older: the Framingham Heart Study. *J Am Geriatr Soc* 1988;36: 1023–1028.
26. Harris TB, Makuc DM, Kleinman JC, et al. Is the serum cholesterol-coronary heart disease relationship modified by activity level in older persons? *J Am Geriatr Soc* 1991;39:747–754.
27. Shipley MJ, Pocock SJ, Marmot MG. Does plasma cholesterol concentration predict mortality from coronary heart disease in elderly people? 18 year follow up in Whitehall study. *BMJ* 1991;303:89–92.
28. Kronmal RA, Cain KC, Ye Z, et al. Total serum cholesterol levels and mortality risk as a function of age. *Arch Intern Med* 1993;153:1065–1073.
29. Zimetbaum P, Frishman WH, Ooi WL, Derman MP, et al. Plasma lipids and lipoproteins and the incidence of cardiovascular disease in the very elderly. *Arterioscler Thromb* 1992;12:416–423.
30. Krumholz HM, Seeman TE, Merrill SS, et al. Lack of association between cholesterol and coronary

- heart disease mortality and morbidity and all-cause mortality in persons older than 70 years. *JAMA* 1994;272:1335-1340.
31. Hulley SB, Newman TB. Cholesterol in the elderly: is it important? *JAMA* 1994;272:1372-1373.
 32. Wald NJ, Law M, Watt H, et al. Apolipoproteins and ischemic heart disease: implications for screening. *Lancet* 1994;343:75-79.
 33. Toronto Working Group on Cholesterol Policy. Asymptomatic hypercholesterolemia: a clinical policy review. *J Clin Epidemiol* 1990;43:1029-1121.
 34. Pooling Project Research Group. Relationship of blood pressure, serum cholesterol, smoking habit, relative weight and ECG abnormalities to incidence of major coronary events. Final report of the Pooling Project. *J Chronic Dis* 1978;31:201-306.
 35. Goldman L, Gordon DJ, Rifkind BS, et al. Cost and health implications of cholesterol lowering. *Circulation* 1992;85:1960-1968.
 36. The Lipid Research Clinics Coronary Primary Prevention Trial Results. I. Reduction in incidence of coronary heart disease. *JAMA* 1984;251:351-364.
 37. Kannel WB, Wilson PF. Risk factors that attenuate the female coronary disease advantage. *Arch Intern Med* 1995;155:57-61.
 38. Benafante R, Reed D. Is elevated serum cholesterol a risk factor for coronary heart disease in the elderly? *JAMA* 1990;263:393-396.
 39. Rubin SM, Sidney S, Black DM, et al. High blood cholesterol in elderly men and the excess risk for coronary heart disease. *Ann Intern Med* 1990;113:916-920.
 40. NIH Consensus Development Panel on Triglyceride, High-density Lipoprotein and Coronary Heart Disease. Triglyceride, high-density lipoprotein, and coronary heart disease. *JAMA* 1993;269:505-510.
 41. Castelli WP, Garrison RJ, Wilson PW, et al. Incidence of coronary heart disease and lipoprotein cholesterol levels: the Framingham Study. *JAMA* 1986;256:2835-2838.
 42. Gordon DJ, Probstfield JL, Garrison RJ, et al. High-density lipoprotein cholesterol and cardiovascular disease: four prospective American studies. *Circulation* 1989;79:8-15.
 43. Rubins HB, Robins SJ, Iwane MK, et al. Rationale and design of the Department of Veterans Affairs High-Density Lipoprotein Cholesterol Intervention Trial (HIT) for secondary prevention of coronary artery disease. *Am J Cardiol* 1993;71:45-52.
 44. Kinosian B, Glick H, Garland G. Cholesterol and coronary heart disease: predicting risk by levels and ratios. *Ann Intern Med* 1994;121:641-647.
 45. Report of Meeting of Physicians and Scientists. Hypertriglyceridemia and vascular risk. *Lancet* 1993;342:781-787.
 46. Criqui MH, Heiss G, Cohn R. Plasma triglyceride level and mortality from coronary heart disease. *N Engl J Med* 1993;328:1220-1225.
 - 46a. Tenkanen L, Pietilä K, Manninen V. The triglyceride issue revisited. *Arch Intern Med* 1994;154:2714-2720.
 47. Chung SJ. Formulas predicting the percentile of serum cholesterol levels by age in adults. *Arch Pathol Lab Med* 1990;114:869-875.
 48. van Beresteijn ECH, Korevaar JC, Huijbregts PCW, et al. Premenopausal increase in serum cholesterol: a 10-year longitudinal study. *Am J Epidemiol* 1993;137:383-392.
 49. Sempos C, Cleeman JI, Carroll MD, et al. Prevalence of high blood cholesterol levels among US adults. An update based on guidelines from the second report of the National Cholesterol Education Program Adult Treatment Panel. *JAMA* 1993;269:3009-3014.
 50. Cooper GR, Myers GL, Smith SJ, et al. Blood lipid measurements. Variations and practical utility. *JAMA* 1992;267:1652-1660.
 51. Laboratory Standardization Panel, National Cholesterol Education Program. Recommendations for improving cholesterol measurement. Bethesda: National Institutes of Health, National Heart, Lung and Blood Institute, 1990. (NIH Publication no. 90-2964.)
 52. Ross JW, Myers GL, Gilmore BF, et al. Matrix effects and the accuracy of cholesterol analysis. *Arch Pathol Lab Med* 1993;117:393-400.
 53. Kaufman HW, McNamara JR, Anderson KM, et al. How reliably can compact chemistry analyzers measure lipids? *JAMA* 1990;263:1245-1249.
 54. Belsey R, Vandenbark M, Goitein RK, et al. Evaluation of a laboratory system intended for use in physicians' offices. II. Reliability of results produced by health care workers without formal or professional training. *JAMA* 1987;258:357-361.

55. Greenland P, Bowley NL, Meiklejohn B, et al. Blood cholesterol concentration: fingerstick plasma vs. venous serum sampling. *Clin Chem* 1990;36:628–630.
56. Belsey R, Baer DM. Cardiac risk classification based on lipid screening. *JAMA* 1990;263:1250–1252.
57. National Cholesterol Education Program. Report of the Expert Panel on Blood Cholesterol Levels in Children and Adolescents. Bethesda: National Institutes of Health, National Heart, Lung and Blood Institute, 1991. (NIH Publication no. 91-2732.)
58. Webber LS, Srinivasan SR, Wattigney WA, Berenson GS. Tracking of serum lipids and lipoproteins from childhood to adulthood. The Bogalusa Heart Study. *Am J Epidemiol* 1991;133:884–899.
59. Stuhldreher WL, Orchard TJ, Donahue RP, et al. Cholesterol screening in childhood: sixteen-year Beaver County Lipid Study experience. *J Pediatr* 1991;119:551–556.
60. Lauer RM, Clarke WR. Use of cholesterol measurements in childhood for the prediction of adult hypercholesterolemia. The Muscatine Study. *JAMA* 1990;264:3034–3038.
61. Mjos OD, Thelle DS, Forde OH, Vik-Mo H. Family study of high density lipoprotein cholesterol and the relation to age and sex. The Tromso Heart Study. *Acta Med Scand* 1977;201:323–329.
62. Croft JB, Cresanta JL, Webber LS, et al. Cardiovascular risk in parents of children with extreme lipoprotein cholesterol levels: the Bogalusa Heart Study. *South Med J* 1988;81:341–349.
63. Wong ND, Hei TK, Qaqundah PY, et al. Television viewing and pediatric hypercholesterolemia. *Pediatrics* 1992;90:75–79.
64. Bell MM, Joseph S. Screening 1140 fifth graders for hypercholesterolemia: family history inadequate to predict results. *J Am Board Fam Pract* 1990;3:259–263.
65. Davidson DM, Van Camp J, Iftner CA, et al. Family history fails to detect the majority of children with high capillary blood total cholesterol. *J School Health* 1991;61:75–80.
66. Steiner MH, Neinstein LS, Pennbridge J. Hypercholesterolemia in adolescents: effectiveness of screening strategies based on selected risk factors. *Pediatrics* 1991;88:269–275.
67. Shea S, Basch CE, Irigoyen M, et al. Failure of family history to predict high blood cholesterol among Hispanic preschool children. *Prev Med* 1990;19:443–455.
68. Benuck I, Gidding SS, Donovan M, et al. Usefulness of parental serum total cholesterol levels in identifying children with hypercholesterolemia. *Prev Cardiol* 1992;69:713–717.
69. Mabuchi H, Koizumi J, Shimizu M, et al. Development of coronary heart disease in familial hypercholesterolemia. *Circulation* 1989;79:225–232.
70. Scientific Steering Committee on behalf of the Simon Broome Register Group. Risk of fatal coronary heart disease in familial hypercholesterolemia. *BMJ* 1991;303:893–896.
71. Williams RR, Hunt SC, Schumacher C. Diagnosing heterozygous familial hypercholesterolemia using new practical criteria validated by molecular genetics. *Am J Cardiol* 1993;72:171–176.
72. Williams RR, Schumacher C, Barlow G, et al. Documented need for more effective diagnosis and treatment of familial hypercholesterolemia according to data from 502 heterozygotes in Utah. *Am J Cardiol* 1993; 72(suppl):18D–24D.
73. Smith S, Cooper G, Myers G, Sampson E. Biological variability in concentrations of serum lipids: sources of variation among results from published studies and composite predicted values. *Clin Chem* 1993;39:1012–1022.
74. Bachorik PS, Cloey TA, Finney CA, et al. Lipoprotein-cholesterol analysis during screening: accuracy and reliability. *Ann Intern Med* 1991;114:741–747.
75. Christenson RH, Roeback JR, Watson TE, Hla KM. Improving the reliability of total and high-density lipoprotein cholesterol measurements. *Arch Pathol Lab Med* 1991;115:1212–1216.
76. McQueen MJ, Henderson AR, Patten RL, et al. Results of a province-wide quality assurance program assessing the accuracy of cholesterol, triglycerides, and high-density lipoprotein cholesterol measurements and calculated low-density lipoprotein cholesterol in Ontario, using fresh human serum. *Arch Pathol Lab Med* 1991;115: 1217–1222.
77. Frank E, Winkleby M, Fortmann S, et al. Improved cholesterol-related knowledge and behavior and plasma cholesterol levels in adults during the 1980s. *JAMA* 1992;268:1566–1572.
78. National Heart, Lung and Blood Institute. Daily dietary fat and total food-energy intake—Third National Health and Nutrition Examination Survey, Phase 1, 1988–91. *MMWR* 1994;43:116–117, 123–125.
79. Johnson CL, Rifkind BM, Sempos CT, et al. Declining serum total cholesterol levels among US adults. The National Health and Nutrition Examination Surveys. *JAMA* 1993;269:3003–3008.
80. Imperial Cancer Research Fund OXCHECK Study Group. Effectiveness of health checks conducted by nurses in primary care: final results of the OXCHECK study. *BMJ* 1995;310:1099–1104.

81. Family Heart Study Group. Randomised controlled trial evaluating cardiovascular screening and intervention in general practice: principal results of British family heart study. *BMJ* 1994;308:313–320.
82. Murray D, Luepker R, Pirie P, et al. Systematic risk factor screening and education: a community-wide approach to prevention of coronary heart disease. *Prev Med* 1986;15:661–672.
83. Elton PJ, Ryman A, Hammer M, et al. Randomised controlled trial in northern England of the effect of a person knowing their own serum cholesterol level. *J Epidemiol Community Health* 1994; 48:22–25.
84. Robertson I, Phillips A, Mant D, et al. Motivational effects of cholesterol measurement in general practice health checks. *Br J Gen Pract* 1992;42:469–472.
85. Resnicow K, Corss D, Lacosse J, et al. Evaluation of a school-site cardiovascular risk factor screening intervention. *Prev Med* 1993;22:838–856.
86. Hong MK, Mintz GS, Popma JJ, et al. Limitations of angiography for analyzing coronary atherosclerosis progression or regression. *Ann Intern Med* 1994;121:348–354.
87. Brown BG, Zhao XQ, Sacco DE, Albers JJ. Lipid lowering and plaque progression. New insights into prevention of plaque disruption and clinical events in coronary disease. *Circulation* 1993;87:1781–1791.
88. Rossouw JE. The effects of lowering serum cholesterol on coronary heart disease risk. *Med Clin North Am* 1994;78:181–195.
89. Law M, Thompson S, Wald N. Assessing possible hazards of reducing serum cholesterol. *BMJ* 1994;308:373–379.
90. Scandinavian Simvastatin Survival Study Group. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994;344:1383–1389.
91. Davey Smith G, Song F, Sheldon TA. Cholesterol lowering and mortality: the importance of considering initial level of risk. *BMJ* 1993;306:1367–1373.
92. Report from the Committee of Principal Investigators. A cooperative trial in the primary prevention of ischaemic heart disease using clofibrate. *Br Heart J* 1978;40:1069–1118.
93. Heady JA, Morris JN, Oliver MF. WHO clofibrate/cholesterol trial: clarifications. *Lancet* 1992;340:1405–1406.
94. Frick MH, Elo O, Haapa K, et al. Helsinki Heart Study: primary prevention trial with gemfibrozil in middle-aged men with dyslipidemia. Safety of treatment, changes in risk factors, and incidence of coronary heart disease. *N Engl J Med* 1987;317:1237–1245.
95. Manninen V, Tenkanen L, Koskinen P, et al. Joint effects of serum triglyceride and LDL cholesterol and HDL cholesterol concentrations on coronary heart disease in the Helsinki Heart Study. Implications for treatment. *Circulation* 1992;85:37–45.
96. Dorr AE, Gundersen K, Schneider JC, et al. Colestipol hydrochloride in hypercholesterolemic patients—effect on serum cholesterol and mortality. *J Chronic Dis* 1978;31:5–14.
97. Bradford RH, Shear CL, Chremos AN, et al. Expanded clinical evaluation of lovastatin (EXCEL) study results. 1. Efficacy in modifying plasma lipoproteins and adverse event profile in 8245 patients with moderate hypercholesterolemia. *Arch Intern Med* 1991;151:43–49.
98. Dayton S, Pearce ML, Hashimoto S, et al. A controlled clinical trial of a diet high in unsaturated fat in preventing complications of atherosclerosis. *Circulation* 1969;40(Suppl II):1–63.
99. Miettinen M, Turpeinen O, Karvonen MJ, et al. Effect of cholesterol-lowering diet on mortality from coronary heart disease and other causes. *Lancet* 1972;2:835–838.
100. Frantz ID, Dawson EA, Ashman PL, et al. Test of effect of lipid lowering by diet on cardiovascular risk. The Minnesota Coronary Survey. *Arteriosclerosis* 1989;9:129–135.
101. World Health Organization European Collaborative Group. European collaborative trial of multifactorial prevention of coronary heart disease: final report on the 6-year results. *Lancet* 1986; 1:869–872.
102. Multiple Risk Factor Intervention Trial Research Group. Multiple Risk Factor Intervention Trial: risk factor changes and mortality results. *JAMA* 1982;248:1465–1477.
103. Hjerermann I, Velve Byre K, Holme I, et al. Effect of diet and smoking intervention on the incidence of coronary heart disease. Report from the Oslo Study Group of a randomized trial in healthy men. *Lancet* 1981;2:1303–1310.
104. Wilhelmssen L, Berglund G, Elmfeldt D, et al. The multifactor primary prevention trial in Goteborg, Sweden. *Eur Heart J* 1986;7:279–288.

105. Miettinen TA, Huttunen JK, Naukkarinen V, et al. Multifactorial prevention of cardiovascular diseases in middle-aged men. Risk factor changes, incidence and mortality. *JAMA* 1985;254:2097–2102.
106. Caggiula AW, Christakis G, Farrand M, et al. The Multiple Risk Factor Intervention Trial (MRFIT). IV. Intervention on blood lipids. *Prev Med* 1981;10:443–475.
107. Multiple Risk Factor Intervention Trial Research Group. Coronary heart disease death, nonfatal acute myocardial infarction and other clinical outcomes in the Multiple Risk Factor Intervention Trial. *Am J Cardiol* 1986;58:1–13.
108. Hegsted DM, Ausman LM, Johnson JA, et al. Dietary fat and serum lipids: an evaluation of the experimental data. *Am J Clin Nutr* 1993;57:875–883.
109. Truswell AS. Review of dietary intervention studies: effect on coronary events and on total mortality. *Aust NZ J Med* 1994;24:98–106.
110. Denke MA. Cholesterol-lowering diets. A review of the evidence. *Arch Intern Med* 1995;155:17–26.
111. Ramsay LE, Yeo WW, Jackson PR. Dietary reduction of serum cholesterol concentration: time to think again. *BMJ* 1991;303:953–957.
112. Murray DM, Kurth C, Mullis R, et al. Cholesterol reduction through low-intensity interventions: results from the Minnesota Heart Health Program. *Prev Med* 1990;19:181–189.
113. Lindholm LH, Ekblom T, Dash C, et al. The impact of health care advice given in primary care on cardiovascular risk. *BMJ* 1995;310:1105–1109.
114. Henkin Y, Garber DW, Osterlund LC, et al. Saturated fats, cholesterol and dietary compliance. *Arch Intern Med* 1992;152:1167–1174.
115. Bae CY, Keenan JM, Wenz J, et al. A clinical trial of the American Heart Association Step One Diet for treatment of hypercholesterolemia. *J Fam Pract* 1991;33:249–254.
116. Gibbins RL, Riley M, Brimble P. Effectiveness of programme for reducing cardiovascular risk for men in one general practice. *BMJ* 1993;306:1652–1656.
117. Wynder E, Harris R, Haley N. Population screening for plasma cholesterol: community results from Connecticut. *Am Heart J* 1989;117:649–656.
118. Gemson D, Sloan R, Messeri P, Goldberg I. A public health model for cardiovascular risk reduction. Impact of cholesterol screening with brief non-physician counseling. *Arch Intern Med* 1990;150:985–989.
119. Ammerman A, Caggiula A, Elmer PJ, et al. Putting medical practice guidelines into practice: the cholesterol model. *Am J Prev Med* 1994;10:209–216.
120. Ornish D, Brown SE, Scherwitz LW, et al. Can lifestyle changes reverse coronary heart disease? *Lancet* 1990;336:129–133.
121. Insull W, Henderson MM, Prentice RL, et al. Results of a randomized feasibility study of a low-fat diet. *Arch Intern Med* 1990;150:421–427.
122. Boyd NF, Cousins M, Beaton M, et al. Quantitative changes in dietary fat intake and serum cholesterol in women: results from a randomized, controlled trial. *Am J Clin Nutr* 1990;52:470–476.
123. Holme I. Relation of coronary heart disease incidence and total mortality to plasma cholesterol reduction in randomized trials. Use of meta-analysis. *Eur Heart J* 1993;69(suppl):S42–S47.
124. Yusuf S, Wittes J, Friedman L. Overview of results of randomized clinical trials in heart disease. II. Unstable angina, heart failure, primary prevention with aspirin and risk factor modification. *JAMA* 1988;260:2259–2263.
125. Muldoon MF, Manuck SB, Matthews KA. Lowering cholesterol concentrations and mortality: a quantitative review of primary prevention trials. *BMJ* 1990;301:309–314.
126. Ravnskov U. Cholesterol lowering trials in coronary heart disease: frequency of citation and outcome. *BMJ* 1992;305:15–19.
127. Canadian Task Force on the Periodic Health Examination. Canadian guide to clinical preventive health care. Ottawa: Canada Communication Group, 1994:649–669.
128. Gordon DJ. Cholesterol lowering and total mortality. In: Rifkind BM, ed. Lowering cholesterol in high-risk individuals and populations. New York: Marcel Dekker, 1995:333–348.
129. Thompson SG. Controversies in meta-analysis: the case of the trials of serum cholesterol reduction. *Stat Methods Med Res* 1993;2:173–192.
130. Knodel LC, Talbert RL. Adverse effects of hypolipidaemic drugs. *Med Toxicol* 1987;2:10–32.
131. Collins R, Keech A, Peto R, et al. Cholesterol and total mortality. Need for larger trials. *BMJ* 1993;306:1689.

132. Wood PD, Stefanick ML, Williams PT, et al. The effects on plasma lipoproteins of a prudent weight-reducing diet, with or without exercise, in overweight men and women. *N Engl J Med* 1991;325:461-466.
133. Moreno GT, Manson JE. Cholesterol and coronary heart disease in women: an overview of primary and secondary prevention. *Coron Artery Dis* 1993;4:580-587.
134. Manolio TA, Ettinger WH, Tracy RP, et al. Epidemiology of low cholesterol levels in older adults. The Cardiovascular Health Study. *Circulation* 1993;87:728-737.
135. Davey Smith G, Shipley MJ, Marmot MG, et al. Plasma cholesterol concentration and mortality. The Whitehall Study. *JAMA* 1992;267:70-76.
136. Carlson LA, Rosenhamer G. Reduction of mortality in the Stockholm Ischaemic Heart Disease Secondary Prevention Study by combined treatment with clofibrate and nicotinic acid. *Acta Med Scand* 1988;223:405-418.
137. Leren P. The effect of plasma cholesterol lowering diet in male survivors of MI. *Acta Med Scand* 1966;(Suppl 466):1-92.
138. LaRosa JC, Applegate W, Crouse JR III, et al. Cholesterol lowering in the elderly: results of the Cholesterol Reduction in Seniors Program (CRISP) pilot study. *Arch Intern Med* 1994;154:529-539.
139. Ives DG, Kuller LH, Traven N. Use and outcomes of cholesterol-lowering intervention in rural elderly subjects. *Am J Prev Med* 1993;9:274-281.
140. Levine GN, Keaney JF, Vita JA. Cholesterol reduction in cardiovascular disease. Clinical benefits and possible mechanisms. *N Engl J Med* 1995;332:512-521.
141. Hill JS, Hayden MR, Frohlich J, et al. Genetic and environmental factors affecting incidence of coronary artery disease in heterozygous familial hypercholesterolemia. *Arterioscler Thromb* 1991;11:290-297.
142. Gomel M, Oldenburg B, Simpson JM, et al. Work site cardiovascular risk reduction: a randomized trial of health risk assessment, education, counseling, and incentives. *Am J Public Health* 1993;83:1231-1238.
143. Nicklas TA, Farris RP, Smoak CG, et al. Dietary factors relate to cardiovascular risk factors in early life. Bogalusa Heart Study. *Arteriosclerosis* 1988;8:193-199.
144. Shea S, Basch CE, Irigoyen M, et al. Relationships of dietary fat consumption to serum total and low-density lipoprotein cholesterol in Hispanic preschool children. *Prev Med* 1991;20:237-249.
145. Walker R, Heller R, Redman S, et al. Reduction of ischemic heart disease risk markers in the teenage children of heart attack patients. *Prev Med* 1992;21:616-629.
146. Shannon BM, Tershakovec AM, Achterberg CL, et al. Cholesterol education for at-risk children. *Pediatrics* 1994;94:923-927.
147. Lapinleimu H, Viikari J, Jokinen E, et al. Prospective randomised trial in 1062 infants of diet low in saturated fat and cholesterol. *Lancet* 1995;345:471-476.
148. The Writing Group for the DISC Collaborative Research Group. Efficacy and safety of lowering dietary intake of fat and cholesterol in children with elevated low-density lipoprotein cholesterol: the Dietary Intervention Study in Children (DISC). *JAMA* 1995;273:1429-1435.
149. Quivers ES, Driscoll DJ, Garvey CD, et al. Variability in response to a low-fat, low-cholesterol diet in children with elevated low-density lipoprotein cholesterol levels. *Pediatrics* 1992;89:925-929.
150. Vartiainen E, Puska P, Pietinen P, et al. Effects of dietary fat modifications on serum lipids and blood pressure in children. *Acta Paediatr Scand* 1986;75:396-401.
151. Kuehl KS, Cockerham JT, Hitchings M, et al. Effective control of hypercholesterolemia in children with dietary interventions based in pediatric practice. *Prev Med* 1993;22:154-166.
152. Polonsky SM, Bellet PS, Sprecher DL. Primary hyperlipidemia in a pediatric population: classification and effect of dietary treatment. *Pediatrics* 1993;91:92-96.
153. Riggs TW, Holmes RD, Gendelman BS, Knust MM. Dietary treatment of childhood hypercholesterolemia. *Ann NY Acad Sci* 1991;623:457-459.
154. Linder CW, DuRant RH, Mahoney OM. The effect of physical conditioning on serum lipids and lipoproteins in white male adolescents. *Med Sci Sports Exerc* 1983;15:232-236.
155. Savage MP, Petratis MM, Thomson WH, et al. Exercise training effects on serum lipids of prepubescent boys and adult men. *Med Sci Sports Exerc* 1986;18:197-204.
156. Sasaki J, Shindo M, Tanaka H, et al. A long-term aerobic exercise program decreases the obesity index and increases the high-density lipoprotein cholesterol concentration in obese children. *Int J Obesity* 1987;11:339-345.

157. Widhalm K, Maxa E, Zyman H. Effect of diet and exercise upon the cholesterol and triglyceride content of plasma lipoproteins in overweight children. *Eur J Pediatr* 1978;127:121–126.
158. Fripp RR, Hodgson JL. Effect of resistive training on plasma lipid and lipoprotein levels in male adolescents. *J Pediatr* 1987;111:926–931.
159. Weltman A, Janney C, Rians CB, et al. The effects of hydraulic-resistance strength training on serum lipid levels in prepubertal boys. *Am J Dis Child* 1987;141:777–780.
160. Liacouras CA, Coates PM, Gallagher PR, et al. Use of cholestyramine in the treatment of children with familial combined hyperlipidemia. *J Pediatr* 1993;122:477–482.
161. Hulley SB, Newman TB, Grady D, et al. Should we be measuring blood cholesterol levels in young adults? *JAMA* 1993;269:1416–1419.
162. Brett AS. Psychologic effects of the diagnosis and treatment of hypercholesterolemia: lessons from case studies. *Am J Med* 1991;91:642–647.
163. Irvine MJ, Logan AG. Is knowing your cholesterol number harmful? *J Clin Epidemiol* 1994; 47:131–145.
164. Kinlay S, Heller R. Effectiveness and hazards of case finding for a high cholesterol concentration. *BMJ* 1990;300:1545–1547.
165. Shea S, Basch CE, Stein AD, et al. Is there a relationship between dietary fat and stature or growth in children three to five years of age? *Pediatrics* 1993;92:579–586.
166. Nicklas TA, Webber LS, Koschak ML, Berenson GS. Nutrient adequacy of low fat intakes for children: the Bogalusa Heart Study. *Pediatrics* 1992;89:221–228.
167. Copperman N, Schebendach J, Arden M, Jacobson MS. Nutrient adequacy of pediatric office-based dietary therapy of hyperlipidemia. *Ann NY Acad Sci* 1991;623:419–421.
168. Pugliese MT, Weyman-Daum M, Lifshitz F. Parental health beliefs as a cause of nonorganic failure to thrive. *Pediatrics* 1987;80:175–182.
169. Lifshitz F, Moses N. Growth failure: a complication of dietary treatment of hypercholesterolemia. *Am J Dis Child* 1989;143:537–542.
170. Fuchs GJ, Farris RP, DeWier M, et al. Effect of dietary fat on cardiovascular risk factors in infancy. *Pediatrics* 1994;93:756–763.
171. Kimm SYS, Payne GH, Lakatos E, et al. Primary care physicians and children's blood cholesterol. *Prev Med* 1992;21:191–202.
172. Editor. Use of cholesterol-lowering drugs. United States, 1992. *Stat Bull Metrop Life Insur Co* 1993;74:10–17.
173. Frick MH, Heinonen OP, Huttunen JK, et al. Efficacy of gemfibrozil in dyslipidemic subjects with suspected heart disease. An ancillary study in the Helsinki Heart Disease Study frame population. *Ann Med* 1993;25:41–45.
174. Huttunen JK, Heinonen OP, Manninen V, et al. The Helsinki Heart Study: an 8.5 year safety and mortality follow-up. *J Intern Med* 1994;235:31–39.
175. Dattilo AM, Kris-Etherton PM. Effects of weight reduction on blood lipids and lipoproteins: a meta-analysis. *Am J Clin Nutr* 1992;56:320–328.
176. Rosenson RS. Low levels of high-density lipoprotein cholesterol (hypoalphalipoproteinemia). *Arch Intern Med* 1993;153:1528–1538.
- 176a. King AC, Haskell WL, Young DR, et al. Long-term effects of varying intensities and formats of physical activity on participation rates, fitness, and lipoproteins in men and women aged 50 to 65 years. *Circulation* 1995; 91: 2596–2604.
177. Steinberg D, Pearson TA, Kuller LH. Alcohol and atherosclerosis. *Ann Intern Med* 1991; 114:967–976.
178. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Summary of the second report of the National Cholesterol Education Program (NCEP) Expert Panel on the Detection, Evaluation, and Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel II). *JAMA* 1993;269:3015–3023.
179. American Academy of Family Physicians. Age charts for periodic health examination. Kansas City, MO: American Academy of Family Physicians, 1994. (Reprint no. 510.)
180. American College of Obstetricians and Gynecologists. The obstetrician-gynecologist in primary preventive health care: a report of the ACOG Task Force on Primary and Preventive Health Care. Washington, DC: American College of Obstetricians and Gynecologists, 1993.
181. American College of Physicians. Serum cholesterol, high-density lipoprotein cholesterol, and triglycerides screening tests for the prevention of coronary heart disease in adults. *Ann Intern Med* 1996 (in press).

182. Committee on Nutrition, American Academy of Pediatrics. Statement on cholesterol. *Pediatrics* 1992;90:469–473.
183. Green M, ed. *Bright Futures: guidelines for health supervision of infants, children and adolescents*. Arlington VA: National Center for Education in Maternal and Child Health, 1994.
184. American Medical Association. *Guidelines for adolescent preventive services (GAPS): recommendations and rationale*. Chicago: American Medical Association, 1994.
185. Anderson KM, Wilson PWF, Odell PM, et al. An updated coronary risk profile. A statement for health professionals. *Circulation* 1991;83:356–362.
186. Krahn M, Naylor C, Basinski A, Detsky A. Comparison of an aggressive (U.S.) and a less aggressive (Canadian) policy for cholesterol screening and treatment. *Ann Intern Med* 1991;115:248–255.
- 186a. Verschuren WMM, Jacobs DR, Bloemberg BPM, et al. Serum total cholesterol and long-term coronary heart disease mortality in different cultures: twenty-five year follow-up of the Seven Countries Study. *JAMA* 1995;274:131–136.
187. Kristiansen IS, Eggen AE, Thelle DS. Cost effectiveness of incremental programmes for lowering serum cholesterol concentration: is individual intervention worth while? *BMJ* 1991;302:1119–1122.
188. Goldman L, Weinstein M, Goldman PA, Williams LW. Cost-effectiveness of HMG-CoA reductase inhibition for primary and secondary prevention of coronary heart diseases. *JAMA* 1991;265:1145–1151.
189. Hamilton VH, Racicot F-E, Zowall H, et al. The cost-effectiveness of HMG-CoA reductase inhibitors to prevent coronary heart disease. *JAMA* 1995;273:1032–1038.
190. Corti M-H, Guralnik JM, Salive ME, et al. HDL cholesterol predicts coronary heart disease mortality in older persons. *JAMA* 1995;274:539–544.