


# Effects of Cattle Manure Handling and Management on Fate and Transport of Hormones in the Feedlot and the Field

EPA-G2006-STAR-M1: Fate and Effects of Hormones in Waste From Concentrated Animal Feeding Operations (CAFOs)



# Research Team

- **Dan Snow**, SNR Water Sciences Laboratory
- **Shannon Bartelt-Hunt**, Civil Engineering
- **Bill Kranz**, Biological Systems Engineering
- **Terry Mader**, Animal Science
- **Charles Shapiro**, Soil Scientist - Crop Nutrition
- **David Shelton**, Biological Systems Engineering
- **Tian Zhang**, Civil Engineering
- **Simon von Donk**, Water Resources Engineer
- **David Tarkalson**, USDA-ARS NW Irrigation and Soils Research Lab, Kimberly, ID
- **Steve Ensley**, College of Veterinary Medicine, Iowa State University, Ames. IA



*Pioneering new frontiers.*

# EPA Request for Proposals

## Purpose of Program


- “Characterize the occurrence, magnitude, and extent of the impact of natural and synthetic steroid hormones in liquid and solid animal waste from concentrated animal feeding operations (CAFOs) on the environment and human health”
- “Determine the impact of current CAFO waste management strategies (i.e. storage and disposal) on the transport, fate, and effects of steroid hormones originating from CAFOs”

# Background

- Estimated 1.3 million livestock operations in the U.S
- Roughly 20% confined generate ~500 million tons of waste annually
- Large facilities have limited land available for effective use of the generated waste as fertilizer
- Potential for natural and synthetic steroid hormones in livestock waste to reach groundwater and surface waters through many pathways (e.g. surface runoff, leaching from holding tanks and lagoons and composting facilities)



**Cattle and Calves - Inventory: 2002**



**2000 Census  
People 1.7 million head  
Hogs 3.0 million head  
Cattle 6.2 million head**


1 Dot = 10,000 Cattle and Calves

United States Total  
95,497,994




# Steroid Hormones

- Steroids (terpenoid lipids) that act as hormones
- Hormone is a chemical messenger from one cell or group of cells to another (3 classes)
- Natural steroid hormones are generally synthesized from cholesterol




General  
chemical  
structure




# Steroid Hormones in Cattle Waste

- Endogenous (natural)


- Estrodiol
- Estrone
- Estriol
- Testosterone
- Androstenedione
- Progesterone




17β-estradiol



estrone




Testosterone




Progesterone

- Exogenous (synthetic)


- Trenbolone
- Zearalonol
- Melengestrol Acetate



17β-trenbolone



Melengestrol acetate




α-zearalonol

# Steroid hormones excreted in livestock manure

- Levels are highly variable
  - up to 1  $\mu\text{g/g}$
  - <100 ng/g
  - depends on compound, sex, diet, reproductive status
- Conjugated forms can be converted back to free hormones



Figure 2. Biliary 3-ketotrienic metabolites in the bile of a heifer



1. denotes percentage of biliary activity  
 2. double arrows indicate possible interconversions  
 3. denote tentatively identified structure configurations of 1-hydroxyls of compounds IX and XI are unspecified



# Chemical Properties

<i>Common Name</i>	<i>Aqueous solubility (<math>\mu\text{g/L}</math>)@ 25°C</i>	<i>Log P<sub>ow</sub></i>
<i>Estrogens</i>		
17 $\beta$ -estradiol	12,960	4.01
Estrone	12,420	3.13
Estriol	13,250	2.45
<i>Androgens</i>		
Testosterone	5,570	3.23
Androstenedione	27,000	2.9
Androsterone	8,750	3.69
<i>Progestins</i>		
Progesterone	6,600	4.0
<i>Synthetic Steroids/progestins</i>		
Trenbolone	20,000	--
Zearalanol	--	3.9
Melengestrol Acetate	1,060	3.67
Ethinylestradiol	483	3.67

**Atrazine solubility = 30,000  $\mu\text{g/L}$  Log Pow=2.3**

# Persistence





Figure 4. Degradation of (A) TbOH-17 $\alpha$ , (B) TbOH-17 $\beta$ , and (C) TbO in liquid manure during storage.

**“The Fate of Trenbolone Acetate and Melengestrol Acetate after Application as Growth Promoters in Cattle: Environmental Studies” *Bettina Schiffer, Andreas Daxenberger, Karsten Meyer, and Heinrich H.D. Meyer.* Environmental Health Perspectives • VOLUME 109 | NUMBER 11 | November 2001.**

# Project Objectives

- 1) *Quantify hormones in various stages of the manure pathway in cattle feedlots.*
- 2) *Determine the effects of different handling practices of cattle feedlot wastes on the stability and availability of hormones.*
- 3) *Determine the effects of different land application strategies on the fate and transport of hormones in vadose zone soils.*
- 4) *Determine if grasses grown in conservation buffers assimilate hormones.*

# Environmental Fate and Transport of Steroid Hormones



# Research Hypotheses

- 1) Hormone levels in manure are greater for cattle treated with hormones compared to untreated cattle.
- 2) Composting manure will facilitate the degradation of hormones in manure compared to stockpiling.
- 3) Hormone losses in runoff will be greater when manure is surface applied compared to when it is incorporated into the soil
- 4) Hormone losses in runoff will be related to the duration and timing of rainfall events with respect to land application of manure
- 5) Grasses commonly used in conservation buffer strips will assimilate hormones
- 6) Hormones will persist in soils and will leach through soils toward groundwater

# Project Tasks

- Task 1: Survey existing feedlots and agricultural fields to determine the fate of hormones in the manure handling pathway over a climatic gradient.
- Task 2: Quantify the fate of hormones as influenced by manure handling practices such as stockpiling, composting, and runoff retention basins.
- Task 3: Conduct runoff studies with controlled rainfall simulators to determine the effect of manure application strategies on hormone transport.
- Task 4: Quantify the uptake of hormones applied in manures to selected grass species commonly used in grass buffer strips.
- Task 5: Determine the fate and transport of hormones in vadose zone soil after land application of manure.

# Task 1. Survey of hormones in pens, run-off basins, stockpiles and fertilized cropland

- Where: Four University of Nebraska feedlots and manure-fertilized crop land
  - Collect manure from 3 pens and 2 stockpiles
  - Composite liquid from 2 run-off basins
  - Collect 12 composite samples at 3 depths from fields having received manure application within 1 month, 1 year, 2 year, and 3 years
- Collect data on feeding strategies, pen cleaning history, implant history, and manure handling





Figure 2. Locations of University of Nebraska Feedlots that will be used for Task 1.

# Task 2. Measure differences in cattle waste as influence by handling practices

- Where: Haskell Ag Lab
- 96 heifers split between six pens
- 48 will receive implants (TBA, Z, E) and MGA supplement
- Composite pen surface samples and fresh manure collected at 0, 7, 45, and 125 days after implanting
- Waste removed after 150 days, stockpiled and composted
- Samples of stockpiled and composted manure will be collected to determine changes in hormone levels
- Reactors used to characterize degradation rates and products




<sup>14</sup>C-labelled estradiol,  
testosterone, trenbolone

Aerobic vs anaerobic bioreactors





# Task 3. Effect of tillage on hormones levels in field run-off

- Where: Haskell Ag Lab
- Five manure sources
  - treated compost and stockpile manure
  - untreated compost and stockpile manure
  - control with no manure applied
- Compare surface application (no-till), single disk incorporation, mold-board plow
- Simulated rainfall




## Task 4. Uptake of hormones applied in manures to selected grass species commonly used in grass buffer strips

- Where: Haskell Ag Lab
- Hormone concentrations measured in buffer strip grasses receiving manure from steroid treated cattle
- Grass harvested after 30, 60, 90, 180, and 365 days after manure application




# Task 5. Fate and transport of hormones in vadose zone soil

- Where: WREC North Platte
- Lysimeter-instrumented test plots will be fertilized with stockpiled and composted manure from treated cattle
- Soil water will be collected from lysimeters over a 6-month period




# Method Development: Extraction and analysis of steroid hormones

## Solid Samples



## Liquid Samples











Evaporate solvent/add internal standards

Redissolve in mobile phase

Analyze by liquid chromatography-tandem mass spectrometry

# LC/MS/MS of Steroid Hormones



Quattro micro triple quadrupole

# LC/MS/MS Detection Limits


<b>Compound</b>	<b>Direct Injection</b>	<b>on-line SPE</b>
	<b>ppb</b>	<b>ppb</b>
17 $\beta$ -estradiol	1.07	0.0036
Estrone	0.77	0.0034
17 $\alpha$ -ethynyl estradiol	0.92	0.0056
Estriol	1.00	0.0064
$\alpha$ -Zearalanol	0.64	0.0032
Testosterone	1.46	0.0011
11-Ketotestosterone	0.58	0.0040
4-Androstenedione	1.71	0.0005
Progesterone	1.00	0.0008
Melengestrel Acetate	1.03	0.0016
17 $\beta$ -trenbolone	0.93	0.0018
Androsterone	3.16	0.0038

Spark Holland Symbiosys  
Environ Extraction System



# LC/MS/MS of steroid hormones

- Steroids
  - difficult to ionize by electrospray (ESI)
  - matrix effects severe
- Atmospheric pressure photoionization (APPI)
  - more selective
  - improved ionization efficiency
  - better for steroid analysis



<b>Preliminary On-Column Detection Limit (pg)</b>				
<b>Compound</b>	<b>ESI</b>	<b>Recovery</b>	<b>APPI</b>	<b>Recovery</b>
<b>Testosterone</b>	36.5	116 %	30.5	97 %
<b>11-Ketotestosterone</b>	14.5	91 %	13.8	103 %
<b>4-Androstenedione</b>	42.8	118 %	33.5	156 %
<b>Progesterone</b>	25.0	99 %	23.3	104 %
<b>Melengestrol Acetate</b>	25.8	96 %	20.5	125 %
<b>17<math>\beta</math>-Trenbolone</b>	23.3	154 %	18.0	142 %

# Contact Information

Daniel D. Snow  
Director and Associate Professor  
202 Water Sciences Laboratory  
University of Nebraska – Lincoln  
Lincoln, NE 68583-0844  
[dsnow1@unl.edu](mailto:dsnow1@unl.edu)  
402-472-7539



*Pioneering new frontiers.*