

Table 117. Death Rates from Malignant Neoplasms, by Selected Characteristics: 1990 to 2004

[Rates per 100,000 population. Beginning 2000, cause of death is coded according to ICD-10; for explanation, see text, this section. For explanation of age adjustment, see headnote, Table 102. See Appendix III]

Characteristic	1990	1995	2000	2001	2002	2003	2004
Total, age adjusted	216.0	209.9	199.6	196.0	193.5	190.1	185.8
Total, crude	203.2	202.2	196.5	194.4	193.2	191.5	188.6
Under 1 year	2.3	1.8	2.4	1.6	1.8	1.9	1.8
1 to 4 years	3.5	3.1	2.7	2.7	2.6	2.5	2.5
5 to 14 years	3.1	2.7	2.5	2.5	2.6	2.6	2.5
15 to 24 years	4.9	4.5	4.4	4.3	4.3	4.0	4.1
25 to 34 years	12.6	11.6	9.8	10.1	9.7	9.4	9.1
35 to 44 years	43.3	40.1	36.6	36.8	35.8	35.0	33.4
45 to 54 years	158.9	140.4	127.5	126.5	123.8	122.2	119.0
55 to 64 years	449.6	412.3	366.7	356.5	351.1	343.0	333.4
65 to 74 years	872.3	863.3	816.3	802.8	792.1	770.3	755.1
75 to 84 years	1,348.5	1,355.4	1,335.6	1,315.8	1,311.9	1,302.5	1,280.4
85 years and over	1,752.9	1,797.7	1,819.4	1,765.6	1,723.9	1,698.2	1,653.3
AGE-ADJUSTED RATES							
Male	280.4	267.5	248.9	243.7	238.9	233.3	227.7
Female	175.7	173.6	167.6	164.7	163.1	160.9	157.4
White male	272.2	260.6	243.9	239.2	235.2	230.1	224.4
Black male	397.9	374.3	340.3	330.9	319.6	308.8	301.2
American Indian, Alaska Native male	145.8	169.0	155.8	155.3	141.9	139.9	147.1
Asian, Pacific Islander male	172.5	164.3	150.8	147.0	137.9	137.2	136.3
Hispanic male, ¹	174.7	180.9	171.7	168.2	161.4	156.5	151.2
Non-Hispanic, White male, ¹	276.7	263.5	247.7	243.1	239.6	234.6	229.2
White female	174.0	172.1	166.9	163.9	162.4	160.2	157.0
Black female	205.9	203.8	193.8	191.3	190.3	187.7	182.5
American Indian, Alaska Native female	106.9	117.7	108.3	114.1	112.9	105.6	108.6
Asian, Pacific Islander female	103.0	107.4	100.7	99.3	95.9	96.7	92.0
Hispanic female, ¹	111.9	110.8	110.8	108.6	106.1	105.9	101.4
Non-Hispanic, White female, ¹	177.5	174.7	170.0	167.2	165.9	163.8	160.9
DEATH RATES FOR MALIGNANT NEOPLASM OF BREASTS FOR FEMALES							
All ages, age adjusted	33.3	30.5	26.8	26.0	25.6	25.3	24.4
All ages, crude	34.0	32.2	29.2	28.6	28.3	28.2	27.5
Under 25 years	(B)	(B)	(B)	(B)	(B)	(B)	(B)
25 to 34 years	2.9	2.6	2.3	2.4	2.1	2.1	2.0
35 to 44 years	17.8	14.9	12.4	12.4	12.0	12.2	11.3
45 to 54 years	45.4	41.0	33.0	32.8	31.4	30.4	29.3
55 to 64 years	78.6	69.4	59.3	57.5	56.2	56.6	55.8
65 to 74 years	111.7	102.8	88.3	85.8	84.4	82.6	81.6
75 to 84 years	146.3	140.1	128.9	125.8	125.9	123.7	119.5
85 years and over	196.8	200.2	205.7	188.9	191.5	189.4	178.6
DEATH RATES FOR MALIGNANT NEOPLASM OF TRACHEA, BRONCHUS, AND LUNG							
All ages, age adjusted	59.3	58.4	56.1	55.3	54.9	54.1	53.2
All ages, crude	56.8	56.8	55.3	54.8	57.7	54.4	53.8
Under 25 years	(Z)	(Z)	(Z)	(Z)	(Z)	(Z)	(Z)
25 to 34 years	0.7	0.6	0.5	0.4	0.4	0.4	0.4
35 to 44 years	6.8	6.0	6.1	6.2	6.0	5.6	5.5
45 to 54 years	46.8	37.5	31.6	30.7	30.3	30.3	29.9
55 to 64 years	160.6	141.6	122.4	117.7	115.3	111.0	106.9
65 to 74 years	288.4	295.4	284.2	279.7	275.0	269.3	265.5
75 to 84 years	333.3	358.9	370.8	371.4	377.6	377.8	373.6
85 years and over	242.5	279.9	302.1	302.7	297.2	298.9	297.5

B Base figure too small to meet statistical standards for reliability of a derived figure. Z Less than 0.05. ¹ Data for 1990 and 1995 exclude data from states lacking an Hispanic-origin item on their death certificates. See text, this section. Persons of Hispanic origin may be any race.