


EVALUATION OF THE BIRD CONSERVATION AREA CONCEPT IN THE NORTHERN TALLGRASS PRAIRIE

ANNUAL REPORT
2000



U.S. Department of the Interior
U.S. Geological Survey
Northern Prairie Wildlife Research Center
Jamestown, North Dakota

In cooperation with:
State University of New York
and
University of Minnesota, Crookston



EVALUATION OF THE BIRD CONSERVATION AREA CONCEPT
IN THE NORTHERN TALLGRASS PRAIRIE

ANNUAL REPORT: 2000

Maiken Winter

College of Environmental Science and Forestry
State University of New York
Syracuse, New York

Douglas H. Johnson

U.S. Geological Survey
Northern Prairie Wildlife Research Center
Jamestown, North Dakota

Therese M. Donovan

U.S. Geological Survey
Vermont Cooperative Fish and Wildlife Research Unit
University of Vermont
Burlington, Vermont

W. Daniel Svedarsky

Northwest Research and Outreach Center
University of Minnesota
Crookston, Minnesota

Funding provided by:
U.S. Geological Survey,
U.S. Fish and Wildlife Service,
Minnesota Department of Natural Resources

Cover Photo: Colleen LeFevre recording measurements on a mist-netted Savannah Sparrow
Cover Photo taken by: Maiken Winter

Executive summary

In 1998 we initiated a test of the concept that Bird Conservation Areas (BCA's) can maintain populations of breeding grassland birds. The underlying hypothesis is that large core areas of quality habitat (such as native prairie) that are surrounded by neutral habitats (such as small-grain fields), and that are isolated from hostile habitats (such as woody vegetation) will result in avian densities and reproductive rates sufficient to at least maintain population levels of breeding birds. This concept was proposed by the Midwest Working Group of Partners In Flight (e.g., Pashley and Fitzgerald 1996) and endorsed also by the Prairie Pothole Joint Venture of the North American Waterfowl Management Plan. This evaluation is being conducted in the northern tallgrass prairie, but the concept may be more generally applicable.

In 1999, we added 11 study plots in Sheyenne National Grassland in southeastern North Dakota to the existing 33 study plots in northwestern Minnesota. All study plots were assigned to one of four categories: 1) small core area surrounded by neutral landscape, 2) small core area surrounded by hostile landscape, 3) large core area surrounded by neutral landscape, and 4) large core area surrounded by hostile landscape. On each of the 44 study plots we collected data on population density of breeding birds by censusing each plot twice during the field season. Data on nesting success, predation, and brood parasitism were obtained from a subset of 29 study plots. In addition, we color-banded birds on four of the study plots, focusing on Clay-colored Sparrow, Savannah Sparrow and Bobolink. In 2000, we recorded 54 species on our census plots (compared to 41 and 53 in 1998 and 1999, respectively), found 679 nests of 39 species (compared to 293 of 19 species, and 793 of 34 species in 1998 and 1999), and color-banded 334 birds (compared to 263 in 1999).

Background and Justification

Grasslands are recognized by many as the most imperiled ecosystem worldwide (Samson and Knopf 1994, Noss et al. 1995). The avian assemblages associated with grasslands also are at risk - grassland bird populations have shown steeper, more consistent, and more geographically widespread declines than any other guild of North American species (Department of the Interior 1996). Breeding Bird Survey data from 1966-1993 indicate that almost 70 percent of 29 grassland bird species adequately

surveyed by BBS data had negative population trends; more than half of these was statistically significant.

In addition to range-wide population declines, the distribution and abundance of many grassland species are highly variable in space and time (Igl and Johnson 1999), which complicates conservation plans for grassland bird species. At both local and range-wide scales, variation in numbers from year to year may be driven by (1) climate patterns, which may significantly alter vegetation characteristics of the site and hence habitat cues used by birds in selecting breeding territories (Igl and Johnson 1999, Price 1995); (2) changes in the habitat caused by management actions or natural disturbances; (3) success of birds in raising young at that location in previous years, which may influence return rates and hence population stability at a site; or (4) changes in landscape structure caused by agriculture, urban sprawl, or other human activities. The relative importance of each of these factors has not been well established for grassland species, yet such knowledge is crucial to understanding patterns of range-wide population declines and local-scale fluctuations in grassland bird populations.

In an attempt to reverse population declines of grassland birds, the management concept of Bird Conservation Areas (BCA's) was suggested as a means to conserve grassland songbird populations (Pashley and Fitzgerald 1996). The notion behind BCA's is that core areas of quality habitat (such as native prairies) that are isolated from hostile habitats (such as woody vegetation) will result in reproductive rates sufficient to maintain population levels of breeding birds (Henderson and Sample 1995). The BCA concept implies that the value of high-quality core areas depends on the habitat composition of the landscape matrix in which the core areas are embedded. This concept is being promoted despite the absence of data that validate its usefulness in maintaining viable populations of grassland songbirds.

Moreover, the U.S. Department of the Interior (DOI) Conservation Strategy for declining birds in grassland ecosystems (DOI 1996) calls for information on the effects of habitat and landscape features on population viability of grassland birds. High-priority information needs identified by the DOI include effects of habitat structure and composition on avian communities and effects of landscape context (e.g., patch distribution, surrounding land use, and proximity to hostile environments) on avian numbers and nesting success. Furthermore, factors associated with highly variable population numbers (climate, habitat changes, nesting success) are needed to determine

causes of population stability or instability over time. This information is critical for developing long-term conservation objectives that will benefit grassland birds, but is lacking for many grassland bird species.

The BCA concept was proposed by the Midwest Working Group of Partners In Flight and supported also by the Prairie Pothole Joint Venture. It was included in the draft of the Landbird Conservation Plan for Physiographic Area 40: the Northern Tallgrass Prairie. This evaluation of the BCA concept in the northern tallgrass prairie is intended to determine whether BCA's do, in fact, meet their intended objectives. The effort addresses needs identified in the Landbird Conservation Plan by evaluating its assumptions.

For this evaluation, we are considering native prairie (parts of which may have been restored) to be high quality habitat; heavily wooded vegetation, which can harbor high numbers of predators and brood parasites, to constitute hostile habitat; and small-grain and hayfields to be neutral habitats.

Objectives

1. To estimate distribution, abundance, and reproductive success of grassland bird species in large and small core habitats embedded within hostile and neutral landscape matrices.
2. To estimate between-year site fidelity of grassland songbirds and factors that influence site fidelity.

Study Areas

In 2000 the study was conducted in three areas in the northern tallgrass prairie: (1) east of Moorhead, MN, in Becker, Mahnomon, and Clay counties; (2) east of Crookston, MN, in Polk County; and (3) in southeastern North Dakota at the Sheyenne National Grassland in Richland and Ransom counties. Study sites include tracts owned by the U.S. Fish and Wildlife Service, U.S. Forest Service, Minnesota Department of Natural Resources, and The Nature Conservancy (Table 1).

Methods

Study Design.--We are using a two-way factorial experimental design to address three major questions: (1) Does size of core habitat patch influence density and nesting success of birds? (2) Does landscape matrix (extent of woody vegetation surrounding the core habitat) influence density and nesting success of birds? and (3) Do patch size and landscape matrix show interactive effects? Main effects in the design are habitat size and landscape matrix, with several replicate plots within each size*landscape combination. All study plots were within native or restored prairie of similar vegetation structure and composition and varied between 1.0 and 16 ha in size.

In total, 21 study plots were established within core areas that are "small" in size (<50 ha), and 23 study plots were established within core areas that are "large" in size (>250 ha). We searched for nests in 15 study plots within small core areas and in 13 study plots within large core areas. Differences in abundance and reproductive measures between large and small core areas will be referred to as "main effects of size." Twenty-one study plots (11 in small core areas and 10 in large core areas) were established within hostile landscapes. Hostile landscapes include landscapes that contain large areas of woodland habitat within 5 km of the core habitat. Twenty-three study plots (10 in small core areas and 13 in large core areas) were established within neutral landscapes. Neutral landscapes include landscapes that consist of habitats that are thought to have little or no negative impact on bird populations within the core areas, such as small-grain fields, hay-meadows, or Conservation Reserve Program fields. Differences in abundance and reproductive measures between hostile and neutral landscapes will be referred to as "main effects of landscape composition." Interactive effects between core habitat size and landscape matrix also will be examined.

Field methods.--On all 44 study plots, we measured vegetation characteristics and bird abundance. Nesting success was investigated on a subset of 29 of the study plots (Table 1). Study plots were marked with flags or wooden laths at 50-m intervals along transects that were 100 m apart. Vegetation was assessed at 10 to 34 measuring points within each study plot, systematically located throughout each plot. The number of measuring points taken within a plot varied with the size of the study plot. Vegetation was measured once, in early to mid July. Measurements included vegetation height, percentage cover by

growth form (grass, forb, woody, bare ground, litter, and standing residual) based on a 20x50 cm Daubenmire frame, height-density (Robel readings), number of small (≤ 30 cm tall) and large (>30 cm tall) woody stems, and litter depth. Vegetation characteristics in each study plot were evaluated to determine the associations between habitat characteristics, local (patch size) features, landscape features, and density of each species.

Abundance of breeding birds of all species was determined on each study plot by strip-transect censuses (Stewart and Kantrud 1972). Censuses were conducted twice between 22 May and 24 June. The maximum count of a species was used to determine density (number of males/100 ha).

We assessed reproductive success of birds by searching for nests and monitoring eggs and young until fledging. The observers located nests by walking through fields with or without flushing-sticks and looking for nests after flushing or observing birds. Nests were marked with a flag 5 m to the north and were revisited every 3 days to ascertain its status and the incidence of brood parasitism. Nest success was determined using the Mayfield method (Mayfield 1961). A nest was considered successful if it fledged at least one young of the parental species, and it was considered parasitized if it contained at least one Brown-headed Cowbird egg or chick. We focused our nest searching efforts on three species: Savannah Sparrow, Clay-colored Sparrow, and Bobolink.

Nest vegetation was characterized within one week after activity at a nest had ceased. Vegetation was measured at five sites near each nest: directly at the nest and at a distance of 0.5 m from the nest in each cardinal direction. At each of the five points we measured vegetation in the same manner as described above for plot vegetation. Vegetation characteristics at the nest were evaluated to determine the associations between reproductive success by species and microhabitat (vegetation), local (patch size), and landscape features.

Four of the 44 study plots (two plots in large core areas surrounded by neutral landscape, and two plots in small core areas surrounded by hostile landscape) were designated as intensive sampling plots. On these we captured and marked birds to assess factors associated with population stability at a local site over time, again focusing on Savannah Sparrows, Clay-colored Sparrows, and Bobolinks. Birds were banded with an aluminum federal band and a combination of three color-bands. In 1999 and 2000, the four sites were monitored throughout the season to determine the number and identity of

individuals that returned from previous nesting seasons. Unbanded birds nesting on the plot were then targeted for banding. We then focused on monitoring banded birds to determine their season-long fecundity and movements within a plot. The goal of the intensive-sampling plots is to evaluate the number of young fledged per year and site fidelity for each adult of the three focal species. Site fidelity was measured in terms of returning to a site between years.

In 2000 we added one more aspect to our study: we employed miniature video cameras (Pietz and Granfors 2000) at nest sites of the three focal species to determine the types of nest predators that affect nests in our study area. Cameras were employed within and close to study sites in the Crookston area at nests of Savannah Sparrows, Clay-colored Sparrows, and Bobolinks.

Results

We recorded 54 species of birds on our study sites in 2000 (Table 2). The four most common species were Savannah Sparrow, Bobolink, Le Conte's Sparrow, and Clay-colored Sparrow. Le Conte's Sparrows and Savannah Sparrows generally occurred in each patch size in relatively high densities. However, Le Conte's Sparrows appeared to have higher densities in large plots within neutral landscapes, whereas Savannah Sparrows seemed to be affected primarily by landscape structure. In contrast to the Savannah Sparrow, Clay-colored Sparrows preferred hostile to neutral landscapes, also independent of patch size. Bobolinks seemed to prefer either large prairie patches or patches situated in neutral landscapes. The only grassland-nesting species that was restricted to large plots was the Greater Prairie-Chicken. Short-eared Owls were found only in plots in neutral landscapes.

Species composition differed slightly among the three regions (Table 2). Some species were detected in only one of the three regions, such as some duck species, Wilson's Phalarope, and Field Sparrow, which were found only at Sheyenne National Grassland. Further, species' densities varied among regions (Table 2); for example, Savannah and Le Conte's sparrows reached highest densities in the Crookston region, whereas Western Meadowlarks and Grasshopper Sparrows were recorded most frequently at Sheyenne National Grassland. Bobolinks and Clay-colored Sparrows had similar densities across the three regions.

We found 679 nests of 39 species (Table 3), compared to 1999 (where we found 793 nests of 34 species), and 1998 (where we found 295 nests of 19 species). The total number of nests found during the last three field seasons thus totals 1767. Most of the nests found belonged to the three focal species, Savannah Sparrow, Clay-colored Sparrow, and Bobolink (Table 3), so that the total number of nests belonging to one of the focal species is 1180. The most unusual nest we found this year was of a Dickcissel in the Glyndon area, a species that usually does not breed so far north.

Nesting success of the focal species was lowest for Bobolinks. For nests pooled over all plots, the Mayfield probability of daily nest survival was 0.937 for Savannah Sparrows, 0.923 for Clay-colored Sparrows, and 0.906 for Bobolinks. These translate to 21.8%, 28.1%, and 9.3% probabilities that a nest survives the incubation and nestling periods.

Nesting success varied greatly among regions, as did patterns in relation to patch size and landscape (Table 4). Contrary to last year's results, nesting success of both Savannah and Clay-colored sparrows did not seem to be affected by either patch size or landscape structure. Although patterns vary among regions, Bobolinks generally had higher nest success in large prairie patches surrounded by neutral landscape.

The main cause of nest failure was depredation (Table 5). Other causes of nest failure included 1) nest abandonment due to partial predation, cowbird parasitism, or unknown reasons, 2) cowbird predation (complete predation of nests by cowbirds occurred only at Sheyenne National Grassland), 3) trampling (mainly by cows), 4) weather related factors (mainly drowning in the nest), and 5) unknown causes.

Cowbird parasitism was low, with 6.4% (34/531) of all grassland passerine nests parasitized (Table 6). The most heavily parasitized species were Red-winged Blackbird, Western Meadowlark, Brewer's Blackbird, and Savannah Sparrow. All other grassland passerines had overall parasitism rates below 5%, even though in a single region parasitism rates were higher for some species.

We deployed cameras at 27 nests, 11 of which were depredated. Nest predators were of a wide array of species: thirteen-lined ground squirrel, short-tailed weasel, long-tailed weasel, American badger, red fox, striped skunk, Northern Harrier, American Kestrel, Brown-headed Cowbird, and one unidentified predator. In addition, a Plains garter snake caused one forced fledging. No single predator species prevailed for any bird species, or in any patch size/landscape configuration. On small neutral plots, 14

nests were videotaped; of those, five nests were depredated, by Brown-headed Cowbird, striped skunk, red fox, American badger, and an unidentified predator. On large neutral plots, cameras were employed at nine nests, of which three were depredated, two by short-tailed weasels and one by a thirteen-lined ground squirrel. On large hostile plots, cameras were employed at four nests, three of which were depredated, two by Northern Harriers and one by an American Kestrel. Due to limited accessibility, cameras were not set up on small hostile plots.

We color-banded 334 birds, 280 of which were focal species: 158 Savannah Sparrows, 92 Clay-colored Sparrows, and 30 Bobolinks. Twenty-nine of these banded birds had been banded in previous years: one female Le Conte's Sparrow (in 1999), 19 Savannah Sparrows (4 in 1998, and 15 in 1999, four Clay-colored Sparrows (1 in 1998, and 3 in 1999), and 5 Bobolinks (1 in 1998, and 4 in 1999). Four male Savannah Sparrows have been caught during each year of the study. We found 52 nests for which at least one of the two parents was banded (36 Savannah Sparrows, 11 Clay-colored Sparrows, 5 Bobolinks). Ten of these nests had both parents banded (6 Savannah Sparrow nests, 3 Clay-colored Sparrow nests, and 1 Bobolink nest). Contrary to 1999, we did not document any renesting or double-brooding in 2000.

Discussion and Future Plans

This study demonstrates the dynamic nature of grasslands and their avifauna. Not only do densities of breeding birds vary regionally and temporally, but so do interactions with their predators and brood parasites. This variability is reflected in the effects associated with habitat patch size and landscape features. After the third field season we have studied more than 1700 nests. As yet, however, it would be premature to conclude how grassland-nesting birds respond to patch size and landscape structure. The variation in avian density, nesting success, and their responses to explanatory variables demonstrates the importance of conducting studies over multiple years, as well as at multiple sites. Basing recommendations on the results of a single year of investigation could be unsound indeed.

In the fourth and final field season we will again census and nest-search the study sites in each of the three regions, color-band and resight birds on the four intensive study

plots in the Crookston region, and set up cameras at Savannah Sparrow and Clay-colored Sparrow nests to better determine the nest predators involved.

Acknowledgments

Primary funding for the third year of the evaluation was provided by the U.S. Geological Survey, through the Northern Prairie Wildlife Research Center, and the U.S. Fish and Wildlife Service, Region 3. Additional funding was provided by the U.S. Fish and Wildlife Service, Region 6.

Access to study areas was kindly granted by The Nature Conservancy, U.S. Fish and Wildlife Service, U.S. Forest Service, and Minnesota Department of Natural Resources (Division of Fish and Wildlife and Division of Parks and Recreation).

We appreciate efforts made by the field assistants: Catherine Brennan, Kathleen Calf, Andrea Claassen, Jamie Cymbaluk, Shawn Hawks, Colleen LeFevre, Sacha Mkheidze, Andy Radomski, Annika Samuelson, Dan Shustack, Brian Sprague, Sarah Stahl, Christine Vatovec, Jeremy Whitted, and Doug Zimmerman. Thanks to Peter Jones and Paul Prior for giving expert training in mist-netting and handling birds to the intensive study crew.

We are indebted for generous assistance with housing and other logistics to Rick Julian of the U.S. Fish and Wildlife Service, Brian Winter and Sonia Winter of The Nature Conservancy, and Bryan Stotts of the U.S. Forest Service.

We are grateful to Stephen J. Lewis of the U.S. Fish and Wildlife Service for his - continuing support of this evaluation.

Jane E. Austin, David P. Fellows, Pamela J. Pietz, and Marsha A. Sovada of the Northern Prairie Wildlife Research Center graciously loaned equipment and supplies. Pam Pietz and Diane A. Granfors were of invaluable help in collecting and organizing the data with the miniature camera system. Betty R. Euliss assisted the project in numerous ways.

Literature Cited

- Department of the Interior Grassland Bird Working Group (DOI). 1996. Declining birds in grassland ecosystems: a Department of the Interior conservation strategy. Fort Collins, Colorado.
- Henderson, R. A., and D. W. Sample. 1995. Grassland communities. Pages 116-129 in J. Addis, editor. Wisconsin's biodiversity as a management issue: a report to Department of Natural Resources' managers. Wisconsin Department of Natural Resources, Madison, Wisconsin.
- Igl, L. D. and D. H. Johnson. 1999. Le Conte's Sparrows breeding in Conservation Reserve Program fields: precipitation and patterns of population change. Pages 178-186 in P. D. Vickery and J. R. Herkert, editors. Ecology and conservation of grassland birds of the Western Hemisphere. *Studies in Avian Biology* 19.
- Mayfield, H. F. 1961. Nest success calculated from exposure. *Wilson Bulletin* 73:255-261.
- Noss, R. F., E. T. LaRoe III, and J. M. Scott. 1995. Endangered ecosystems of the United States: a preliminary assessment of loss and degradation. Biological Report 28. National Biological Service, Washington, D.C.
- Pietz, P. J., and D. A. Granfors. 2000. Identifying predators and fates of grassland passerine nests using miniature video cameras. *Journal of Wildlife Management* 64:71-87.
- Pashley, D., and J. Fitzgerald. 1996. Results of Prairie Pothole Joint Venture/Partners in Flight meeting, 11 & 12 July 1996. Unpublished memorandum.
- Price, J. T. 1995. Potential impacts of global climate change on the summer distributions of some North American grassland birds. Ph.D. dissertation. Wayne State University, Detroit, Michigan.
- Samson, F., and F. F. 1994. Prairie conservation in North America. *BioScience* 44:418-421.
- Stewart, R. E., and H. A. Kantrud. 1972. Population estimates of breeding birds in North Dakota. *Auk* 89:766-788.

Table 1. Study sites in northwestern Minnesota and southeastern North Dakota tallgrass prairie patches.

Prairie	Size category	Landscape	Plot size (ha)	Nest search?	Ownership
<u>Crookston Region (Minnesota)</u>					
Foxboro	small	neutral	6.00	Yes	SNA ¹
Mentor SW	small	neutral	9.00	Yes	WMA ²
Shypoke	small	neutral	6.00	Yes	WMA
Dugdale	small	neutral	6.00	No	WMA
Chicog NW	small	hostile	3.75	Yes	WMA
Mentor W	small	hostile	8.75	Yes	WMA
Tilden	small	hostile	8.25	Yes	WMA
Mentor NW	small	hostile	3.00	No	WMA
Pankratz S	large	neutral	16.00	Yes	TNC ³
Tympanuchus	large	neutral	16.00	Yes	WMA
Pankratz N	large	neutral	16.00	No	TNC
Pembina Trail	large	neutral	16.00	No	TNC
Burnham Creek	large	hostile	12.00	Yes	WMA
Pankratz/Kertzonville	large	hostile	15.00	Yes	TNC/WMA
Chicog W	large	hostile	12.00	No	WMA
<u>Glyndon Region (Minnesota)</u>					
Hoykens	small	neutral	5.75	Yes	WPA ⁴
"Refuge" ⁵	small	neutral	10.50	Yes	State Game Refuge
Spring Creek	small	neutral	12.00	Yes	WPA
Zimmerman	small	neutral	6.00	No	TNC
Eide	small	neutral	1.00	No	WPA
Sagebraaten	small	neutral	3.00	No	WPA
"Private" ⁶	small	hostile	4.50	Yes	WPA
Ulen	small	hostile	5.50	Yes	WMA
Buffalo E	small	hostile	3.00	No	State Park
Bicentennial	large	neutral	16.00	Yes	SNA
Bluestem N	large	neutral	15.25	Yes	TNC
Margherita	large	neutral	12.00	No	TNC
Blazing Star	large	neutral	16.00	No	TNC
Rice-Elliott	large	neutral	16.00	No	TNC
Bluestem S	large	hostile	15.00	Yes	TNC
Buffalo W	large	hostile	16.00	Yes	State Park
Flickertail	large	hostile	12.00	No	WPA
Fuglie	large	hostile	16.00	No	WPA

Table 1 cont. Study sites in northwestern Minnesota and southeastern North Dakota tallgrass prairie patches.

Prairie	Size category	Landscape	Plot size (ha)	Nest search	Ownership
<u>Sheyenne National Grassland (North Dakota)</u>					
Shrike	small	hostile	2.75	Yes	USFS
Pileated	small	hostile	7.00	Yes	USFS
Camp	small	hostile	7.25	Yes	USFS
Surprise	small	hostile	1.50	No	USFS
North	large	neutral	16.00	Yes	USFS
Highway	large	neutral	16.00	Yes	USFS
Plum	large	neutral	16.00	Yes	USFS
Southeast	large	neutral	16.00	No	USFS
Savannah	large	hostile	16.00	Yes	USFS
Eagle Point	large	hostile	14.00	Yes	USFS
Hammock	large	hostile	16.00	No	USFS

¹ Scientific and Natural Area

² Wildlife Management Area

³ The Nature Conservancy

⁴ Waterfowl Production Area

⁵ State Game Refuge on north side of Hwy 26 at large rock pile, east of Hwy 9.

⁶ State Game Refuge immediately south of the Shrike Unit

Table 2. Mean bird density/100 ha (\pm standard error) in small and large plots in neutral and hostile landscapes, by region, in 2000. Species are in taxonomic order.

Species	Crookston				Glyndon				Sheyenne		
	Small		Large		Small		Large		Small	Large	
	Neutral N= 4	Hostile N=4	Neutral N= 4	Hostile N= 3	Neutral N= 6	Hostile N= 3	Neutral N= 5	Hostile N= 4	Hostile N= 4	Neutral N= 5	Hostile N= 2
Mallard	0	0	0	0	18.2 " 16.4	0	2.9 " 1.8	0	0	2.5 " 1.5	3.6 " 3.6
Green-winged Teal	0	0	0	0	0	0	0	0	0	2.5 " 2.5	0
Blue-winged Teal	0	0	0	0	0	0	0	0	0	8.7 " 5.4	0
Northern Shoveler	0	0	0	0	0	0	0	0	0	1.2 " 1.2	3.6 " 3.6
Northern Pintail	0	0	0	0	0	0	0	0	0	1.2 " 1.2	0
Wood Duck	0	0	0	0	2.9 " 2.9	0	0	0	0	0	0
Sora	0	0	0	0	2.4 " 2.4	0	0	0	0	0	0
Marbled Godwit	4.2 " 4.2	0	1.6 " 1.6	0	0	0	4.2 " 1.7	0	0	5.0 " 2.3	0
Wilson's Phalarope	0	0	0	0	0	0	0	0	0	11.2 " 6.1	0
Killdeer	0	0	1.6 " 1.6	0	2.9 " 2.9	0	0	0	0	2.5 " 1.5	3.6 " 3.6
Common Snipe	6.9 " 4.2	6.7 " 6.7	7.8 " 1.6	2.2 " 2.2	2.9 " 2.9	0	1.2 " 1.2	0	0	1.2 " 1.2	14.0 " 0.2
Upland Sandpiper	2.8 " 2.8	0	3.1 " 1.8	2.2 " 2.2	0	0	2.5 " 1.5	0	0	1.2 " 1.2	6.9 " 6.9
Red-tailed Hawk	0	8.3 " 8.3	0	0	0	0	0	0	0	0	0
Northern Harrier	4.2 " 4.2	0	0	0	1.6 " 1.6	0	4.2 " 1.7	3.2 " 1.8	0	0	0
American Kestrel	0	0	0	0	0	0	1.2 " 1.2	1.6 " 1.6	0	0	0
Short-eared Owl	0	0	1.6 " 1.6	0	1.6 " 1.6	0	2.5 " 1.5	0	0	0	0
Greater Prairie-Chicken	0	0	9.4 " 7.4	0	0	0	2.9 " 1.8	0	0	1.2 " 1.2	3.4 " 3.4
Morning Dove	0	0	0	0	0	0	0	0	0	1.2 " 1.2	0
Northern Flicker	4.2 " 4.2	0	0	0	0	0	0	0	0	0	0
Eastern Kingbird	4.2 " 4.2	0	1.6 " 1.6	7.7 " 0.5	4.8 " 4.8	18.5 " 9.8	2.6 " 1.6	3.6 " 2.1	19.6 " 7.9	3.7 " 2.5	10.6 " 3.7
Willow Flycatcher	0	2.9 " 2.9	0	0	0	0	0	0	0	0	0
Alder Flycatcher	0	0	0	0	0	0	1.3 " 1.3	0	0	0	0
Bank Swallow	0	0	0	0	0	0	0	7.8 " 7.8	0	0	0
Barn Swallow	0	3.0 " 3.0	1.6 " 1.6	0	2.0 " 2.0	6.1 " 6.1	3.8 " 1.6	2.1 " 2.1	18.2 " 18.2	12.5 " 6.6	3.6 " 3.6
Tree Swallow	0	0	0	0	5.3 " 3.4	28.3 " 19.9	18.7 " 10.5	4.7 " 4.7	0	0	10.3 " 10.3
American Crow	0	0	0	0	0	0	0	0	7.1 " 7.1	0	0
Marsh Wren	0	0	0	5.3 " 5.3	17.6 " 7.2	0	0	0	0	1.2 " 1.2	0
Sedge Wren	30.6 " 17.8	8.6 " 8.6	56.2 " 28.1	26.0 " 13.9	5.5 " 3.9	14.8 " 14.8	11.3 " 5.7	17.2 " 11.8	0	5.0 " 5.0	6.9 " 6.9
House Wren	0	0	0	0	0	11.1 " 11.1	0	2.1 " 2.1	0	0	0

Table 2 cont. Mean bird density/100 ha (\pm standard error) in small and large plots in neutral and hostile landscapes, by region, in 2000. Species are in taxonomic order.

Species	Crookston				Glyndon				Sheyenne		
	Small		Large		Small		Large		Small	Large	
	Neutral N= 4	Hostile N=4	Neutral N= 4	Hostile N= 3	Neutral N= 6	Hostile N= 3	Neutral N= 5	Hostile N= 4	Hostile N= 4	Neutral N= 5	Hostile N= 2
American Robin	0	0	0	0	0	0	0.6 " 0.6	0	0	0	0
Gray Catbird	0	0	0	2.2 " 2.2	0	0	1.3 " 1.3	1.6 " 1.6	0	0	3.6 " 3.6
Warbling Vireo	0	0	0	0	0	0	0	0	9.1 " 9.1	0	3.6 " 3.6
Yellow Warbler	0	0	0	5.6 " 5.6	5.3 " 3.4	17.2 " 9.6	7.8 " 5.3	2.1 " 2.1	0	0	3.6 " 3.6
Common Yellowthroat	8.3 " 4.8	6.7 " 6.7	7.8 " 4.7	9.9 " 1.7	26.5 " 9.4	13.5 " 6.8	16.7 " 7.7	18.8 " 9.8	6.9 " 6.9	2.5 " 2.5	10.7 " 10.7
Grasshopper Sparrow	0	15.1 " 15.1	3.1 " 1.8	0	11.1 " 11.1	0	37.2 " 15.8	0	48.9 " 30.2	67.5 " 21.4	59.7 " 4.6
Le Conte's Sparrow	55.6 " 26.1	65.9 " 31.8	107.8 " 29.1	79.4 " 39.9	19.5 " 8.6	52.5 " 27.2	76.4 " 24.4	19.5 " 7.1	0	31.2 " 8.8	24.7 " 11.0
Vesper Sparrow	0	0	0	2.7 " 2.7	0	0	0	0	17.7 " 13.5	0	3.6 " 3.6
Savannah Sparrow	162.5 " 34.9	73.5 " 10.5	173.4 " 5.3	86.8 " 11.0	91.1 " 25.8	46.5 " 23.3	104.0 " 36.8	34.2 " 12.0	0	101.2 " 18.0	63.8 " 36.2
Song Sparrow	0	0	0	0	12.9 " 6.8	17.2 " 9.6	1.3 " 1.3	13.7 " 5.2	0	0	3.6 " 3.6
Field Sparrow	0	0	0	0	0	0	0	0	10.5 " 6.6	0	0
Clay-colored Sparrow	20.8 " 8.0	52.9 " 10.5	34.4 " 14.5	58.3 " 32.8	53.9 " 15.9	76.1 " 6.6	38.5 " 17.3	76.2 " 27.8	87.1 " 30.6	10.0 " 8.5	46.3 " 39.4
Swamp Sparrow	8.3 " 8.3	0	0	0	10.8 " 5.5	0	0	0	0	0	0
Nelson's Sharp-tailed Sparrow	0	0	1.6 " 1.6	0	0	0	0	0	0	0	0
Bobolink male	22.2 " 15.7	3.0 " 3.0	51.6 " 6.4	27.4 " 3.0	73.0 " 14.6	39.1 " 21.2	40.6 " 3.4	26.0 " 14.8	47.1 " 31.5	60.0 " 23.2	10.2 " 1.2
Western Meadowlark	6.9 " 4.2	3.0 " 3.0	10.9 " 3.0	0	3.5 " 2.3	0	9.7 " 2.1	1.6 " 1.6	16.1 " 7.5	31.2 " 3.9	24.7 " 11.0
Red-winged Blackbird	15.3 " 6.9	0	10.9 " 5.3	5.3 " 5.3	81.3 " 20.1	0	16.2 " 9.6	0	0	41.2 " 18.1	53.1 " 25.5
Yellow-headed Blackbird	0	0	0	0	0	0	0	0	0	5.0 " 5.0	0
Brewer's Blackbird	12.5 " 12.5	0	3.1 " 3.1	0	0	0	0	0	0	0	0
Brown-headed Cowbird	8.3 " 8.3	0	3.1 " 3.1	0	2.9 " 2.9	11.1 " 11.1	0	1.6 " 1.6	9.1 " 9.1	18.7 " 6.2	21.3 " 14.4
Brown-headed Cowbird male	5.6 " 5.6	0	4.7 " 4.7	2.2 " 2.2	6.9 " 3.2	11.1 " 11.1	8.2 " 3.3	3.3 " 3.3	9.1 " 9.1	22.5 " 6.1	21.3 " 14.4
Common Grackle	0	0	0	0	0	0	0	0	0	2.5 " 1.5	17.9 " 17.9
Northern Oriole	0	0	0	0	0	0	0	0	0	0	3.6 " 3.6
American Goldfinch	0	12.7 " 7.4	0	0	9.5 " 5.4	29.6 " 19.6	3.9 " 2.6	0	16.2 " 9.5	0	10.7 " 10.7
Total number of species	19	13	22	16	27	16	28	20	15	29	29

Table 3. Numbers of nests found, by study area, in 1998-2000, ordered by total number.

Species	Crookston			Glyndon			Sheyenne		Total			All
	1998	1999	2000	1998	1999	2000	1999	2000	1998	1999	2000	
Clay-colored Sparrow	20	86	27	71	119	93	56	66	91	261	186	538
Savannah Sparrow	41	150	141	21	25	25	27	44	62	202	210	474
Bobolink	23	27	24	25	20	18	9	22	48	56	64	168
Red-winged Blackbird	6	2	7	6	4	4	28	13	12	34	24	70
Western Meadowlark	1	6	2	3	1	2	25	17	4	32	21	57
Mallard	5	8	3	11	5	6	12	5	16	25	14	55
Le Conte's Sparrow	5	21	5	2	0	4	0	0	7	21	9	37
Brewer's Blackbird	12	8	9	4	3	0	0	0	16	11	9	36
Blue-winged Teal	0	0	0	3	1	4	17	10	3	18	14	35
Song Sparrow	2	2	0	5	8	10	3	0	7	13	10	30
Field Sparrow	0	0	0	0	0	0	13	16	0	13	16	29
Grasshopper Sparrow	0	0	1	3	4	0	8	8	3	12	9	24
Sedge Wren	0	10	6	3	3	1	0	0	3	13	7	23
Upland Sandpiper	0	1	6	7	3	2	3	1	7	7	9	23
Lark Sparrow	0	0	0	0	0	0	16	3	0	16	3	19
Vesper Sparrow	0	0	0	0	0	0	8	10	0	8	10	18
Mourning Dove	0	0	0	2	2	0	4	9	2	6	9	17
Eastern Kingbird	0	0	0	4	5	2	1	4	4	6	6	16
American Goldfinch	0	0	5	0	4	5	0	1	0	4	11	15
Greater Prairie-Chicken	2	1	3	2	2	2	0	0	4	3	5	12
Common Yellowthroat	0	0	0	2	4	3	0	1	2	4	4	10
Yellow Warbler	0	1	0	0	2	6	0	0	0	3	6	9
Common Snipe	2	3	0	1	1	0	1	1	3	5	1	9
Marbled Godwit	0	0	1	1	1	0	2	1	1	3	2	6
American Woodcock	0	1	0	0	1	0	0	4	0	2	4	6
Northern Harrier	0	2	1	0	1	1	0	0	0	3	2	5
Northern Shoveler	0	0	0	0	0	0	1	3	0	1	3	4
Gadwall	0	0	0	0	0	0	3	0	0	3	0	3
Wilson's Phalarope	0	0	0	0	0	0	2	1	0	2	1	3
Gray Catbird	0	1	0	0	0	2	0	0	0	1	2	3
Swamp Sparrow	0	1	0	0	1	1	0	0	0	2	1	3
Green-winged Teal	0	0	0	0	0	0	0	2	0	0	2	2
American Bittern	0	0	0	0	0	0	1	0	0	1	0	1
Redhead	0	0	0	0	0	1	0	0	0	0	1	1
Tree Swallow	0	0	0	0	0	1	0	0	0	0	1	1
Brown Thrasher	0	0	0	0	1	0	0	0	0	1	0	1
Marsh Wren	0	0	0	0	0	1	0	0	0	0	1	1
Chipping Sparrow	0	0	0	0	0	0	1	0	0	1	0	1
Yellow-headed Blackbird	0	0	0	0	0	0	0	1	0	0	1	1
Dickcissel	0	0	0	0	0	1	0	0	0	0	1	1
Total	119	331	241	176	221	195	241	243	295	793	679	1767

Table 4. Mayfield estimates of daily nest survival (May), its standard error (SE), percentage of nests parasitized by Brown-headed Cowbirds (Cow), and sample size (N) in small and large plots in neutral and hostile landscapes, by region, in 2000.

a) Crookston Region

Species	Small						Large					
	Neutral			Hostile			Neutral			Hostile		
	May " SE	Cow	N	May " SE	Cow	N	May " SE	Cow	N	May " SE	Cow	N
Savannah Sparrow	0.94 " 0.01	0	45	0.93 " 0.02	1	10	0.93 " 0.01	3	64	0.86 " 0.03	0	19
Clay-colored Sparrow	0.77 " 0.08	0	8	0.95 " 0.03	0	6	0.87 " 0.05	20	5	0.94 " 0.04	0	6
Bobolink	0.70 " 0.12	0	4	-	0	1	0.87 " 0.03	0	13	0.85 " 0.07	0	5

b) Glyndon Region

Species	Small						Large					
	Neutral			Hostile			Neutral			Hostile		
	May " SE	Cow	N	May " SE	Cow	N	May " SE	Cow	N	May " SE	Cow	N
Savannah Sparrow	0.93 " 0.03	0	9	0.94 " 0.04	0	3	0.88 " 0.04	10	10	0.92 " 0.05	0	3
Clay-colored Sparrow	0.94 " 0.02	0	15	0.94 " 0.02	0	16	0.92 " 0.02	0	28	0.91 " 0.02	3.4	34
Bobolink	0.92 " 0.04	0	5	0.82 " 0.07	16.6	6	0.94 " 0.03	0	6	-	0	1

c) Sheyenne National Grassland

Species	Small						Large					
	Neutral			Hostile			Neutral			Hostile		
	May " SE	Cow	N	May " SE	Cow	N	May " SE	Cow	N	May " SE	Cow	N
Savannah Sparrow	-	-	-	-	-	0	0.93 " 0.01	39.4	33	0.93 " 0.03	0	11
Clay-colored Sparrow	-	-	-	0.92 " 0.02	3.6	36	-	-	0	0.91 " 0.02	19.2	26
Bobolink	-	-	-	0.89 " 0.05	0	4	0.98 " 0.02	0	10	0.80 " 0.06	0	8

Table 5. Causes of nest failure, by study region, in 2000.

	Crookston	Glyndon	Sheyenne
Depredated:	136	83	90
Egg-laying stage	7	0	1
Incubation stage	42	41	64
Nestling stage	37	42	25
Cowbird predation	0	0	18
Weather-related	10	7	0
Trampled	1	1	13
Unknown	0	1	3
Abandoned:	24	17	40
Partial predation	3	7	23
Partial cowbird predation	0	0	4
Cowbird parasitism	0	1	5
Unknown	21	9	8

Table 6. Percentage of grassland passerine nests that were parasitized by Brown-headed Cowbirds in 2000. Total numbers of nests are shown in parentheses.

Species	Crookston	Glyndon	Sheyenne	Total
	% (N)	% (N)	% (N)	% (N)
Red-winged Blackbird	14.3 (7)	0 (4)	23.1 (13)	16.7 (24)
Western Meadowlark	0 (2)	50.0 (2)	11.8 (17)	14.3 (21)
Brewer's Blackbird	11.1 (9)	-	-	11.1 (9)
Savannah Sparrow	2.3 (129)	4 (25)	29.5 (44)	8.6 (198)
Clay-colored Sparrow	4.8 (21)	1.1 (93)	9.7 (62)	4.5 (176)
Bobolink	0 (19)	5.6 (18)	0 (22)	1.7 (59)
Field Sparrow	-	-	0 (16)	0 (16)
Grasshopper Sparrow	0 (1)	-	0 (8)	0 (9)
Le Conte's Sparrow	0 (4)	0 (4)	-	0 (8)
Vesper Sparrow	-	-	0 (10)	0 (10)
Sedge Wren	0 (6)	0 (1)	-	0 (7)
Lark Sparrow	-	-	0 (3)	0 (3)
Dickcissel	-	0 (1)	-	0 (1)
Total	3.0 (198)	2.7 (147)	12.8 (187)	6.4 (531)

Appendix 1: Avian density/ 100 ha on study plots in Minnesota State Parks in 2000.

Buffalo River State Park west (large hostile)

Clay-colored Sparrow	131.2
Savannah Sparrow	68.7
Sedge Wren	50.0
Common Yellowthroat	43.7
Le Conte's Sparrow	25.0
Song Sparrow	25.0
Tree Swallow	18.7
Bobolink	18.7
American Kestrel	6.2
Northern Harrier	6.2
Western Meadowlark	6.2

Buffalo River State Park east (small hostile)

Tree Swallow	66.7
Clay-colored Sparrow	66.7
American Goldfinch	66.7
Eastern Kingbird	33.3
House Wren	33.3
Yellow Warbler	33.3
Song Sparrow	33.3
Brown-headed Cowbird	33.3

Appendix 1: Avian density/ 100 ha on study plots on properties of The Nature Conservancy in 2000.

Blazing Star (large neutral)

Le Conte's Sparrow	118.7
Savannah Sparrow	93.7
Tree Swallow	56.2
Clay-colored Sparrow	37.5
Grasshopper Sparrow	37.5
Bobolink	37.5
Common Yellowthroat	25.0
Marbled Godwit	6.2
Upland Sandpiper	6.2
Northern Harrier	6.2
Short-eared Owl	6.2
Barn Swallow	6.2
Western Meadowlark	6.2
Brown-headed Cowbird	6.2
Red-winged Blackbird	6.2
American Goldfinch	6.2

Bluestem LN (large neutral)

Clay-colored Sparrow	80.0
Savannah Sparrow	53.3
Le Conte's Sparrow	46.7
Bobolink	46.7
Common Yellowthroat	40.0
Grasshopper Sparrow	40.0
Yellow Warbler	26.7
Brown-headed Cowbird	20.0
American Goldfinch	13.3
Eastern Kingbird	6.7
Alder Flycatcher	6.7
Barn Swallow	6.7
Gray Catbird	6.7
Sedge Wren	6.7
Song Sparrow	6.7
Western Meadowlark	6.7

Bluestem LH (large hostile)

Clay-colored Sparrow	59.0
Savannah Sparrow	32.8
Le Conte's Sparrow	19.7
Song Sparrow	13.1
Brown-headed Cowbird	13.1
Northern Harrier	6.6
Gray Catbird	6.6
Common Yellowthroat	6.6

Margherita (large neutral)

Savannah Sparrow	191.7
Le Conte's Sparrow	141.7
Bobolink	50.0
Tree Swallow	25.0
Sedge Wren	25.0
Red-winged Blackbird	25.0
Western Meadowlark	16.7
Mallard	8.3
Marbled Godwit	8.3
Northern Harrier	8.3
Greater Prairie-Chicken	8.3
Grasshopper Sparrow	8.3
Brown-headed Cowbird	8.3

Pankratz LH (large hostile)

Le Conte's Sparrow	113.3
Savannah Sparrow	80.0
Sedge Wren	53.3
Clay-colored Sparrow	46.7
Bobolink	33.3
Common Yellowthroat	13.3
Common Snipe	6.7
Upland Sandpiper	6.7
Eastern Kingbird	6.7
Gray Catbird	6.7
Brown-headed Cowbird	6.7

Pankratz N (large neutral)

Savannah Sparrow	187.5
Le Conte's Sparrow	143.7
Bobolink	68.7
Sedge Wren	50.0
Red-winged Blackbird	12.5
Common Snipe	6.2
Upland Sandpiper	6.2
Short-eared Owl	6.2
Barn Swallow	6.2
Western Meadowlark	6.2

Pankratz S (large neutral)

Savannah Sparrow	168.7
Le Conte's Sparrow	168.7
Sedge Wren	137.5
Bobolink	50.0
Clay-colored Sparrow	25.0
Red-winged Blackbird	25.0
Mallard	6.2
Common Snipe	6.2
Greater Prairie-Chicken	6.2
Nelson's Sharp-tailed Sparrow	6.2
Western Meadowlark	6.2

Pembina Trail (large neutral)

Savannah Sparrow	162.5
Le Conte's Sparrow	75.0
Clay-colored Sparrow	43.7
Bobolink	37.5
Greater Prairie-Chicken	31.2
Sedge Wren	18.7
Western Meadowlark	18.7

Brown-headed Cowbird	18.7
Common Yellowthroat	12.5
Brewer's Blackbird	12.5
Killdeer	6.2
Common Snipe	6.2
Eastern Kingbird	6.2
Grasshopper Sparrow	6.2
Red-winged Blackbird	6.2

Rice-Elliott (large neutral)

Savannah Sparrow	181.2
Le Conte's Sparrow	68.7
Red-winged Blackbird	50.0
Bobolink	31.2
Sedge Wren	25.0
Mallard	6.2
Common Snipe	6.2
Greater Prairie-Chicken	6.2
Northern Harrier	6.2
American Kestrel	6.2
Short-eared Owl	6.2
Grasshopper Sparrow	6.2
Western Meadowlark	6.2

Zimmerman (small neutral)

Savannah Sparrow	185.7
Red-winged Blackbird	114.3
Bobolink	100.0
Le Conte's Sparrow	57.1
Clay-colored Sparrow	57.1
Marsh Wren	42.9
Common Grackle	42.9
Eastern Kingbird	28.6
Sora	14.3
Tree Swallow	14.3
Yellow Warbler	14.3
Swamp Sparrow	14.3
Brown-headed Cowbird	14.3
American Goldfinch	14.3

Appendix 1: Avian density/ 100 ha on study plots in Scientific and Natural Areas in 2000.

Bicentennial (large neutral)

Grasshopper Sparrow	93.7
Clay-colored Sparrow	75.0
Bobolink	37.5
Common Yellowthroat	18.7
Tree Swallow	12.5
Yellow Warbler	12.5
Western Meadowlark	12.5
Marbled Godwit	6.2
Upland Sandpiper	6.2
Eastern Kingbird	6.2
Barn Swallow	6.2
American Robin	6.2
Le Conte's Sparrow	6.2
Brown-headed Cowbird	6.2

Foxboro (small neutral)

Savannah Sparrow	266.7
Le Conte's Sparrow	66.7
Clay-colored Sparrow	33.3
Marbled Godwit	16.7
Common Snipe	16.7
Eastern Kingbird	16.7
Common Yellowthroat	16.7
Red-winged Blackbird	16.7

Appendix 1: Avian density/100 ha on study plots in Wildlife Management Areas in 2000.

Burnham (large hostile)

Le Conte's Sparrow	125.0
Savannah Sparrow	108.3
Bobolink	25.0
Sedge Wren	16.7
Yellow Warbler	16.7
Eastern Kingbird	8.3
Common Yellowthroat	8.3
Clay-colored Sparrow	8.3

Chicog LH (small hostile)

Clay-colored Sparrow	120.0
Savannah Sparrow	72.0
Bobolink	24.0
Marsh Wren	16.0
Red-winged Blackbird	16.0
Eastern Kingbird	8.0
Sedge Wren	8.0
Common Yellowthroat	8.0
Vesper Sparrow	8.0

Chicog SH (small hostile)

Savannah Sparrow	53.3
Common Snipe	26.7
Common Yellowthroat	26.7
Le Conte's Sparrow	26.7
Clay-colored Sparrow	26.7
American Goldfinch	26.7

Dugdale (small neutral)

Savannah Sparrow	133.3
Bobolink	66.7
Brewer's Blackbird	50.0
Red-winged Blackbird	33.3
Northern Harrier	16.7

Shypoke (small neutral)

Savannah Sparrow	116.7
Sedge Wren	66.7
Le Conte's Sparrow	33.3
Swamp Sparrow	33.3
Northern Flicker	16.7
Common Yellowthroat	16.7
Clay-colored Sparrow	16.7
Western Meadowlark	16.7

Mentor NW (small hostile)

Savannah Sparrow	100.0
Le Conte's Sparrow	100.0
Clay-colored Sparrow	66.7
Red-tailed Hawk	33.3

Mentor SE (small hostile)

Le Conte's Sparrow	137.1
Savannah Sparrow	80.0
Clay-colored Sparrow	45.7
Sedge Wren	34.3
Willow Flycatcher	11.4

Mentor SN (small neutral)

Savannah Sparrow	133.3
Le Conte's Sparrow	122.2
Sedge Wren	55.6
Clay-colored Sparrow	33.3
Bobolink	22.2
Brown-headed Cowbird	22.2
Common Snipe	11.1
Upland Sandpiper	11.1
Western Meadowlark	11.1
Red-winged Blackbird	11.1

Ulen (small hostile)

Le Conte's Sparrow	90.9
Savannah Sparrow	72.7
Clay-colored Sparrow	72.7
Bobolink	72.7
Barn Swallow	18.2
Tree Swallow	18.2
Common Yellowthroat	18.2
Yellow Warbler	18.2
Song Sparrow	18.2

Tilden (small hostile)

Clay-colored Sparrow	72.7
Grasshopper Sparrow	60.6
Savannah Sparrow	60.6
American Goldfinch	24.2
Barn Swallow	12.1
Bobolink	12.1
Western Meadowlark	12.1

Tympanuchus (large neutral)

Savannah Sparrow	175.0
Clay-colored Sparrow	68.7
Bobolink	50.0
Le Conte's Sparrow	43.7
Sedge Wren	18.7
Common Yellowthroat	18.7
Common Snipe	12.5
Western Meadowlark	12.5
Marbled Godwit	6.2
Upland Sandpiper	6.2
Grasshopper Sparrow	6.2

Appendix 1: Avian density/ 100 ha on study plots in Waterfowl Production Areas in 2000.

Eide (small neutral)

Mallard	100.0
Bobolink	100.0
Common Yellowthroat	50.0
Savannah Sparrow	50.0
Clay-colored Sparrow	50.0
Red-winged Blackbird	50.0

Flickertail (large hostile)

Clay-colored Sparrow	108.3
Le Conte's Sparrow	33.3
Common Yellowthroat	25.0
Savannah Sparrow	16.7
Song Sparrow	16.7
Bobolink	16.7
Barn Swallow	8.3
Eastern Kingbird	8.3
House Wren	8.3
Yellow Warbler	8.3

Hoykens (small neutral)

Savannah Sparrow	121.7
Clay-colored Sparrow	121.7
Red-winged Blackbird	121.7
Bobolink	69.6
Common Yellowthroat	52.2
Song Sparrow	34.8
Wood Duck	17.4
Killdeer	17.4
Common Snipe	17.4
Tree Swallow	17.4
Marsh Wren	17.4
Yellow Warbler	17.4
Le Conte's Sparrow	17.4
Swamp Sparrow	17.4
Brown-headed Cowbird	17.4

Fuglie (large hostile)

Bobolink	68.7
Bank Swallow	31.2
Sedge Wren	18.7
Savannah Sparrow	18.7
Eastern Kingbird	6.2
Clay-colored Sparrow	6.2

Private (small hostile)

Clay-colored Sparrow	88.9
Le Conte's Sparrow	66.7
Savannah Sparrow	66.7
Sedge Wren	44.4
Bobolink	44.4
Eastern Kingbird	22.2
Common Yellowthroat	22.2
American Goldfinch	22.2

Sagebraaten (small neutral)

Red-winged Blackbird	133.3
Bobolink	100.0
Common Yellowthroat	33.3
Marsh Wren	33.3
Song Sparrow	33.3
Swamp Sparrow	33.3
American Goldfinch	33.3

Spring Creek (small neutral)

Savannah Sparrow	94.1
Bobolink	58.8
Red-winged Blackbird	58.8
Clay-colored Sparrow	47.0
Common Yellowthroat	23.5
Sedge Wren	23.5
Le Conte's Sparrow	23.5
Barn Swallow	11.8
Marsh Wren	11.8
Western Meadowlark	11.8

Appendix 1: Avian density/ 100 ha on study plots in State Game Refuges in 2000.

Refuge (small neutral)

Savannah Sparrow	95.2
Grasshopper Sparrow	66.7
Clay-colored Sparrow	47.6
Le Conte's Sparrow	19.0
Mallard	9.5
Northern Harrier	9.5
Short-eared Owl	9.5
Sedge Wren	9.5
Song Sparrow	9.5
Bobolink	9.5
Western Meadowlark	9.5
Red-winged Blackbird	9.5
Brown-headed Cowbird	9.5
American Goldfinch	9.5

Appendix 1: Avian density/ 100 ha on study plots in Sheyenne National Grassland in 2000.

Camp (small hostile)

Grasshopper Sparrow	124.1
Clay-colored Sparrow	96.6
Bobolink	55.2
Eastern Kingbird	27.6
Common Yellowthroat	27.6
Field Sparrow	27.6
Vesper Sparrow	13.8
Western Meadowlark	13.8

<u>Pileated (small hostile)</u>	
Clay-colored Sparrow	142.8
Grasshopper Sparrow	71.4
Vesper Sparrow	57.1
American Crow	28.6
American Goldfinch	28.6
Eastern Kingbird	14.3
Field Sparrow	14.3
Western Meadowlark	14.3

<u>Surprise (small hostile)</u>	
Bobolink	133.3

<u>Highway (large neutral)</u>	
Savannah Sparrow	150.0
Grasshopper Sparrow	62.5
Le Conte's Sparrow	37.5
Western Meadowlark	31.2
Brown-headed Cowbird	31.2
Yellow-headed Blackbird	25.0
Blue-winged Teal	18.7
Marbled Godwit	12.5
Eastern Kingbird	12.5
Bobolink	12.5
Red-winged Blackbird	12.5
Wilson's Phalarope	6.2
Killdeer	6.2
Common Snipe	6.2
Morning Dove	6.2
Marsh Wren	6.2

<u>North (large neutral)</u>	
Bobolink	118.7
Grasshopper Sparrow	87.5
Savannah Sparrow	81.2
Barn Swallow	37.5
Western Meadowlark	43.7
Red-winged Blackbird	31.2
Le Conte's Sparrow	18.7
Mallard	6.2
Marbled Godwit	6.2
Upland Sandpiper	6.2

Plum (large neutral)

Savannah Sparrow	137.5
Grasshopper Sparrow	131.2
Western Meadowlark	31.2
Le Conte's Sparrow	25.0
Brown-headed Cowbird	18.7
Marbled Godwit	6.2
Barn Swallow	6.2
Eastern Kingbird	6.2
Bobolink	6.2

Southeast (large neutral)

Red-winged Blackbird	100.0
Savannah Sparrow	81.2
Le Conte's Sparrow	62.5
Bobolink	56.2
Clay-colored Sparrow	43.7
Wilson's Phalarope	31.2
Western Meadowlark	31.2
Sedge Wren	25.0
Barn Swallow	12.5
Common Yellowthroat	12.5
Northern Shoveler	6.2
Greater Prairie-Chicken	6.2
Common Grackle	6.2

Shrike (small hostile)

Clay-colored Sparrow	109.1
Barn Swallow	72.7
Eastern Kingbird	36.4
Warbling Vireo	36.4
Western Meadowlark	36.4
Brown-headed Cowbird	36.4
American Goldfinch	36.4

Eagle (large hostile)	
<hr/>	
Savannah Sparrow	100.0
Clay-colored Sparrow	85.7
Red-winged Blackbird	78.6
Bobolink	71.4
Grasshopper Sparrow	64.3
Le Conte's Sparrow	35.7
Common Grackle	35.7
Western Meadowlark	35.7
Brown-headed Cowbird	35.7
Common Yellowthroat	21.4
American Goldfinch	21.4
Common Snipe	14.3
Eastern Kingbird	14.3
Mallard	7.1
Northern Shoveler	7.1
Killdeer	7.1
Barn Swallow	7.1
Gray Catbird	7.1
Warbling Vireo	7.1
Yellow Warbler	7.1
Northern Oriole	7.1
Song Sparrow	7.1
Vesper Sparrow	7.1
<hr/>	

Hammock (large hostile)	
<hr/>	
Bobolink	68.9
Grasshopper Sparrow	55.2
Savannah Sparrow	27.6
Red-winged Blackbird	27.6
Tree Swallow	20.7
Upland Sandpiper	13.8
Common Snipe	13.8
Sedge Wren	13.8
Le Conte's Sparrow	13.8
Western Meadowlark	13.8
Greater Prairie-Chicken	6.9
Eastern Kingbird	6.9
Clay-colored Sparrow	6.9
Brown-headed Cowbird	6.9
<hr/>	

Savannah (large hostile)

Bobolink	106.2
Red-winged Blackbird	62.5
Savannah Sparrow	56.2
Grasshopper Sparrow	56.2
Brown-headed Cowbird	31.2
Blue-winged Teal	25.0
Wilson's Phalarope	18.7
Western Meadowlark	18.7
Green-winged Teal	12.5
Le Conte's Sparrow	12.5
Mallard	6.2
Northern Pintail	6.2
Killdeer	6.2
Barn Swallow	6.2
Clay-colored Sparrow	6.2
Common Grackle	6.2

Appendix 2: Number of nests found, number of nests successful, and number of nests parasitized by Brown-headed Cowbirds in Waterfowl Production areas in 2000.

Hoykens (small neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Mallard	3	3	0
Blue-winged Teal	2	1	0
Redhead	1	0	0
Marsh Wren	1	1	0
Savannah Sparrow	2	1	0
Song Sparrow	1	?	0
Swamp Sparrow	1	1	0
Bobolink	2	0	0
Red-winged Blackbird	1	1	0

Private (small hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Eastern Kingbird	1	?	0
Sedge Wren	1	1	0
Clay-colored Sparrow	3	3	0
Song Sparrow	1	1	0

Refuge (small neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Mallard	1	0	0
Savannah Sparrow	4	1	0
Clay-colored Sparrow	4	1	0
Western Meadowlark	2	0	1
Dickcissel	1	1	0
Red-winged Blackbird	1	0	0
American Goldfinch	2	?	0

Spring Creek (small neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Blue-winged Teal	2	2	0
Savannah Sparrow	3	2	0
Clay-colored Sparrow	3	0	0
Bobolink	3	1	0
Red-winged Blackbird	2	2	0

Appendix 2: Number of nests found, number of nests successful, and number of nests parasitized by Brown-headed Cowbirds in study plots on properties of The Nature Conservancy in 2000.

BluestemLH (large hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Yellow Warbler	1	0	0
Le Conte's Sparrow	1	0	0
Song Sparrow	2	2	0
Clay-colored Sparrow	17	2	1

BluestemLN (large neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Mallard	2	0	0
Upland Sandpiper	1	?	0
Northern Harrier	1	0	0
Gray Catbird	2	0	0
Yellow Warbler	2	0	0
Le Conte's Sparrow	2	0	0
Savannah Sparrow	10	3	1
Song Sparrow	1	0	1
Clay-colored Sparrow	12	6	0
Bobolink	3	1	0

Pankratz east (large hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Savannah Sparrow	3	2	0
Clay-colored Sparrow	6	6	0
Bobolink	2	0	0

Pankratz north (large neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Upland Sandpiper	1	0	0
Greater Prairie-Chicken	1	0	0
Sedge Wren	1	0	0
Le Conte's Sparrow	1	?	0
Savannah Sparrow	26	13	0
Bobolink	5	0	0
Red-winged Blackbird	1	0	0

Pankratz south (large neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Mallard	2	2	0
Marbled Godwit	1	0	0
Sedge Wren	4	4	0
Le Conte's Sparrow	1	?	0
Savannah Sparrow	28	13	1
Clay-colored Sparrow	1	0	1
Bobolink	3	0	0
Red-winged Blackbird	3	1	1

Appendix 2: Number of nests found, number of nests successful, and number of nests parasitized by Brown-headed Cowbirds in study plots in Scientific and Natural Areas in 2000.

Bicentennial (large neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Upland Sandpiper	1	1	0
Greater Prairie-Chicken	2	0	0
Eastern Kingbird	1	1	0
Clay-colored Sparrow	16	5	0
Bobolink	3	1	0
American Goldfinch	1	?	0

Foxboro (small neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Savannah Sparrow	25	11	0
Clay-colored Sparrow	6	0	0

Appendix 2: Number of nests found, number of nests successful, and number of nests parasitized by Brown-headed Cowbirds in study plots in Minnesota State Parks in 2000.

Buffalo (large hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Tree Swallow	1	1	0
Common Yellowthroat	3	0	0
Le Conte's Sparrow	1	0	0
Savannah Sparrow	3	1	0
Song Sparrow	2	1	0
Clay-colored Sparrow	17	3	0
Bobolink	1	1	0
American Goldfinch	2	?	0

Appendix 2: Number of nests found, number of nests successful, and number of nests parasitized by Brown-headed Cowbirds in Wildlife Management Areas in 2000.

Burnham (large hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Le Conte's Sparrow	2	1	0
Savannah Sparrow	14	2	0
Clay-colored Sparrow	2	0	0

Dugdale (small neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Upland Sandpiper	1	0	0
Savannah Sparrow	3	2	0
Bobolink	2	0	0
Red-winged Blackbird	2	0	0
Brewer's Blackbird	9	2	1

Mentor (small hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Le Conte's Sparrow	1	0	0
Savannah Sparrow	7	3	1
Clay-colored Sparrow	3	2	0

Mentor (small neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Upland Sandpiper	4	1	0
Greater Prairie-Chicken	1	0	0
Sedge Wren	1	0	0
Savannah Sparrow	9	5	0
Clay-colored Sparrow	2	1	0
Bobolink	2		0
Western Meadowlark	1	0	0

Shypoke (small neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Savannah Sparrow	8	5	0
Western Meadowlark	1	1	0
Red-winged Blackbird	1	0	0

Tilden (small hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Grasshopper Sparrow	1	1	0
Savannah Sparrow	3	0	0
Clay-colored Sparrow	3	1	0
Bobolink	1	1	0

Tympanuchus (large neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Greater Prairie-Chicken	1	1	0
Mallard	1	0	0
Northern Harrier	1	0	0
Savannah Sparrow	15	8	1
Clay-colored Sparrow	4	0	0
Bobolink	5	0	0

Ulen (small hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Yellow Warbler	3	2	0
Savannah Sparrow	3	1	0
Song Sparrow	3	3	0
Clay-colored Sparrow	13	4	0
Bobolink	6	1	1

Appendix 2: Number of nests found, number of nests successful, and number of nests parasitized by Brown-headed Cowbirds on study plots in Sheyenne National Grassland in 2000.

Camp (small hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Morning Dove	1	0	0
Eastern Kingbird	1	0	0
Vesper Sparrow	2	1	0
Field Sparrow	6	4	0
Lark Sparrow	1	?	0
Clay-colored Sparrow	18	5	0
Bobolink	4	0	0

Eagle Point (large hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Blue-winged Teal	1	0	0
Northern Shoveler	1	0	0
Morning Dove	4	0	0
Eastern Kingbird	3	1	0
Common Yellowthroat	1	0	0
Yellow Warbler	1	0	0
Grasshopper Sparrow	5	3	0
Vesper Sparrow	1	0	0
Savannah Sparrow	7	2	0
Clay-colored Sparrow	22	6	5
Bobolink	3	0	0
Western Meadowlark	2	0	0
Red-winged Blackbird	6	3	2

Highway (large neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Blue-winged Teal	5	1	0
Green-winged Teal	1	0	0
Northern Shoveler	1	0	0
Marbled Godwit	1	0	0
Wilson's Phalarope	4	0	0
Morning Dove	1	1	0
Savannah Sparrow	11	4	6
Western Meadowlark	4	1	1
Red-winged Blackbird	3	2	1
Yellow-headed Blackbird	1	0	0

Savannah (large hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Blue-winged Teal	1	1	0
Green-winged Teal	1	1	0
Northern Shoveler	1	0	0
Common Snipe	1	0	0
Morning Dove	2	0	0
Grasshopper Sparrow	1	1	0
Vesper Sparrow	1	0	0
Savannah Sparrow	4	3	0
Clay-colored Sparrow	4	0	0
Bobolink	6	0	0
Western Meadowlark	3	1	0
Red-winged Blackbird	1	1	0
American Goldfinch	1	?	0

North (large neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Mallard	4	2	0
Blue-winged Teal	3	0	0
Savannah Sparrow	2	0	0
Bobolink	9	8	0
Western Meadowlark	2	1	0
Red-winged Blackbird	3	2	0

Plum (large neutral)

Species	No. found	No. successful	No. parasitized by cowbirds
Mallard	1	0	0
Upland Sandpiper	1	0	0
Grasshopper Sparrow	2	0	0
Savannah Sparrow	20	7	7
Western Meadowlark	6	0	1

Pileated (small hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Clay-colored Sparrow	16	6	0
Vesper Sparrow	6	4	0
Lark Sparrow	2	1	0
Field Sparrow	10	5	0
Chipping Sparrow	1	1	0

Shrike (small hostile)

Species	No. found	No. successful	No. parasitized by cowbirds
Morning Dove	1	0	0
Clay-colored Sparrow	2	1	1