

Plants and Lichens as Indicators of Atmospheric Deposition

USGS is studying the pathways used by atmospheric contaminants in National Parks using plants and lichens.

The natural world is being continuously exposed to atmospheric emissions from man's activities. Even remote, supposedly pristine areas like Isle Royale National Park in Lake Superior and areas of Alaska are exposed. This is due to the magnitude of emissions from industries, urban areas, and power generation facilities, and the ability of the atmosphere to transport the emissions great distances. Elements such as mercury and lead are found in ecosystems in national parks throughout our country. These are toxic to many organisms as they are passed up food chains. One USGS scientist is conducting long-term studies of the geographic extent and degree of contamination of these

elements and several others in our National Parks.

In the terrestrial world, deposition of air pollutants occurs to foliage of plants and the soil surface. Rooted plants, therefore, may take up elements through leaves directly, or through roots in the soil. Mobility of these elements inside plants is determined physiologically, and therefore analysis of plant parts elucidates the pathways of deposition. Arsenic, for example, is highly toxic, and is usually confined to roots. Analysis of soils is also needed to determine the element concentrations normally present. These studies also help determine if there are naturally high levels of certain elements present from geologic anomalies. For example, naturally high levels of mercury are found in some plants in Yellowstone National Park due to natural emissions from geothermal features.


Lichens are small, symbiotic plants that grow on trees, rocks and soils. They have no roots and are totally dependent on the atmosphere for their nutrition. They receive no sustenance from their hosts, just physical support. Thus, elevated concentrations of certain elements in lichens are a sure sign of atmospheric deposition. Studies of the presence and absence of sensitive



Hypogymnia physodes, one of the most common lichens in the world, is shown here growing on birch bark.

lichen species, and of element concentrations in their tissues are ways of discovering atmospheric impacts in an area. Some studies are performed along transects downwind of point sources (tall stacks, for example) to determine the amount and geographic distribution of fallout from the source. Transplants of lichens from pristine to polluted areas are done to determine the continued presence of atmospheric pollutants. Common species such as *Hypogymnia physodes*, that are found in many localities, are used for sampling element deposition because of their widespread distributions. Nutritional elements are commonly studied, in addition to elements known to be emitted anthropogenically, such as lead, mercury, cadmium and chromium.

Since 1982, studies like these have been conducted in 26 national parks. Almost 2000 lichen samples representing 49 species have been collected from 333 localities in these parks. These samples have been measured for 22 chemical elements, providing over 40,000 data points for analysis. The data are archived for use by other researchers and for comparisons over




Statistical analysis of arsenic deposits in wild rice (Zizania palustris) from northern Wisconsin shows the highest concentrations in the roots.

time. This research helps to document the effects of air pollution on park vegetation and assists park managers by alerting them to potential air pollution

problems. Work on additional parks (and continuation of efforts in currently sampled parks) will continue. About 2 parks are sampled each year.

For additional information, please contact:
 Dr. James P. Bennett
 U.S. Geological Survey
 National Wildlife Health Center
 6006 Schroeder Road, Madison WI 53711-6223
 (608) 262-5489 FAX (608) 265-2993
<http://www.nwhc.usgs.gov/>

National Parks Studied for Lichen Chemistry



<i>Acadia, ME</i>	26	<i>Kings Canyon, CA</i>	3
<i>Apostle Islands, WI</i>	17	<i>Olympic, WA</i>	1
<i>Chaco Culture, NM</i>	8	<i>Pictured Rocks, MI</i>	20
<i>Chiricahua, AZ</i>	7	<i>Redwood, CA</i>	2
<i>Cuyahoga Valley, OH</i>	23	<i>Saint Croix, MN & WI</i>	15
<i>Delaware Water Gap, PA</i>	25	<i>Saguaro, AZ</i>	6
<i>Effigy Mounds, IA</i>	16	<i>Sequoia, CA</i>	4
<i>Grand Portage, MN</i>	18	<i>Shenandoah, VA</i>	24
<i>George Washington Carver, MO</i>	12	<i>Sleeping Bear Dunes, MI</i>	21
<i>Homestead, NE</i>	10	<i>Theodore Roosevelt, ND</i>	9
<i>Hot Springs, AR</i>	13	<i>Voyageurs, MN</i>	14
<i>Indiana Dunes, IN</i>	22	<i>Wilson's Creek, MO</i>	11
<i>Isle Royale, MI</i>	19	<i>Yellowstone, WY</i>	5

