

Southwestern Willow Flycatcher Breeding Site and Territory Summary – 2000

Prepared by:

Mark K. Sogge, U.S. Geological Survey, Forest and Rangeland Ecosystem Science Center, Colorado Plateau Field Station, Flagstaff, AZ

Susan J. Sferra, U.S. Bureau of Reclamation, Phoenix Area Office, Phoenix, AZ

Tracy McCarthy, Arizona Game and Fish Department - Nongame Branch, Phoenix, AZ

Sartor O. Williams III, New Mexico Department of Game and Fish, Santa Fe, NM

Barbara E. Kus, U.S. Geological Survey, Western Ecological Research Center, San Diego Field Station, San Diego, CA

Prepared for:

The Southwestern Willow Flycatcher Recovery Team, U.S. Fish and Wildlife Service, Region 2, Albuquerque, NM

November 2001

Southwestern Willow Flycatcher Breeding Site and Territory Summary - 2000

Table of Contents

Introduction	2
Changes in number of known territories over time	5
Recency of survey data	6
Population sizes of breeding sites	7
Distribution of territories by state	8
Distribution of territories by drainage	9
Distribution of territories by Recovery Unit	10
Elevation range of breeding territories	11
Use of native and exotic habitats	12
Dominant tree species at breeding sites	13
Administration/management of sites and territories	14
Summary	15
Literature cited / data sources	16
Acknowledgements	19

Southwestern Willow Flycatcher Breeding Site and Territory Summary - 2000

Introduction

The Southwestern Willow Flycatcher (*Empidonax traillii extimus*) is an endangered bird that is known to breed only in dense riparian habitats in six southwestern states (southern California, extreme southern Nevada, southern Utah and Colorado, Arizona, and New Mexico). Since 1993, hundreds of Willow Flycatcher surveys have been conducted each year, and many new flycatcher breeding sites located. This document arose from efforts to synthesize information on all known Southwestern Willow Flycatcher breeding sites, primarily as a tool for the U.S. Fish and Wildlife Service's Southwestern Willow Flycatcher Recovery Team. Established in 1998, this team depends on access to all available current information in order to effectively plan for the conservation and recovery of the flycatcher.

This rangewide data synthesis was designed to meet these objectives:

- 1 – identify all known Southwestern Willow Flycatcher breeding sites, and
- 2 – assemble data on population size, location, habitat, and other information for all breeding sites, for as many years as possible, from 1993 through 2000.

This report provides data summaries in terms of the number of flycatcher sites and the number of territories. When interpreting and using this information, the following must be kept in mind:

A site is defined as a location where one or more Willow Flycatchers establish a territory in which they attempt to breed. Sites with unpaired territorial males are considered breeding sites even if no nesting attempts were documented. A site is often a discrete patch of habitat; however, there is no standardized definition for site and its use varies among states. For example, five occupied habitat patches along a 10 km stretch of river might be considered as five different sites in one state, but as only a single site in another state. This makes comparison of information based on “site” problematic. For this report, we deferred to the statewide summary documents, or to local managers and researchers, when delineating a site for inclusion in the database. Due to differences in site definitions, one should not evaluate the relative importance of a geographic region (drainage, watershed, state, etc.) based simply on the number of flycatcher sites.

A territory is an exclusive defended area within a breeding site. Although detailed monitoring studies have identified unpaired territorial males and/or polygynous males at some flycatcher breeding sites, for purposes of this report a territory is roughly equivalent to a pair of flycatchers. The concept of territory is more similar between states and among different investigators, so this is a more “robust” unit to use for summaries and comparisons.

For each breeding site, we referred to reports or spoke directly with researchers and managers to gather information such as management entity/agency, location (state, drainage, elevation), gross habitat type (native, exotic, or mixed; dominant tree species), and flycatcher population size (number of territories).

Gathering and synthesizing the information on more than 200 breeding sites was at times challenging because of inconsistencies in practices of collecting and reporting flycatcher information across states. Rangelwide, annual survey reporting requirements are not standardized, and the nature and degree of readily available information varied widely. Most states and USFWS regions require standard data sheets be submitted each year, and produce detailed statewide summary reports; these resources were tremendously helpful in producing this report. The task was more challenging for areas such as California, where synthesis of annual Willow Flycatcher survey data is minimal. Furthermore, the USFWS in southern California does not require that surveyors submit the standardized survey forms, further complicating data gathering. The lack of standardized, annual state-based synthesis and reporting is the most immediate obstacle to rangelwide synthesis of data. We sincerely thank the many people who generously provided information from the sites they were surveying and monitoring (see following sections listing data sources and contacts, and acknowledgements).

This report includes all flycatcher breeding sites reported between 1993 and 2000. The statistics included herein are based on survey data from the most recent year during which surveys were conducted, whether flycatchers were detected or not. Therefore, 56 sites that had no flycatchers in the most recent survey year (as judged by the agencies consolidating statewide survey data) are still included in the site tallies if they had resident flycatchers during one or more years since 1993. This report does not include data from sites where only migrant Willow Flycatchers were detected.

Every effort was made to locate and include all survey information for every known Southwestern Willow Flycatcher breeding site. However, there may be some extant sites that have not yet been reported and are therefore not included herein. Hopefully, the preparation and dissemination of this report will prompt additional and more comprehensive reporting, such that future annual rangelwide summaries become more complete with each iteration.

Additional Considerations in Using and Interpreting the Data in this Report: We used data from a wide variety of sources, and the amount of information and level of detail varied greatly among sites. Because survey methodology and effort varied among sites and/or between years, these summary data should be interpreted and used in context. Following is a discussion of cautions to consider when using these data.

Subspecies status of each site: The Willow Flycatcher sites entered into this database all fall within the geographic range of the southwestern subspecies (*E.t. extimus*), as defined by Unitt (1987), Browning (1993) and Sogge et al. (1997). Recent studies of flycatcher genetics (e.g., Paxton 2000) and song patterns (e.g., Sedgwick 2001) support a more southern range boundary for *E.t. extimus* than was used for the 1999 summary (Sogge et al. 2000). Thus, the Beaver Creek and Clear Creek sites (in the Dolores National Forest, CO) were removed from consideration as *extimus*. Future research may provide more insight into subspecies range boundaries; therefore, additional sites may eventually be removed from management as *extimus*, and/or new geographic areas and sites could be added. This should be considered when producing updates in future years, and when making rangewide comparisons among years.


Population estimates: Population estimates are just that – **estimates**. Their accuracy and precision vary with survey effort, surveyor experience, habitat density, and even background noise levels. The population estimates often represent the minimum number of flycatchers present; i.e., if surveyors suspected 12 to 14 flycatchers, the lower (more conservative) number was used. Therefore, although estimates may be very accurate for some intensively surveyed sites, the overall statistics presented in this report should be recognized as approximate.

DATA SUMMARIES

Changes in the number of known territories over time

Since 1993, extensive survey effort in Arizona, California, Colorado, New Mexico and Utah has greatly increased the number of known breeding sites and breeding territories. From a 1993 estimate of roughly 30 sites and 111 territories, we now have data for 933 territories, located among 209 sites (Figure 1). This increase should NOT be interpreted entirely as a Southwestern Willow Flycatcher population increase. Rather, it is to a great extent a function of increased survey effort over time. Although population increases and decreases undoubtedly occur at some sites, movements of birds among sites and lack of standardized survey effort/reporting make it difficult to separate population trends from variances in survey effort. Determination of trends (positive or negative) can be made in only a few cases, and original data sources (e.g., reports, survey data sheets, etc.) must be consulted when trying to elucidate population trends.

FIGURE 1



Recency of survey data

The information used in this report is based on the most recent available survey data for each site. Although there were a few sites for which we could not find survey data more recent than 1995 or earlier, 1999 and/or 2000 data were available for 84% of sites (accounting for 91% of territories). Thus, the information used for most of the statistics reported herein is quite recent.

Table 1. Most recent year of survey data for sites and territories included in this report.

Year	# Sites	% Total Sites (n=209)	# Territories	% Total Territories (n=933)
1993	3	1	6	0.6
1994	1	0.5	0	0
1995	4	2	5	0.5
1996	4	2	9	1
1997	9	4	44	5
1998	13	6	17	2
1999	19	9	27	3
2000	156	75	825	88


Population sizes of breeding sites

Most southwestern willow flycatcher breeding sites are small, both in terms of population size (hosting five or fewer territories: Figure 2) and habitat patch size. Such small sites are theoretically more susceptible to extirpation, and there is evidence to support this is the case. Willow flycatchers have disappeared from 56 of the 209 sites tracked since 1993. All but two of these extirpated sites were composed of five or fewer territories. The two exceptions – Colorado River inflow to Lake Mead, and PZ Ranch on the San Pedro River – were larger sites where habitat was destroyed by flooding and fire, respectively.

Not all birds at these extirpated sites necessarily died – some may have moved elsewhere. We know this is the case for banded flycatchers which moved from the Verde River Tuzigoot Bridge and PZ Ranch to other sites (Paxton and Sogge 1996, Paxton et al. 1997, Netter et al. 1998).

If we look again at the size distribution of breeding sites and exclude the extirpated sites, the picture remains much the same - the vast majority of sites (73%) have five or fewer territories. Because most of the 56 extirpated sites had very small populations (usually only one or two territories), their loss does not greatly affect the overall rangewide territory estimates, nor many of the territory statistics that we report herein.

Figure 2



Distribution of territories by state

Arizona, New Mexico, and California account for the greatest number of known Southwestern Willow Flycatcher sites and territories (Table 2). Nevada, Colorado, and Utah account for only 11% of territories, primarily because they have few known Willow Flycatcher breeding sites occurring far enough south to fall within the range of *E.t. extimus*. Texas is absent from this table because there were no recent survey data or other records to shed light on current status and distribution within the state. We believe this is an unfortunate data gap and hope that coordinated survey work is soon initiated within southwestern Texas.

Table 2. The number of Southwestern Willow Flycatcher breeding sites and territories by state.

State	# Sites	% of Total Sites (n=209)	# Territories	% of Total Territories (n=933)
AZ	91	44	341	37
CA	65	31	224	24
CO	5	2	37	4
NM	32	15	263	28
NV	10	5	56	6
UT	6	3	12	1
TOTAL	209		933	

Distribution of territories by drainage

More flycatcher territories are found along the Gila River than any other major drainage (Table 3); the largest known population (in the Cliff-Gila Valley, NM) contributes many of the territories within this drainage. Elsewhere in New Mexico, and in southwest Colorado, most territories are along the Rio Grande. The primary flycatcher drainages in California are the Kern, Santa Ana, Santa Margarita, and San Luis Rey rivers.

Table 3. The number of Southwestern Willow Flycatcher breeding sites and territories by major river drainage.

DRAINAGE	# Sites	% of Total Sites (n=209)	# Territories	% of Total Territories (n=933)
Colorado River	37	18	34	4
Gila River	34	16	235	25
Kern River	2	1	27	3
Rio Grande	18	9	109	12
San Luis Rey River	7	3	56	6
San Pedro River	13	6	69	7
Salt River	5	2	87	9
Santa Ana River	16	8	31	3
Santa Margarita River	2	1	19	2
Tonto Creek	1	0.5	28	3
Virgin River	6	3	33	4
All others	68	32	205	22

Distribution of territories by Recovery Unit

We tallied the number of breeding sites and territories by Recovery Unit and Management Unit (Table 4), as defined in the draft Southwestern Willow Flycatcher Recovery Plan. Note that in some Management Units, the number of territories is **less than** the number of sites; this occurs where Management Units include primarily small sites, one or more of which no longer contain territorial flycatchers as of the most recent survey (e.g., “extirpated” sites).

Table 4. The currently known number of flycatcher breeding sites and territories (as of 2000 data), by Recovery Unit and Management Unit.

Recovery Unit	Management Unit	# of Sites	# of Territories
Basin and Mojave	Owens	5	16
	Kern	2	27
	Amargosa	2	6
	Mojave	5	12
	Salton	1	3
	TOTAL	15	64
Coastal California	Santa Ynez	3	34
	Santa Clara	3	3
	Santa Ana	18	32
	San Diego	20	93
	TOTAL	44	162
Gila	Verde	4	5
	Hassayampa - Agua Fria	1	0
	Salt – Tonto	6	115
	San Francisco	2	4
	Upper Gila	16	184
	Gila – San Pedro	30	120
	TOTAL	59	428
Lower Colorado	Pahranagat	5	28
	Virgin	5	32
	Little Colorado	4	7
	Middle Colorado	18	12
	Hoover - Parker	6	19
	Bill Williams	8	41
	Parker - Mexico	14	3
	TOTAL	60	142
Rio Grande	San Luis Valley	2	34
	Upper Rio Grande	13	45
	Middle Rio Grande	6	47
	Lower Rio Grande	2	6
	TOTAL	23	132
Upper Colorado River	Sevier	4	2
	Dolores	<i>Removed from</i>	<i>extimus range</i>
	San Juan	4	3
	TOTAL	8	5
GRAND TOTAL		209	933

Elevational range of breeding territories

As might be expected of a species that ranges over such a wide geographic area, the Southwestern Willow Flycatcher is distributed over a wide elevational range. The majority of sites occur between 0 and 1000 m elevation (Figure 3a). Most territories are found between 0 and 1600 m, with “spikes” at 601-800 m (the Gila/San Pedro River confluence area in AZ) and 1401-1600 m (the Cliff-Gila Valley in NM). Although relatively few territories are known to occur above 2000 m elevation, Willow Flycatchers breed at 3 sites that are above 2500 m.

Figure 3.

Figure 3a. The percentage of flycatcher breeding sites located at different elevations (200 = 0 - 200 m, 400 = 201 – 400 m, etc.).





Figure 3b. The percentage of flycatcher territories occurring at differing elevations (200 = 0 - 200 m, 400 = 201 – 400 m, etc.).



Use of native and exotic habitats

Many (perhaps most) flycatcher breeding sites are comprised of spatially complex habitat mosaics, often including both exotic and native vegetation. Within a site, flycatchers often use only a part of the patch, with territories frequently clumped and/or distributed near the patch edge. Therefore, the vegetative composition of individual territories may differ from the overall composition of the patch.

Although detailed territory-based habitat measurements are lacking for the majority of Southwestern Willow Flycatcher breeding sites, it is important to characterize the use of native and exotic habitats. To do so, we classified the habitat at each site into one of four broad categories, based on the overall species composition of the tree/shrub layer(s) of the site. The categories were:


Native	(>90% native vegetation)
Mixed – >50% Native	(50-90% native vegetation)
Mixed – >50% Exotic	(50-90% exotic vegetation)
Exotic	(>90% exotic vegetation)

Habitat patches comprised of Native vegetation account for approximately half (49%) of the known flycatcher territories (Figure 4). Although only 8% of territories occur at Exotic sites, another 41% are located within sites where the habitat includes native/exotic mixtures. In many of these cases, exotics are contributing significantly to the habitat structure by providing the dense lower-strata vegetation that flycatchers prefer.

Dominance of native vegetation at the single largest flycatcher site (121 territories in the Cliff-Gila Valley, NM) substantially affects this habitat summary. Removing it from the analysis, the proportion of rangewide territories occurring in Native sites decreases to 41%, Mixed – Mostly Native sites account for 25% of territories, 21% of territories occur in Mixed – Mostly Exotic sites, and Exotic sites support 10% of territories.

Figure 4.

Percentage of flycatcher territories occurring within breeding sites of differing compositions of native and exotic vegetation.



Dominant tree species at breeding sites


Most flycatcher breeding sites are comprised of spatially complex mosaics of different tree species. Within a site, flycatchers often use only a part of the patch, with territories frequently clumped and/or distributed near the patch edge. Therefore, the dominant tree species may differ between a patch and an individual territory within that patch. Generally, detailed territory-based habitat measurements are lacking for the majority of Southwestern Willow Flycatcher breeding sites. Despite this limitation, it is useful to characterize the dominant tree species within known flycatcher breeding sites.

To characterize the degree to which flycatchers breed in habitats dominated by particular tree species, we tallied the number of territories occurring in breeding sites dominated by particular tree species. Over half (54%) of territories are found at sites where willow (*Salix spp*) is the dominant tree species (Figure 5). Approximately one-fourth are located at sites where saltcedar (*Tamarix spp*) predominates, and 13% are in patches where boxelder (*Acer spp*) is the most common habitat component. Taken together, sites dominated by all other tree species account for only about 6% of territories.

The large percentage of territories located in boxelder dominated habitats might suggest that boxelder sites are widely used across the Southwestern Willow Flycatcher's range. However, boxelder dominated breeding habitats occur only in the Cliff-Gila Valley, New Mexico. Removing that site from the analysis, no territories are found in boxelder dominated habitats, and the proportions of rangewide territories at willow and saltcedar sites increase to 62% and 31%, respectively.

Figure 5.

Percentage of flycatcher territories occurring within breeding sites dominated by particular tree species.



Administration/management of sites and territories

One factor important in conservation and recovery planning is the nature of ownership or “administration” of a site – e.g., whether management of the site is the responsibility of private landowners, the government, or some other entity. We examined this in two ways – first by site, then by territory.

By Site (Figure 6a): Forty-six percent of known breeding sites are under federal government administration, and 26% are on privately owned lands. State/local/municipal governments account for another 12% of sites, and 6% are administered by Native American tribes.

By Territory (Figure 6b): Federal and private lands each account for equal numbers of flycatcher territories (44%). This underscores the importance of working with private landowners as flycatcher conservation and recovery efforts proceed. Roughly a third (35%) of the flycatcher territories on privately owned lands are in the Cliff-Gila Valley, New Mexico.

Figure 6a

Administration by Sites





Figure 6b

Administration by Territories



SUMMARY

- We have learned of many new breeding sites and territories in the last eight years, thanks to extensive survey efforts throughout the southwest. In 1993, there were only 111 known territories distributed among 30 breeding sites. The current count (as of 2000) is 938 territories located among 209 sites (but remember the earlier caution about lack of standard definition for “site”).
- Most territories are found within small breeding sites (those sites with five or fewer territories). There are fewer than a half-dozen sites with 50 or more territories.
- We know of 56 sites that have been “extirpated” since 1993 - almost all were very small sites (five or fewer territories). Because these were primarily small sites, these extirpations account for only a small percentage of known territories; however, they underscore the vulnerability of small sites to extirpation.
- The states of California, Arizona, and New Mexico account for 89% of known territories. Nevada, Colorado, and Utah collectively have 11% of the known territories. We know virtually nothing about the current status of the Southwestern Willow Flycatcher in Texas.
- Southwestern Willow Flycatchers are distributed over a wide elevation range, with most from sea level to 1600 m, but a few territories are located as high as 2530 m in elevation.
- Just under half of territories are in native habitat another 27 % are in habitats having a 50 % or greater exotic component. A large percentage of the native habitat territories occur at one site –the Cliff-Gila Valley in New Mexico. Over 90% of territories are in habitats where willow, saltcedar, or boxelder are the dominant tree species. Flycatchers breed in boxelder dominated habitats only in the Cliff-Gila Valley, New Mexico.
- Roughly half (46%) of sites are on federally-controlled lands, while 26% are on private lands; these privately owned sites account for 38% of known territories. Approximately one-third (35%) of territories on privately owned sites are found in the Cliff-Gila Valley, New Mexico.

List of literature cited and data sources: Journal Articles and Reports

- Farmer, C., M.A. Holmgren, and S.I. Rothstein. 2001. DRAFT REPORT: Distribution and abundance of southwestern willow flycatchers on Vandenberg Air Force Base and the Lower Santa Ynez River. Dept. Ecology, Evolution, and Marine Biology, University of California, Santa Barbara, California.
- R. Furtek, T. Gallagher, C. Tomlinson. 2000. Southern Region Bird Projects Federal Fiscal year 1999, State Fiscal Year 2000. Nevada Division of Wildlife. September 2000
- Harmsworth Associates. 2001. Least Bell's vireo and southwestern willow flycatcher surveys in the North Ranch Policy Plan Area, Orange County, California. Draft report prepared for USFWS, CDFG, County of Orange, Metropolitan Water Dist. of Southern California, Irvine Ranch District, Transportation Corridor Agencies, and the Irvine Company. February.
- Hoffman, S. M. 2000 Status and management of the least Bell's vireo and southwestern willow flycatcher on the Santa Ana River in Riverside county above Prado Basin including Hidden Valley 2000. Prepared for the Santa Ana Watershed Association, Orange County Water District, U.S.F.W.S. and California Department of Fish and Game.
- Konecny Biological Services. 2000. Annual report to the U.S. Fish and Wildlife Service, Carlsbad Field Office, CA.
- Kus, B.E. 2001. Distribution, abundance, and breeding activities of the southwestern willow flycatcher at Marine Corps Base Camp Pendleton, California, in 2000. Final Report prepared for Assistant Chief of Staff, Environmental Security, Marine Corps Base Camp Pendleton.
- Langridge, S.M. and M.K. Sogge. 1998. Banding and genetic sampling of Willow Flycatchers in Utah: 1997 and 1998. USGS Colorado Plateau Field Stn Report to the U.S. Bureau of Reclamation, Salt Lake City, UT.
- McKernan, R.L. and G. Braden. 2001. Status, distribution, and habitat affinities of the Southwestern Willow Flycatcher along the Lower Colorado River, Year 5 – 2000. Report by the San Bernardino County Museum, Redlands, CA.
- Netter, M.R., E.H. Paxton and M.K. Sogge. 1998. Banding and movements of the Southwestern Willow Flycatcher at Roosevelt Lake and San Pedro River/Gila River confluence, Arizona – 1998. USGS Colorado Plateau Field Station Report. 48 pp
- Owen, J. C. and M. K. Sogge. 1997. Banding and Genetic Sampling of Willow Flycatchers in Colorado - 1996 & 1997 Summary Report. USGS Colorado Plateau Field Station: Flagstaff, Arizona.
- Paradzick, C.E., T.D. McCarthy, R.F. Davidson, J.W. Rourke, M.W. Sumner, and A.B. Smith. 2001 Southwestern Willow Flycatcher 2000 survey and nest monitoring report. Arizona Game and Fish Department, Nongame and Endangered Wildlife Technical Report 175. 93 pp.
- Paxton, E.H. 2000. Molecular genetic structuring and demographic history of the Willow Flycatcher. Masters Thesis. Northern Arizona University, Flagstaff, AZ.
- Paxton, E. and M.K. Sogge. 1996. Banding and population genetics of Southwestern Willow Flycatchers in Arizona – 1996 Summary Report. USGS Colorado Plateau Field Station Report. 25 pp.
- Pike, J., D. Pelligrini, S. Reynolds, and L.H. Hays. 2000. The status and management of the least Bell's vireo and southwestern willow flycatcher within the Prado Basin, California, 1986- 2000. Prepared for the Orange County Water District, U.S. Army Corps of Engineers, Los Angeles District, U.S. Fish and Wildlife Service, and California Department of Fish and Game.

- Porter, A. and K.S. Day. 2000. Summary of Southwestern Willow Flycatcher (*Empidonax traillii extimus*) Surveys Conducted in the Virgin River Drainage, Washington County, Utah in 2000. Utah Division of Wildlife Resources Publication No. 00-32, Cedar City, UT.
- Recon. 2000. Post-survey notification- focused survey results for the least Bell's vireo and southwestern willow flycatcher on the Windmill Creek Channel Improvements in Oceanside, California. August 14. Report to U.S. Fish and Wildlife Service.
- Recon. 2000b. Post-survey notification- focused survey results for the least Bell's vireo and southwestern willow flycatcher on the proposed bridge and thoroughfare District No. 4 in Carlsbad, California. August 14. Report to the U.S. Fish and Wildlife Service.
- Sedgwick, J.A. 2001. Geographic variation in the song of Willow Flycatchers: differentiation between *Empidonax traillii adustus* and *E.t. extimus*. *Auk* 118:366-379.
- Sogge, M.K., R.M. Marshall, S.J. Sferra and T.J. Tibbitts. 1997. A Southwestern Willow Flycatcher natural history summary and survey protocol. National Park Service Technical Report NPS/NAUCPRS/NRTR-97/12. 37 pp.
- Sogge, M.K., S.J. Sferra, T.D. McCarthey, S.O. Williams and B.E. Kus. 2000. Southwestern Willow Flycatcher breeding site and territory summary - 1999. Report to the U.S. Fish and Wildlife Service, Albuquerque, NM.
- Sugnet and Associates. 1998. Presence or absence survey for Southwestern Willow Flycatcher (*Empidonax traillii extimus*) for Colorado Department of Transportation Project #FC-NH (CX) 160-2(048) (SH160 Farmington Hill to Bayfield). Report to US Fish and Wildlife Service, Grand Junction, CO.
- Sugnet and Associates. 2000. Presence or absence survey for Southwestern Willow Flycatcher (*Empidonax traillii extimus*) at Piano Creek Ranch. Report to US Fish & Wildlife Serv., Grand Junction, CO.
- Sweetwater Authority. 2001. Endangered species monitoring report for the Sweetwater Reservoir Habitat Management Program and Urban Runoff Diversion System, Phase II. Prepared by P. Famolaro and L. Tikkanen Reising for the U.S. Fish and Wildlife Service and Calif. Dept. of Fish and Game.
- Turnbull, Jennifer L. 2001. Tonner Canyon southwestern willow flycatcher surveys, 2000. Report of findings to Planning Consultants Research, Irvine, CA.
- Varanus Biological Services, Inc. 2000. Report of baseline surveys of existing species, San Felipe Creek and adjoining CDFG acquisition properties, San Diego County, California. Report prepared under contract to California Department of Fish and Game.
- Varanus Biological Services, Inc. 2000. Report: Results of general surveys for arroyo toad and protocol surveys for least Bell's vireo, southwestern willow flycatcher, California gnatcatcher, Salesian Mission property, California 76, Pala, San Diego County, California. Report prepared for URS Geiner Corporation.
- Varanus Biological Services, Inc. 2000. Least Bell's vireo and willow flycatcher surveys and monitoring, Cannon Road Extension Wetlands Mitigation Project, Macario Canyon, Carlsbad, San Diego County, California. City Project No. 3184. Report prepared for the City of Carlsbad.
- Wells, J., and J. Turnbull. 2000. Year 2000 western San Luis Rey River least Bell's vireo and southwestern willow flycatcher survey and nest monitoring program. Unpublished report prepared for the U.S. Army Corps of Engineers, Los Angeles District.
- Whitfield, M.J. and J.C. Lynn. 2001. Southwestern willow flycatcher (*Empidonax traillii extimus*) surveys, nest monitoring, and removal of brown-headed cowbirds on the South Fork Kern River, California in 2000. Final report prepared for the U.S. Army Corps of Engineers and California Department of Fish and Game.
- Williams, S.O. III. 2001. Summary of Willow Flycatcher surveys in New Mexico during 2000. Unpublished summary provided by New Mexico Department of Game and Fish, Santa Fe, NM.

List of data sources: Personal communications and unpublished data

Bloom, Peter. 13611 Hewes Ave, Santa Ana, CA 92705.

Braden, Gerald. San Bernardino County Museum, 2024 Orange Tree Lane, Redlands, CA 92374. 909-307-2669 x232; gbraden@ccr.co.san-bernardino.ca.us

Michael Cassell. U.S. Bureau of Land Management, San Luis Valley, CO. Michael_Cassell@co.blm.gov

Day, Keith. BLM Cedar City Office, Cedar City, UT. Keith_Day@ut.blm.gov

Greaves, Jim. 2416 De la Vina St #2, Santa Barbara, CA 93105. 805-563-2905: greaves@silcom.com

Guthrie, Danial. Keck Science Center, 925 N. Mills Ave, Claremont, CA 91711. dguthrie@jso.claremont.edu

Haas, William. Varanus Biological Service, 10601-A Tierra Santa Blvd. Suite 127, San Diego, CA 92124. 858-536-8762; wehaas@ibm.net

Hays, Loren. U.S. Fish and Wildlife Service, 2730 West Loker Ave. West, Carlsbad, CA 92008. 760-431-9440; Loren_Hays@nbs.gov

Ireland, Terry. U.S. Fish and Wildlife Service, 764 Horizon Drive, South Annex A Grand Junction, CO 81506. 970-243-2778 ext 16; Terry_Ireland@fws.gov

Krueger, Jeri. U.S. Fish and Wildlife Service, 1510 N. Decatur Blvd, Las Vegas, NV 89108. Jeri_Krueger@fws.gov

Laymon, Steve. P.O. Box 1236, Weldon, CA 93283. slaymon@lightspeed.net

Lucero, Jill. U.S. Bureau of Land Management, San Luis Valley, CO. Jill_Lucero@co.blm.gov

McKernan, Marnie. San Bernardino County Transportation Department/Flood Control, 825 E. Third Street, San Bernardino CA 92415

McKernan, Robert. San Bernardino County Museum, 2024 Orange Tree Lane, Redlands, CA 92374. 909-307-2669 x232; rmckernan@ccr.co.san-bernardino.ca.us

Paxton, Eben. Colorado Plateau Field Station, P.O. Box 5614, Northern Arizona University, Flagstaff, AZ 86011. 520-556-7311 ext 229; Eben.Paxton@nau.edu.

Stone, Kelli. Alamosa National Wildlife Refuge, Alamosa, CO. Kelli_Stone@fws.gov

Summer, Priscilla. USFS Cedar City Ranger District, 1789 N Wedgewood Lane, Cedar City, UT. 94720. psummers@fs.fed.gov

Tomlinson, Cris. Nevada Division of Wildlife, 474 Vegas Drive, Las Vegas, NV 89108. 702-486-5127 x 3217; wctommy@aol.com

Utah Division of Wildlife Resources, 1594 W. North Temple St, Suite 2110, Salt Lake City, UT 84114-6301. 801-538-4764

Wallen, Rick. Division of Resources Management, Bryce Cyn National Park, UT. Rick_Wallen@nps.gov

Whitfield, Mary. Kern River Research Center, P.O. Box 662, Weldon, CA 92383. 760-378-2402; wifl@lightspeed.net

Acknowledgements

A synthesis of data from so many sites over such a broad geographic range was only made possible by the efforts of a great many cooperators. Originally, these data were gathered by literally hundreds of agency and non-governmental biologists surveying for thousands of hours, often in very difficult field conditions. Their dedication and efforts are greatly appreciated. Further, the ability to report specific information for each site was aided by agencies and people that provided detailed summary information; our sincere thanks go to the individuals listed below.

For Arizona: Rebecca Davidson, Charles Paradzick, James Rourke, Alex Smith, and Michael Sumner (all of the Arizona Game and Fish Department); Tim Tibbitts (National Park Service); and Kerry Christensen (Hualapai Tribe). **For California:** Peter Bloom, Gerald Braden (San Bernardino County Museum), Marnie Crook; Pete Famolaro (Sweetwater Authority); Jim Greaves; William Haas (Varanus Biological Services); Loren Hays (USFWS); Mark Holmgren, Robert McKernan (San Bernardino County Museum); David Perekesta (USFWS); and Mary Whitfield (Kern River Research Center). **For Colorado:** Terry Ireland (USFWS), Jill Lucero (BLM), Kelli Stone (USFWS). **For Nevada:** Jeri Krueger (USFWS), Robert McKernan (SBCM), Cris Tomlinson (Nevada Division of Wildlife). **For New Mexico:** Steve Albert (Pueblo of Zuni; Darrel Ahlers and Larry White (US Bureau of Reclamation); Scott Stoleson (US Forest Service); and Pat Zenone. **For Utah:** Keith Day (BLM) and Frank Howe (Utah Division of Wildlife Resources).

Terry Arundel (USGS), Charles Drost (USGS), and Eben Paxton provided essential technical assistance with this database. This project was sponsored in part using federal funds from the U.S. Geological Survey and the U.S. Bureau of Reclamation (Phoenix Area Office).