



Figure 1. Precambrian cover (uplift) (folded) of Rocky Mountain region.

Figure 2. Aeromagnetic anomaly map of northwestern United States showing major Precambrian tectonic boundaries and principal Proterozoic faults and shear zones. CB, Cheyenne belt suture; CF, Cedar Creek suture zone; GFSZ, Great Falls suture zone; LC, Lewis and Clark fault zone; SF, Snake River fault zone; SR, Snake River Plain volcanic belt. Aeromagnetic map prepared by North American Magnetic Anomaly Group (2002).

INTRODUCTION

Newly updated aeromagnetic data of Montana, in conjunction with the known geologic framework of basement rocks in the State, have been combined to produce a new interpretive Precambrian basement map of Montana. Crystalline rocks of Precambrian age underlie the entire State, but have been exposed only in the coals of Laramide-age mountains in southwestern Montana (Fig. 1). A thick cover of Phanerozoic rocks of Paleozoic and Mesozoic age overlies the Precambrian crystalline rocks in western Montana (Wilson, 1986; Winston and Link, 1993) and Phanerozoic strata cover the basement in eastern Montana. The Belt Supergroup and equivalent rocks of Proterozoic age are not included as part of the Precambrian basement in the geologic map.

This report confirms and extends the hypothesis of O'Neill (1999) that the Great Falls tectonic zone, defined mainly from tectonic features and gravity anomalies in the Wyoming province, is a major tectonic province of the Precambrian basement in the northern part of the State. The Great Falls tectonic zone is a major tectonic province of the Precambrian basement in the northern part of the State. The Great Falls tectonic zone is a major tectonic province of the Precambrian basement in the northern part of the State. The Great Falls tectonic zone is a major tectonic province of the Precambrian basement in the northern part of the State.

GENERAL GEOLOGY OF CRYSTALLINE BASEMENT

The basement geologic framework of Montana is complex, recording a history of at least 3 billion years of time. Archean granite-granulite intrusions and gneissic rocks contain early orogenic, tectonic, and tectonic features. The Paleoproterozoic and Paleoproterozoic rocks contain early orogenic, tectonic, and tectonic features. The Paleoproterozoic and Paleoproterozoic rocks contain early orogenic, tectonic, and tectonic features. The Paleoproterozoic and Paleoproterozoic rocks contain early orogenic, tectonic, and tectonic features.

TECTONIC MODEL

We introduce the term "Trans-Montana orogeny" in this report for the orogenic assembly comprising the Precambrian terranes in the northern part of the State. The orogenic assembly comprising the Precambrian terranes in the northern part of the State. The orogenic assembly comprising the Precambrian terranes in the northern part of the State.

TECTONIC MODEL

A conceptual model for evolution of the Trans-Montana orogen is shown in Figure 3. Stage 1 (Fig. 3a) represents the initial tectonic assembly of the Trans-Montana orogen, which is a major tectonic province of the Precambrian basement in the northern part of the State. The orogenic assembly comprising the Precambrian terranes in the northern part of the State. The orogenic assembly comprising the Precambrian terranes in the northern part of the State.

ACCRETED TERRAINS

The two terranes accreted to the northern margin of the Wyoming craton in Montana during the Trans-Montana orogeny are interpreted as having been

ACCRETED TERRAINS

The two terranes accreted to the northern margin of the Wyoming craton in Montana during the Trans-Montana orogeny are interpreted as having been

ACCRETED TERRAINS

The two terranes accreted to the northern margin of the Wyoming craton in Montana during the Trans-Montana orogeny are interpreted as having been

PRECAMBRIAN BASEMENT GEOLOGIC MAP OF MONTANA—AN INTERPRETATION OF AEROMAGNETIC ANOMALIES

By P.K. Sims, J.M. O'Neill, Viki Bankey, and E. Anderson 2004