

## **Reported Historic Asbestos Prospects and Natural Asbestos Occurrences in the Central United States**

By Bradley S. Van Gosen

### **U.S. Geological Survey Open-File Report 2006-1211 Version 1.0**

#### **Introduction**

This map and the accompanying dataset (*asbestos\_sites.xls*) provide information for 26 natural asbestos occurrences in the Central United States (U.S.) using descriptions found in the geologic literature. Data on location, mineralogy, geology, and relevant literature for each asbestos site are provided in the aforementioned digital file. Using the map and digital data in this report, the user can examine the distribution of previously reported asbestos occurrences and their geological characteristics in the Central U.S. This report is part of an ongoing study by the U.S. Geological Survey to identify and map reported natural asbestos occurrences in the U.S., which began with U.S. Geological Survey Open-File Report 2005-1189 (Van Gosen, 2005). These reports are intended to provide State and local government agencies and other stakeholders with geologic information on natural occurrences of asbestos.

The file *asbestos\_sites.xls* was compiled through a systematic State-by-State search of the geologic literature. Although this asbestos dataset represents a thorough study of the published literature, it can not be construed as a complete list. An asbestos site was included only when the literature source specifically mentioned asbestos and (or) described the commonly recognized asbestos minerals as occurring in the asbestiform crystal morphology. No attempt was made to interpret the presence of asbestos from the context of the geology-mineralogy description if asbestos was not explicitly described. The user should refer to the references cited with each asbestos site entry for descriptions of these occurrences. These asbestos occurrences were reported to exist in outcrop exposures or rock exposed by mining operations. Note that these site descriptions apply to the time of each report's publication. No field verification of the sites was performed, nor were evaluations of potential exposure made at these sites. Many of the sites are likely to have been subsequently modified by human activities since their description, sometimes substantially. For example, since the time that the source literature was published there may have been remediation of the site or it may have been either exposed or covered by more recent development.

#### **What is Asbestos?**

The history of asbestos discovery and usage is at least 5,000 years old, extending back to the ancient civilizations in Greece and what is now Italy (see Ross and Nolan, 2003). Historically, asbestos is a generic commercial-industrial term used to describe a group of specific silicate minerals that form as long, very thin mineral fibers, which combine to form bundles. When handled or crushed, asbestos bundles readily separate into individual mineral fibers. The special properties of commercial-grade asbestos—

long, thin, durable mineral fibers and fiber bundles with high tensile strength, flexibility, and resistance to heat, chemicals, and electricity—have made it well suited for a number of commercial applications (Ross, 1981; Zoltai, 1981; Cossette, 1984; Ross and others, 1984; Skinner and others, 1988). Asbestos has been especially used for its insulating and fire-resistant properties in many types of products (see Virta and Mann, 1994; Ross and Virta, 2001).

Currently, commercial and regulatory definitions of asbestos most commonly include chrysotile, the asbestiform member of the serpentine group, and several members of the amphibole mineral group, including the asbestiform varieties of (1) riebeckite (commercially called crocidolite), (2) cummingtonite-grunerite (commercially called amosite), (3) anthophyllite (anthophyllite asbestos), (4) actinolite (actinolite asbestos), and (5) tremolite (tremolite asbestos). Other amphiboles are known to occur in the fibrous or asbestiform habit (Skinner and others, 1988), such as winchite, richterite (Meeker and others, 2003), and fluoro-edenite (Gianfagna and Oberti, 2001; Gianfagna and others, 2003), but to date they have not been specifically listed in the asbestos regulations. The many different ways that asbestos and asbestiform and other related terms have been described are summarized in Lowers and Meeker (2002).

Historically, chrysotile has accounted for more than 90 percent of the world's asbestos production, and it presently accounts for over 99 percent of the world production (Ross and Virta, 2001; Virta, 2002). Mining of crocidolite (asbestiform riebeckite) and amosite (asbestiform cummingtonite-grunerite) deposits accounts for most of the other asbestos production, and small amounts of anthophyllite asbestos have been mined in Finland and the U.S. in the past (Ross and Virta, 2001; Van Gosen, 2005). Asbestos is no longer mined in the U.S. The last U.S. asbestos operation mined chrysotile deposits in California; this mine closed in 2002.

### **Naturally Occurring Asbestos**

Mounting evidence throughout the 20th century indicated that inhalation of asbestos fibers caused respiratory diseases that have seriously affected many workers in certain asbestos-related occupations (Tweedale and McCulloch, 2004; Dodson and Hammar, 2006). Airborne exposures to asbestos have been linked to a number of serious health problems and diseases, including asbestosis, lung cancer, and mesothelioma. Additional asbestos information is available online at <http://www.epa.gov/asbestos/> and <http://www.atsdr.cdc.gov/asbestos/>.

Asbestos-bearing materials (some pipe wrappings and insulation, as examples) are frequently uncovered in older buildings and structures, causing health concerns for those individuals exposed to dusts liberated from these materials. As older structures are continually torn down or remodeled, contact with asbestos-bearing materials will likely be of concern for decades to come. The proper handling and disposal of these man-made asbestos materials is addressed by a number of federal regulations. Less straightforward is the regulation and management of "naturally occurring asbestos" (NOA), which has recently gained the attention of regulatory agencies, health agencies, and citizen groups. NOA is asbestos found in-place in its natural state; that is, asbestos minerals in bedrock exposed by man's excavations or by natural weathering. The geology of asbestos is summarized in Van Gosen (2006).

NOA is of concern due to potential exposures to microscopic fibers that can become airborne if asbestos-bearing rocks are disturbed by natural erosion or human activities (road building, urban excavations, agriculture, mining, crushing, and milling, as just a few examples). Examples of occupational and environmental exposures to asbestos are described in Nolan and others (2001) and Ross and Nolan (2003).

Recent attention towards NOA was spurred by the recognition of high incidences of asbestos-related mortality and respiratory disease in vermiculite miners and residents of Libby, Montana. This disease cluster has been attributed to fibrous and asbestiform amphibole particles within the vermiculite ore body once mined and milled near the town from 1923 to 1990 (Peipins and others, 2003). Meeker and others (2003) described in detail the fibrous and asbestiform amphibole minerals intergrown with the Libby vermiculite deposit.

Large areas of exposed ultramafic bedrock in northern California, some now densely populated by housing and infrastructure, have become the focus of recent attention because they contain chrysotile and possibly tremolite-actinolite asbestos (Churchill and Hill, 2000; Clinkenbeard and others, 2002; Ross and Nolan, 2003; Swayze and others, 2004).

The history and study of naturally occurring asbestos and the multiple, complex issues that surround asbestos are discussed in Campbell and others (1977), Ross (1981), Stanton and others (1981), Zoltai (1981), Levadie (1984), Skinner and others (1988), Mossman and others (1990), Occupational Safety and Health Administration (1992), Guthrie and Mossman (1993), van Oss and others (1999), Nolan and others (2001), Virta (2002), Plumlee and Ziegler (2003), and Dodson and Hammar (2006). Current federal regulations are provided in the Code of Federal Regulations (available online at <http://www.gpoaccess.gov/cfr/>). However, these asbestos regulations do not specifically address exposures to natural occurrences of asbestos.

### **Asbestos in the Central United States**

Based on this study, natural asbestos appears to be relatively sparse in the Central U.S., especially when compared with other regions of the country (Van Gosen, 2005). The asbestos occurrences in the Central U.S. are found in a diverse variety of geologic settings (see *asbestos\_sites.xls*), including altered ultramafic rocks, mafic alkaline igneous intrusions, dolomitic marbles, metamorphosed iron-formations, iron-rich skarns, and talc deposits that replace dolostones. The majority of the sites on the map represent asbestos that occurs as a minor accessory mineral within bedrock or a larger mineral deposit. Only five sites in this region were once prospected for asbestos (evaluated for commercial use), and apparently none of these deposits were further developed. No records were found to indicate that asbestos was ever mined from this region of the U.S.

### **Fibrous Amphiboles in the Central United States**

During this study, several examples were noted in the geologic literature that mentioned the presence of fibrous amphiboles in outcrops or within a metallic ore body (copper, gold, iron). These examples are shown on the map and described in a separate dataset (*fibrous\_amphiboles.xls*). Amphibole asbestos was not specifically mentioned in

the descriptions of these deposits. However, these sites indicate geologic settings with the potential to host asbestos. This is because the geologic settings of these examples are similar to those that elsewhere form and host the reported asbestos, including metamorphosed and metasomatized mafic rock, skarn, marble, and metamorphosed banded iron-formation. Thus, a discovery of asbestos in these areas would not be unusual from a geologic standpoint. Also, the distinction between "fibrous" amphibole and "regulatory" amphibole asbestos is often not clear-cut in natural amphibole-bearing deposits. This regulatory dilemma—clear distinction between "fibrous" particle and "regulatory fiber" (Occupational Safety and Health Administration, 1992)—is well illustrated by the descriptions provided in this dataset (*fibrous\_amphiboles.xls*). For example, elongate amphibole particles, mostly grunerite and cummingtonite, in metamorphosed banded iron-formation of the eastern Mesabi iron range, Minnesota, have been variously described as tabular or fibrous (French, 1968), or asbestiform (Great Lakes Advisory Board, 1975), depending on the author (see Lowers and Meeker, 2002, for published definitions of these terms). Grunerite and cummingtonite are widespread rock-forming minerals in the Biwabik iron-formation in this area, but do not everywhere occur in fibrous form. Concentrations of grunerite or cummingtonite occur widely within contact-metamorphosed banded iron-formation elsewhere in northern Minnesota. They also occur widely in other iron ranges of the Lake Superior region, particularly the western part of the Marquette iron range in Michigan and the western part of the Gogebic iron range in Wisconsin. However, descriptions of these amphibole deposits are not sufficiently detailed to define areas where the amphiboles occur in fibrous form, and thus specific sites could not be shown on this map. These amphiboles may or may not meet the regulatory criteria of asbestos, which requires site-specific detailed microscopic analyses.

### References Cited

- Campbell, W.J., Blake, R.L., Brown, L.L., Cather, E.E., and Sjoberg, J.J., 1977, Selected silicate minerals and their asbestiform varieties—Mineralogical definitions and identification-characterization: U.S. Bureau of Mines Information Circular IC-8751, 56 p.
- Churchill, R.K., and Hill, R.L., 2000, A general location guide for ultramafic rocks in California—Areas more likely to contain naturally occurring asbestos: Sacramento, Calif., California Department of Conservation, Division of Mines and Geology, DMG Open-File Report 2000-19. Available online at <http://www.consrv.ca.gov/>
- Clinkenbeard, J.P., Churchill, R.K., and Lee, Kiyoung, eds., 2002, Guidelines for geologic investigations of naturally occurring asbestos in California: Sacramento, Calif., California Department of Conservation, California Geological Survey Special Publication 124, 70 p.
- Cossette, Marcel, 1984, Defining asbestos particulates for monitoring purposes, *in* Levadie, Benjamin, ed., Definitions for asbestos and other health-related silicates: Philadelphia, Pa., American Society for Testing and Materials, ASTM Special Technical Publication 834, p. 5-50.

- Dodson, R.F., and Hammar, S.P., eds., 2006, *Asbestos—Risk assessment, epidemiology, and health effects*: Boca Raton, Fla., Taylor & Francis Group, 425 p.
- French, B.M., 1968, Progressive contact metamorphism of the Biwabik iron-formation, Mesabi Range, Minnesota: Minneapolis, Minn., University of Minnesota, Minnesota Geological Survey Bulletin 45, 103 p.
- Gianfagna, A., Ballirano, P., Bellatreccia, F., Bruni, B., Paoletti, L., and Oberti, R., 2003, Characterization of amphibole fibres linked to mesothelioma in the area of Biancavilla, eastern Sicily, Italy: *Mineralogical Magazine*, v. 67, no. 6, p. 1221-1229.
- Gianfagna, Antonio, and Oberti, Roberta, 2001, Fluoro-edenite from Biancavilla (Catania, Sicily, Italy)—Crystal chemistry of a new amphibole end-member: *American Mineralogist*, v. 86, p. 1489-1493.
- Great Lakes Research Advisory Board, 1975, *Asbestos in the Great Lakes basin, with emphasis on Lake Superior—A report to the International Joint Commission*: Great Lakes Research Advisory Board, 100 p.
- Guthrie, G.D., and Mossman, B.T., eds., 1993, *Health effects of mineral dusts*: Mineralogical Society of America, *Reviews in Mineralogy*, v. 28, 584 p.
- Levadie, Benjamin, ed., 1984, *Definitions for asbestos and other health-related silicates*: Philadelphia, Pa., American Society for Testing and Materials, ASTM Special Technical Publication 834, p. 1-147.
- Lowers, Heather, and Meeker, Greg, 2002, *Tabulation of asbestos-related terminology*: U.S. Geological Survey Open-File Report 02-458, 74 p. Available online at <http://pubs.usgs.gov/of/2002/ofr-02-458/>
- Meeker, G.P., Bern, A.M., Brownfield, I.K., Lowers, H.A., Sutley, S.J., Hoefen, T.M., and Vance, J.S., 2003, The composition and morphology of amphiboles from the Rainy Creek complex, near Libby, Montana: *American Mineralogist*, v. 88, nos. 11-12, Part 2, p. 1955-1969.
- Mossman, B.T., Bignon, J., Corn, M., Seaton, A., and Gee, J.B.L., 1990, Asbestos—Scientific developments and implications for public policy: *Science*, v. 247, p. 294-301.
- Nolan, R.P., Langer, A.M., Ross, M., Wicks, F.J., and Martin, R.F., eds., 2001, *The health effects of chrysotile asbestos—Contribution of science to risk-management decisions*: The Canadian Mineralogist, Special Publication 5, 304 p.
- Occupational Safety and Health Administration, 1992, 29 CFR Parts 1910 and 1926 [Docket No. H-033-d], Occupational exposure to asbestos, tremolite, anthophyllite and actinolite: *Federal Register*, v. 57, no. 110, Monday, June 8, 1992, p. 24,310-24,331.
- Peipins, L.A., Lewin, M., Campolucci, S., Lybarger, J.A., Miller, A., Middleton, D., Weis, C., Spence, M., Black, B., and Kapil, V., 2003, Radiographic abnormalities and exposure to asbestos-contaminated vermiculite in the community of Libby, Montana, U.S.A.: *Environmental Health Perspectives*, v. 111, no. 14, p. 1753-1759.
- Plumlee, G.S., and Ziegler, T.L., 2003, The medical geochemistry of dusts, soils, and other earth materials, *in* Holland, H.D., and Turekian, K.K., eds., *Treatise on geochemistry*, volume 9—Environmental geochemistry: Amsterdam, Elsevier Pergamon, p. 263-310.

- Ross, Malcolm, 1981, The geologic occurrences and health hazards of amphibole and serpentine asbestos [chapter 6], *in* Veblen, D.R., ed., *Amphiboles and other hydrous pyriboles—Mineralogy: Mineralogical Society of America, Reviews in Mineralogy*, v. 9A, p. 279-323.
- Ross, Malcolm, Kuntze, R.A., and Clifton, R.A., 1984, A definition for asbestos, *in* Levadie, Benjamin, ed., *Definitions for asbestos and other health-related silicates: Philadelphia, Pa., American Society for Testing and Materials, ASTM Special Technical Publication 834*, p. 139-147.
- Ross, Malcolm, and Nolan, R.P., 2003, History of asbestos discovery and use and asbestos-related disease in context with the occurrence of asbestos within ophiolite complexes, *in* Dilek, Yildirim, and Newcomb, Sally, eds., *Ophiolite concept and the evolution of geological thought: Geological Society of America Special Paper 373*, p. 447-470.
- Ross, Malcolm, and Virta, R.L., 2001, Occurrence, production and uses of asbestos, *in* Nolan, R.P., Langer, A.M., Ross, Malcolm, Wicks, F.J., and Martin, R.F., eds., *The health effects of chrysotile asbestos—Contribution of science to risk-management decisions: The Canadian Mineralogist, Special Publication 5*, p. 79-88.
- Skinner, H.C.W., Ross, Malcolm, and Frondel, Clifford, 1988, *Asbestos and other fibrous materials—Mineralogy, crystal chemistry, and health effects: New York, Oxford University Press*, 204 p.
- Stanton, M.F., Layard, Maxwell, Tegeris, Andrew, Miller, Eliza, May, Margaret, Morgan, Elizabeth, and Smith, Alroy, 1981, Relation of particle dimensions to carcinogenicity in amphibole asbestos and other fibrous minerals: *Journal of the National Cancer Institute*, v. 67, p. 965-975.
- Swayze, G.A., Higgins, C.T., Clinkenbeard, J.P., Kokaly, R.F., Clark, R.N., Meeker, G.P., and Sutley, S.J., 2004, Preliminary report on using imaging spectroscopy to map ultramafic rocks, serpentinites, and tremolite-actinolite-bearing rocks in California: U.S. Geological Survey Open-File Report 2004-1304 and California Geological Survey Geologic Hazards Investigation 2004-01, 20 p. Available online at <http://pubs.usgs.gov/of/2004/1304/>
- Tweedale, Geoffrey, and McCulloch, Jock, 2004, Chrysophiles versus chrysophobes—The white asbestos controversy, 1950s-2004: *Isis*, v. 95, p. 239-259.
- Van Gosen, B.S., 2005, Reported historic asbestos mines, historic asbestos prospects, and natural asbestos occurrences in the Eastern United States: U.S. Geological Survey Open-File Report 2005-1189. Available online at <http://pubs.usgs.gov/of/2005/1189/>
- Van Gosen, B.S., 2006, Using the geology of asbestos deposits to predict the presence or absence of asbestos in mining and natural environments, *in* Reid, J.C., ed., *Proceedings of the 42nd Forum on the Geology of Industrial Minerals, May 7-13, 2006, Asheville, North Carolina, USA: Raleigh, N.C., North Carolina Geological Survey Information Circular 34 [CD-ROM]*, p. 412-432.
- van Oss, C.J., Naim, J.O., Costanzo, P.M., Giese, R.F., Jr., Wu, W., and Sorling, A.F., 1999, Impact of different asbestos species and other mineral particles on pulmonary pathogenesis: *Clay and Clay Minerals*, v. 47, no. 6, p. 697-707.
- Virta, R.L., 2002, *Asbestos: U.S. Geological Survey Open-File Report 02-149*, 35 p.

- Virta, R.L., and Mann, E.L., 1994, Asbestos, *in* Carr, D.D., ed., *Industrial minerals and rocks*, 6th edition: Littleton, Colo., Society for Mining, Metallurgy, and Exploration, Inc., p. 97-124.
- Wylie, A.G., and Huggins, C.W., 1980, Characteristics of a potassian winchite-asbestos from the Allamoore talc district, Texas: *Canadian Mineralogist*, v. 18, p. 101-107.
- Zoltai, Tibor, 1981, Amphibole asbestos mineralogy [chapter 5], *in* Veblen, D.R., ed., *Amphiboles and other hydrous pyriboles—Mineralogy*: Mineralogical Society of America, *Reviews in Mineralogy*, v. 9A, p. 237-278.