

## METHODS

### Sample Collection and Chemical Analyses

Methods of sample collection were detailed in the Sample and Analysis Plan (USFWS 2003; see Appendix G). All collections of water quality and fish were conducted between March 29 and April 2, 2004. In general, water samples were collected from the lake by way of motorized boat along two transects bisecting each lake. Fish samples were collected by electrofishing boat or trammel net. Chain-of-custody procedures were followed to ensure samples were collected, protected, stored, handled, chemically analyzed, and disposed of properly by authorized personnel. Each sample was labeled with a unique sample number, date and name of water body. Water and fish samples were placed on ice immediately after processing and kept on ice until delivery to the Texas A&M University laboratory the following day. All sampling equipment used to collect and process fish and water samples were decontaminated prior to use.

#### *Collection of Water and Physical Measurements*

Surface-water was sampled up to eight locations along two perpendicular cross-sections in each lake accessed by a boat. Surface water samples were collected from the epilimnion (*i.e.*, within 36 inches [762 mm] of the lake's surface) using a Teflon, DH-95 surface-water sampler, bottle, cap, and nozzle. Surface-water samples analyzed for trace elements were transferred into a polyurethane plastic bottle and then composited in a polyurethane plastic churn-splitter. The composite water was then transferred from the churn using a peristaltic pump, C-flex tubing, and a 0.45-micron in-line capsule filter into each sample bottle. Sample bottles for any particular analyses were filled and preserved immediately. However, whole-water samples to be analyzed for methylmercury were grab-sampled from the epilimnion at up to four locations from each lake using a gloved hand and a Teflon bottle.

A bottle blank was collected during the Red Lake sampling event using reagent water from the laboratory that prepared the methylmercury sample containers to insure the cleanliness of the bottles prior to use in the field. After decontamination of the churn and pump assembly, an equipment blank was collected into a clean container for each sampling event. In total, up to four grab samples of water for methylmercury analysis and two filtered composited samples of water for element analysis (plus 1 equipment field blank) were collected from each lake. While lake water quality is known to have spatial variability vertically, horizontally, and seasonally, the sampling design could not accommodate these factors within its budget.

At each sampling location along a depth profile every 2 to 5 ft (609 to 1,524 mm), the water column was measured for physical properties (*i.e.*, temperature, pH, electrical conductance, dissolved oxygen [DO], and turbidity as a measure of light scatter) using a Hydrolab multi-probe logger (Hydrolab Corporation 1998). The logger's sensors are designed to meet the criteria and specifications in section 2550 (temperature), section 2520-B (specific conductance), section 4500-O (DO), and section 4500-H+ (pH) in Standard Methods for the Examination of Water and Wastewater (American Public Health Association *et al.* 1995). Prior to use each day, the pH, DO, and conductivity probes were calibrated and

maintained according to the manufacturer's instructions (Hydrolab Corporation 1998). Additionally, a simple measurement of light penetration was made with a secchi disk, which was lowered into the water to record the depth at which it appeared to disappear to the observer. These measurements and additional ecological observations were recorded on standardized field notes (field notes forms provided in Appendix H).

#### *Collection and Processing of Fish Tissue*

The USEPA (2000) guidelines were used for determining sample sizes of fish from each lake while also considering logistics, fish availability and budget. The minimum number of fish that we attempted to collect was determined using Table 6-1 in USEPA (2000). The use of 4 composite samples of 5 fish from each lake had the power between 50 to 90 percent to determine a statistically significant difference between the screening value (0.3 mg/kg), and the geometric mean concentration of each lake's fish samples, depending on the variability of mercury within each fish. Fish of the same species and the similar length ( $\pm 10\%$ ) class were composited for each sample. Rainbow trout and channel catfish were the fishes targeted for sampling, as they are the primary fish permitted by the Navajo Nation Fish and Wildlife Department on coldwater and warmwater lakes, respectively. Additionally, these fish have been observed by biologists with the Navajo Nation Natural Heritage Program as being taken as food items by foraging bald eagles.

Fish tissues were collected to measure contaminant concentrations of selected elements and methylmercury. Fish were collected using an electrofishing boat with an adequate capacity to produce pulsed DC current in order to stun fish. The electrofishing boat has a positive foot-activated power control switch as well as other engineered safety features. All sampling personnel were familiar with safe electrofishing techniques including using net poles made of an insulating material such as fiberglass, wearing personal flotation devices, and wearing insulated lineman gloves. Members of the sampling crew were certified in CPR as well as First Aid. Boat electrofishing is a preferred technique to collect fish, as it is time-efficient and less destructive to the fish community. However, at Morgan Lake, the electrofishing boat generator failed and the remaining fish samples were collected with trammel nets. All fish collected were retained in a live well until sampling was completed and were iced after selection. Only live or freshly dead fish (red gills) were retained.

Fish sample preparation was completed in the field. All equipment used to prepare the samples was cleaned with soap and water, rinsed with a dilute nitric acid solution, and then rinsed with de-ionized water. Equipment was also cleaned between each site. Sample preparation included anesthetizing fish, weighing and measuring, removal and compositing of the fillet portions as well as compositing that portion which remained (the "offal"). Total length was measured to the nearest mm using a plastic measuring board. Total weight (using the spring balance) of fish, fillet composites and offal composites to the nearest gram was measured and recorded. Field data, including total lengths and mass of each fish, were recorded at the time of sample collection and processing onto the standardized field forms.

In total, from each lake, five similarly sized fish were composited into four fillet and four offal samples for chemical analyses. Four composite fillet samples for both methylmercury and element analyses as well as four composite offal samples for element analyses were

collected. Fish were filleted using a stainless steel knife on a plastic resin cutting board. Each composite was placed in a new, clear, colorless plastic food-quality bag. Immediately after they were processed, packaged, and labeled, all samples of trout, catfish, and largemouth bass were placed on wet ice in a chest freezer. All samples were then packed with ice, shipped and received by the analytical laboratory on the following day.

### *Chemical Analyses*

Methods of chemical analyses are detailed in Appendix I. Generally, fish samples were weighed and homogenized at the analytical laboratory. Water and fish samples were digested and analyzed either wet or after freeze-drying. Methylmercury was determined by cold vapor atomic absorption spectrophotometry with a Laboratory Data Control Model 1235 Mercury Monitor equipped with a 300 mm absorption cell (modified from Wagemann *et al.* 1997). Remaining elements in fish and water samples were analyzed by a combination of graphite furnace atomic absorption spectrophotometers, hydride generation atomic fluorescence analyzer, or inductively coupled plasma emitting mass spectroscopy. Normal calibration and quality assurance and quality control procedures were used to determine the concentration of elements and methylmercury. Quality control samples were processed in a manner identical to actual samples, and included reagent blanks, standard reference materials, spiked blanks, duplicates, and spiked samples.

### Data Analysis and Statistical Methods

After conversion to wet weight concentrations, fish that had fillets removed were mathematically re-combined (as the sum of weighted concentrations of the parts of both the fillet and the offal) to yield an ‘re-integrated’ whole body fish concentration, thereby allowing comparisons of contaminant concentrations in whole fish with those mathematically re-integrated fish reported herein. A generalized equation was used to calculate re-integrated fish contaminant concentration (Equation 4).

$$\text{Re-integrated fish concentration} = [(fM/wM) \times cF] + [(oM/wM) \times cO] \quad \text{Equation (4)}$$

Where:

fM	=	mass of a fillet (g WW)
wM	=	mass of fillet + mass of offal (g WW)
cF	=	contaminant concentration in a fillet (mg/kg WW)
oM	=	mass of offal (g WW)
cO	=	contaminant concentration in offal (mg/kg WW)

For statistical purposes as well as simplicity, all results, including integrated fish, which were below the laboratory’s instrument detection limit, were replaced with a value one-half the instrument detection limit (USEPA 1998). Where detectable concentrations were below the laboratory’s instrument detection limit in the offal, no descriptive statistics or comparisons were conducted. Several descriptive statistics (*e.g.*, the geometric mean), statistical analyses, and graphical representations were conducted on concentrations of selected contaminants in samples. For these analyses, the software program STATISTICA (version 6.0 by StatSoft Inc. 2001) was used. Unless otherwise noted, statistical significance

in this report refers to a probability of less than or equal to 0.05. Geometric means were calculated in both dry and wet weight concentrations for selected environmental contaminants. The geometric mean provided a measurement of the central tendency of contaminant distributions and was calculated using data converted to their natural logarithms.

Environmental contaminant concentrations were evaluated using two techniques:

1. Concentrations in water or biota were compared to values reported in the literature as ambient or elevated and to Navajo Nation water quality standards (Navajo Nation 2004).
2. A human health and bald eagle risk assessment, as described below.

Risk assessments generally take the form of hazard identification (outlined above as methylmercury), dose-response (including the reference dose [RfD] identified above), exposure models (*e.g.*, amount and frequency of food ingestion, body weight), and risk characterization (identifying concerns and uncertainty).

### Human Health Risk Assessment Considerations

Methylmercury is a developmental poison that can produce adverse effects following a comparatively brief exposure period (*i.e.*, a few months rather than decades) and therefore, short-term dietary patterns may be important (USEPA 2001). Consequently, estimation of recent patterns of methylmercury consumption from fish is the relevant exposure for the health endpoint of concern for people. Because it is not possible to identify the period of development during which mercury is likely to damage the nervous system of the developing fetus or growing child, exposure of women of childbearing age and to her children to mercury through consumption of fish is a cause for concern. Using the default characteristics provided by USEPA (2000), three hypothetical individual populations were modeled:

1. a woman (weighing 65 kg);
2. a child (less than 14 years old and weighing 14.5 kg); and,
3. a man (weighing 78 kg)

Estimates of health risks to human population consumers of fish were calculated and evaluated according to USEPA (2000) and other published data. The calculation of potential human daily intakes of methylmercury due to fish fillet ingestion was calculated according to the following formula:

$$\text{Intake} = \frac{C_f \times \text{SFIR} \times \text{EF}}{\text{BM} \times \text{AT}} \quad \text{Equation (5)}$$

Where:

- Intake = methylmercury intake rate (mg/kg-day)
- $C_f$  = geometric mean methylmercury concentration in fish fillet (mg/kg WW)
- SFIR = subpopulation fish ingestion rate (kg/day)
- EF = exposure frequency (days/year)
- BM = body mass (kg)
- AT = averaging time (days)

The importance of fish consumption patterns by the local population is critical when deriving protective criteria for local fish tissue concentrations as well as identifying risk or any epidemiological concerns (USEPA 2001). However, Navajo Nation-specific information on fish consumption was not available. In the absence of such information, the average nationwide fish consumption rate (17.5 g/day), and the average nationwide subsistence consumption rate (142.4 g/day) were used from the USEPA (2000) to model the exposure to the fish collected during this study. High-end exposure estimates are useful in estimating population risks and establishing exposure limits because they provided a plausible worst-case scenario at the upper end of the exposure distribution. Table 1 summarizes the assumptions of the human health risk assessment. However, selection of appropriate risk level, consumer body mass, exposure duration, and the average meal size are considered risk management decisions of the Navajo Nation.

Reference dose (RfD)	0.0001 mg methylmercury/kg-bw/day
Consumer body mass	78 kg (man) 65 kg (woman) 14.5 kg (child)
Fish consumption rate	17.5 g/day (nationwide average) 142.4 g/day (nationwide subsistence average)
Meal size	8 oz fish fillet per person per week
Exposure duration	14 days (recreational fishing) 156 days (subsistence fishing)
Geometric mean fish fillet methylmercury concentrations for various lake combinations	Asaayi Lake (0.06 mg/kg WW) Morgan Lake (0.01 mg/kg WW) Red Lake (0.39 mg/kg WW) Wheatfields Lake (0.08 mg/kg WW) Coldwater lakes (0.07 mg/kg WW) Warmwater lakes (0.06 mg/kg WW) All lakes combined (0.065 mg/kg WW)
Averaging Time	365 days per year for a lifetime

The number of days per month that an individual consumes methylmercury from the diet can be estimated from data on frequency of fish consumption. Accordingly, the simplifying assumption that the frequency of fish consumption was made for children is the same as for adults. Children's exposures, therefore, on a per kg body weight basis, are higher than those of adults. Since the methylmercury concentrations in the fish consumed are the same at a given site, exposure is the direct ratio of mass ingested per unit of body weight. Without additional information, successful recreational fishing occurred for 14 days per year and subsistence fishing occurred for 156 days per year was assumed. The assumption that the lakes studied were only source of fish in the diets of subsistence adults and children as well as the recreational angler was assumed. Contaminant concentrations used to estimate daily intakes were the geometric mean concentration for each lake individually, for either the coldwater or the warmwater lakes only, and for a combination of all four lakes. Only non-carcinogenic risks posed by methylmercury in fish fillets were modeled for people.

## Bald Eagle Risk Assessment Considerations

An estimated 10,000 to 12,000 bald eagles inhabit the lower 48 United States (USFWS 1995). Bald eagles migrate into the lower forty-eight states only during the winter months; others are resident throughout the year. Bald eagles, like several other avian species, were adversely impacted by DDT and its metabolites during the 1950s, '60s, and '70s. Due to their status as a federally listed "threatened" species and as a national symbol, the potential threat of mercury exposure to bald eagle survival and recovery is a concern.

The Navajo Nation provides suitable migrating and wintering habitats, but has no current or historic nesting records of bald eagles (G. Tom, Navajo Nation Fish and Wildlife Department, written communication, 1999). Migrating bald eagles have been reported to use at least six interior lakes on the Navajo Nation. Except for Morgan Lake, these lakes can typically become frozen-over by November and remain frozen through February. The San Juan and Little Colorado Rivers are also known foraging and wintering sites for bald eagles.

Since information on body weights and food ingestion rates for bald eagles foraging on the Navajo Nation fishing lakes are not readily available, the bald eagle body weights (5.25 kg) used in calculations below were based on the mean of average female body weights reported nationwide by the USEPA (1993). As the avian reference dose for methylmercury is based on adverse reproductive effects manifested by laying females, it is more appropriate to use average female body weights (Russell 2003).

Information presented by the USEPA (1993) regarding metabolically available energy from various prey types and the ability of bald eagles to assimilate this energy allows for the use of methods to estimate daily food requirements. However, attempting to quantify a specific dietary composition for bald eagles is more difficult than for other species with a narrower range of prey types, and is further confounded by food preferences that may vary both geographically and temporally. An additional difficulty in calculating a general food ingestion rate (FIR) for bald eagles arises because of the composition of the diet can also vary substantially between seasons, locations, or individuals. Therefore, discussion of the energy content of diet items and prey composition that the bald eagle may choose are discussed below as they affect the uncertainties of the risk assessment calculations.

Uncertainty and variability described in predictions of human exposures that result from fish consumption are also applicable to the wildlife. It is interesting to note that on a per kilogram body weight basis, predicted exposures to wildlife are much greater than to humans. Estimates of risks to bald eagle foraging on the Navajo Nation were evaluated according to USEPA (1993). The following equation (Equation 6) was used to estimate daily food intake methylmercury:

$$\text{Intake} = \frac{C_f \times F_{fkI} \times \text{FIR} \times \text{EF} \times \text{ED}}{\text{BM} \times \text{AT}} \quad (\text{Equation 6})$$

Where:

- Intake = methylmercury intake rate (mg/kg WW per day)
- $C_f$  = geometric mean methylmercury concentration in whole fish (mg/kg WW)
- $F_{f_i}$  = fraction of fish ingested compared to whole diet (*i.e.*, a unitless decimal)
- ED = exposure duration (years)
- EF = exposure frequency (days/year; assumption was 30 days for migrant, 180 days for wintering, and 365 days for nesting)
- BM = body mass (kg)
- AT = averaging time (days)

The bald eagle diet has been extensively studied throughout the country. Although generally known as a piscivorous species, bald eagles are opportunistic predators and carrion scavengers (Buehler 2000). Many bald eagles consume a mixture of both aquatic and terrestrially derived prey. A wide variety of various birds, mammals, reptiles, amphibians, and crustaceans may serve as additional bald eagle prey (Buehler 2000). Haywood and Ohmart (1986) reported the diet of nesting bald eagles in Arizona as 58% fish, 14% birds, and 28% mammals. Hunt *et al.* (1992) reported the diet of nesting bald eagles in Arizona as 71% fish, 10% birds, and 18% mammals. Hunt *et al.* (1992) further described the diets of bald eagles as varying with habitat setting. Furthermore, the distribution of prey types they consume may vary seasonally. While there is no definitive diet composition preferred by bald eagles in Arizona – they all show that fish are generally the predominant food item, and particularly, that catfish were the principal prey selected.

The diet composition reported by Hunt *et al.* (1992) of 71.4% fish, 10.3% birds, and 18.3% mammals was used for the generic bald eagle risk assessment calculations. Hunt *et al.* (1992) reported that of the fish, 34.5% were catfish, 25.7% were suckers, 24.3 % were carp, and 15.5% were bass, perch, bluegill, or other Centrarchids. Based on the diets of these fish (Sublette *et al.* 1990; Moyle 2002), suckers were classified as trophic level 2 herbivores, carp as trophic level 3 consumers, and catfish as trophic level 3.5 predators. While channel catfish are opportunistic omnivores, consuming whatever prey they can locate, as catfish increase in size, they become increasingly predatory (Moyle, 2002; USEPA 1995b). The fish lengths determined in this study for channel catfish, suggested that an intermediate trophic level of 3.5 be assigned to these fish when eaten by bald eagles. Using the intermediate trophic level breakdown for catfish, together with the other trophic level 4 fish (trout, largemouth bass), this suggests that about 17 percent of the overall estimated biomass would be comprised of trophic level 4 fish. Of the remainder of the overall fish component to the generic diet, 58 percent is classified as trophic level 3 and as 25 percent is trophic level 2.

While an overall dietary methylmercury concentration can be calculated, the amount of prey consumed from each trophic level is the driving factor influencing the amount of methylmercury ingested on a daily basis. The methylmercury concentration in the overall diet for bald eagle is dependent on both the trophic level composition of its diet and the methylmercury concentrations in each of the trophic levels from which the species feeds. In these situations, the bald eagle could obtain a higher mercury dose from eating other piscivorous birds than it would otherwise receive from a strictly fish-based diet.

Of the bird species consumed by bald eagles (10.3% of total biomass in the overall bald eagle diet as reported by Hunt *et al.* 1992), the most commonly seen in prey remains are coots and ducks, representing approximately 5 percent of the total estimated biomass. Several species were exclusively terrestrial (*e.g.*, mountain quail; <1%), and the remainder was primarily piscivorous: grebes, cormorants, and mergansers (5%). Based on the diets of these birds, coots, ducks, quail, and piscivorous birds were all classified as trophic level 3 consumers. Similarly, the mammal species consumed (18.3% of total biomass in overall bald eagle diet), were classified as 85% trophic level 2 and 15% trophic level 3 consumers.

The final bald eagle Food Ingestion Rate (FIR) was based on this generic bald eagle diet (71.4% fish, 10.3% birds, and 18.3% mammals; as described above) and was calculated using the methodology reported by the USEPA (1993, 1995b), wherein the animal's free-living metabolic rate (FMR, in kilocalories per day) is divided by the metabolizable energy from bald eagle prey. The FMR was determined by Nagy's (1987) allometric equation relating FMR for birds to body weight:

$$\begin{aligned} \text{FMR (kcal/day)} &= 2.601 \times (\text{body weight [g]}^{0.640}) \\ \text{FMR} &= 2.601 \times ((5250)^{0.640}) \\ \text{FMR} &= 625 \text{ kcal/day} \end{aligned}$$

According to the USEPA (1993), metabolizable energy (ME) equals gross energy (GE) of food in kcal/kg wet weight times the assimilation efficiency (AE) of the consumer. The USEPA (1993) gave a GE value of 1.2 kcal/kg for bony fishes, while bird tissue GE is 1.9, and the mammal GE is 1.7. The AEs for a bald eagle consuming birds/mammals and fish is given as 78 and 79 percent, respectively.

$$\begin{aligned} \text{ME}_{\text{fish}} &= 1.2 \text{ kcal/g} \times 0.79 = 0.948 \text{ kcal/g fish} \\ \text{ME}_{\text{birds}} &= 1.9 \text{ kcal/g} \times 0.78 = 1.482 \text{ kcal/g birds} \\ \text{ME}_{\text{mammals}} &= 1.7 \text{ kcal/g} \times 0.78 = 1.326 \text{ kcal/g mammals} \end{aligned}$$

If:  $Y$  = grams of birds consumed, and  
 $6.93Y$  = grams of fish consumed (*i.e.*, 71.4% fish  $\div$  10.3% birds = 6.93)  
 $1.78Y$  = grams of mammals consumed (*i.e.*, 18.3% mammals  $\div$  10.3% birds = 1.78)

Then the FIR for each food can be determined by the equation:

$$\begin{aligned} \text{FMR (625 kcal/day)} &= [Y(\text{g}) \times 1.482 \text{ kcal/g birds}] \\ &+ [6.93Y(\text{g}) \times 0.948 \text{ kcal/g fish}] \\ &+ [1.78Y(\text{g}) \times 1.326 \text{ kcal/g mammals}] \\ 625 \text{ kcal/day} &= [1.482Y + 6.57Y + 2.36Y] \\ 625 \text{ kcal/day} &= 10.41Y \end{aligned}$$

Since  $Y = 60.03$  g birds consumed per day

Then  $6.93Y =$  grams of fish consumed  
 $= 416.1$  g fish/day (*i.e.*, 6.93 x 60.03)



and  $1.78Y = \text{grams of mammals consumed}$   
 $= 106.9 \text{ g mammals/day (i.e., } 1.78 \times 60.03)$

Therefore, the total FIR for a generic bald eagle becomes:

$$\text{FIR} = [60 \text{ g birds} + 416 \text{ g fish} + 107 \text{ g mammals}]/\text{day} = 583 \text{ grams wet weight per day.}$$

Exposure frequency represents how much time a bird will spend feeding at a particular lake each year. The bald eagle was assumed to over winter 182 days, stop to feed for as many as 30 days during migration, or would potentially nest and reside year round. A conservative assumption made for these assessments was that bald eagles would spend their entire exposure frequency and duration time feeding solely at each lake or at a combination of all lakes. Exposure duration represents the longevity of each bird species. Longevities were determined through banding data from a variety of sources. Generally, the age of the oldest banded wild bird was used for exposure duration. This provides a conservative estimate of actual longevities in the wild. Averaging time was calculated as 365 days per year multiplied by the exposure duration of the species. Table 2 summarizes the input parameters used in the bald eagle risk assessment.

Reference dose (RfD)	0.021 mg mercury/kg-bw/day
Bald eagle body mass (BW)	5.25 kg (average mass of female)
Food ingestion rate (FIR)	60 g birds/day 107 g mammals/day 416 g fish/day
Exposure duration (ED)	30 days (migratory bald eagle stopover) 182 days (over wintering bald eagle) 365 days (hypothetical nesting bald eagle)
Geometric Mean Re-integrated fish mercury concentrations for various lake combinations	Asaayi Lake (0.07 mg/kg WW) Morgan Lake (0.01 mg/kg WW) Red Lake (0.23 mg/kg WW) Wheatfields Lake (0.06 mg/kg WW) Coldwater lakes (0.06 mg/kg WW) Warmwater lakes (0.05 mg/kg WW) All lakes combined (0.05 mg/kg WW)
Averaging Time	365 days per year for a 30 year lifetime

Using these parameters, the overall dietary concentration of mercury at or below the reference dose (i.e., the Dietary Value [DV]) was calculated using the following equation:

$$\text{DV} = \frac{\text{RfD} \times \text{BW}}{\sum \text{FIR}_i} \quad (\text{Equation 7})$$

Where:

- DV = Dietary Value (mg/kg in the diet that is at or below the Reference Dose)
- RfD = Reference Dose (0.021 mg/kg-bw/day, see above)
- BW = Body Weight (in kg)
- FIR<sub>i</sub> = Food Ingestion Rate (kg food/day), from the i<sup>th</sup> trophic level, for bald eagle

### *Bald Eagle Water Quality Criterion*

Calculation of protective numeric water quality criteria to protect bald eagles through the consumption of fish is based upon a reference dose approach combined with the extent to which mercury becomes concentrated in the fish from specific water bodies (*i.e.*, “wildlife criteria” or WC). The methods used to calculate this criterion are based on those described in the Great Lakes Water Quality Initiative (USEPA 1995a, 1995b, 1995c). When originally implemented in support of the Great Lakes Water Quality Initiative, this approach yielded a single endpoint, which was the total mercury concentration in water protective of wildlife. In this report, an effort was made to update the WC for mercury by calculating its value using data for methylmercury. It should be noted that a methylmercury-based WC could still be related to total mercury residues in fish or water with appropriate conversion factors.

A WC value for mercury was estimated as the ratio of the RfD to an estimated mercury consumption rate in fish (only) referenced to water concentration using a bioaccumulation factor (BAF). A BAF is the ratio of the mercury concentration in fish to its concentration in water. The WC for bald eagles was calculated using the following equation and the BAF from methylmercury measured in water to the concentration of total mercury in re-integrated fish tissue (on a dry weight basis):

$$WC_{\text{ bald eagle}} = \frac{(TD \times UF \times BW)}{D + (FF \times BAF)} \quad (\text{Equation 8})$$

Where:

- WC = Wildlife Criteria for bald eagle (pg/L)
- TD = Tested Dose (mg/kg-bw/day) (*i.e.*, 0.064 mg/kg-bw/day, from above)
- UF = Uncertainty Factor (unitless) (*i.e.*, 0.33, from above)
- BW = Body Weight (*i.e.*, 5.25 kg)
- D = Drinking water intake (L/d)
- FF = Fraction of diet that is fish (all other sources assumed to be negligible)
- BAF = Bioaccumulation Factor (total mercury in fish/methylmercury in water)

## RESULTS AND DISCUSSION

### Limnological Characteristics of the Lakes

Only the average limnological and water quality characteristics of each lake were reviewed. Surface temperature and bottom temperature were strongly correlated ( $r^2=1.00$ ). Of the lakes studied, there were significant correlations between lake size and water temperature ( $r^2=0.92$ ), pH ( $r^2=0.86$ ), specific conductivity ( $r^2=0.79$ ), and including the dissolved Al ( $r^2=0.80$ ), B ( $r^2=0.96$ ), Ca ( $r^2=0.88$ ), Fe ( $r^2=0.88$ ), Mg ( $r^2=0.91$ ), Na ( $r^2=0.98$ ), Se ( $r^2=0.92$ ), S ( $r^2=0.92$ ), and Sr ( $r^2=0.93$ ) in water. Asaayi Lake had the largest surface area-to-depth ratio ( $53 \times 10^6$  m for every meter of depth), Wheatfields and Morgan had a surface area-to-depth ratio of  $6 \times 10^6$ , while Red Lake had a surface area-to-depth ratio of  $0.6 \times 10^6$ . Surface area-to-depth ratio was correlated with dissolved oxygen ( $r^2=0.91$ ) and low turbidity ( $r^2= -0.80$ ) at the lake surface, with the pH ( $r^2=0.90$ ) measured at the bottom, and with mercury dissolved in the water column ( $r^2= -0.81$ ).

### Environmental Contaminants other than Mercury in Water and Fish

The limits of detection for the analytical results reported in this investigation are found in Table 3. Several elements, such as silver, beryllium, cadmium, cobalt, chromium, copper, nickel, and zinc were not detected in any ambient dissolved water sample. Molybdenum was only detected in one dissolved water sample from Morgan Lake, and vanadium was only detected dissolved in two dissolved water samples from Red Lake. Beryllium, cadmium, boron, cobalt, and vanadium were only found in selected fish offal samples from Red Lake or Morgan Lake. These elements (Ag, Be, Cd, Co, Cr, Cu, Ni, and Zn) were not found above the limits of detection with sufficient frequency to further characterize or summarize in this report. All analytical results of lake water, fish tissues, and quality assurance samples, as well as the limnological characteristics of four recreational lakes studied in 2004 for the Navajo Nation Lake Fish and Water Quality Investigation are reported in Table 4. Trout collected from Wheatfields Lake were likely stocked within the previous month as indicated by their uniform condition and size and stock history (E. Benally, Navajo Nation Department of Game and Fish, oral communication, 2004).

### *Trace Elements Dissolved in Water*

The average concentrations of elements found in ambient dissolved lake water samples are reported in Table 5. Several elements (B, Ca, K, Mg, Na, S, Se, and Sr) were found in dissolved Morgan Lake water at concentrations greater than those found at the other lakes. Several elements (Al, Fe, Hg, and P) were found dissolved in Morgan Lake water at concentrations less than found at the other lakes. Morgan Lake is filled with water from the San Juan River; however, concentrations of some elements (B, Ca, K, Na and Sr) were elevated compared to concentrations reported by Goetz and Abeyta (1987) and Ortiz *et al.* (2000) for the San Juan River. The average concentrations of elements dissolved in ambient lake water samples were compared with the numeric criteria of the Navajo Nation Surface Water Quality Standards (Navajo Nation 2004) in Table 5. Only aluminum was routinely detected at concentrations (up to  $236 \mu\text{g/L}$ ) greater than the chronic aquatic habitat criterion

Element Name	Symbol	Method	Limit of Detection <sup>a</sup> in Wet Weight	
			Water (mg/L)	Fish Tissue (mg/kg)
Aluminum	Al	ICP-AES <sup>b</sup>	0.05	1.11
Arsenic	As	ICP-MS <sup>c</sup>	0.0002	0.51
Barium	Ba	ICP-AES	0.001	0.02
Beryllium	Be	ICP-AES	0.005	0.01
Boron	B	ICP-AES	0.01	0.23
Cadmium	Cd	ICP-MS	0.00005	0.01
Calcium	Ca	ICP-AES	0.01	1.11
Chromium	Cr	ICP-AES	0.005	0.11
Cobalt	Co	ICP-AES	0.005	0.11
Copper	Cu	ICP-AES	0.005	0.11
Iron	Fe	ICP-AES	0.01	0.23
Lead	Pb	ICP-MS	0.00005	0.01
Magnesium	Mg	ICP-AES	0.01	0.23
Manganese	Mn	ICP-AES	0.002	0.05
Mercury	Hg	AFS <sup>d</sup>	0.0000005	--- <sup>e</sup>
Mercury	Hg	CVAAS <sup>f</sup>	--- <sup>e</sup>	0.002
Methylmercury	MeHg	AFS	0.000000011	0.002
Molybdenum	Mo	ICP-AES	0.01	0.25
Nickel	Ni	ICP-AES	0.005	0.11
Phosphorus	P	ICP-AES	0.05	1.11
Potassium	K	ICP-AES	0.10	0.01
Selenium	Se	AFS	0.0004	0.02
Silver	Ag	ICP-AES	0.010	--- <sup>e</sup>
Sodium	Na	ICP-AES	2	44
Strontium	Sr	ICP-AES	0.0005	0.01
Sulfur	S	ICP-AES	0.1	2.5
Vanadium	V	ICP-AES	0.01	0.25
Zinc	Zn	ICP-AES	0.005	0.13

<sup>a</sup> = For tissue, limit of detection reported is the highest detection limit for the sample batch.

<sup>b</sup> = Analysis was by inductively coupled plasma - atomic emission spectroscopy.

<sup>c</sup> = Analysis was by inductively coupled plasma - mass spectroscopy.

<sup>d</sup> = Analysis was by atomic fluorescence.

<sup>e</sup> = Sample media was not analyzed using this method.

<sup>f</sup> = Analysis was by cold-vapor atomic absorption spectroscopy.

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Collection Date	Latitude (decimal degrees)	Longitude (decimal degrees)	Average Fish Length (mm)	Average Fish Weight (grams)	Average Surface Water Temperature (Celsius)
T304-2001	Asaayi Lake	unfiltered grab water	31-Mar-2004	35.980917	-108.930194			9.73
T304-2002	Asaayi Lake	unfiltered grab water	31-Mar-2004	35.980917	-108.930194			9.73
T304-2003	Asaayi Lake	unfiltered grab water	31-Mar-2004	35.980917	-108.930194			9.73
T304-2004	Asaayi Lake	unfiltered grab water	31-Mar-2004	35.980917	-108.930194			9.73
T304-2005	Wheatfields Lake	unfiltered grab water	30-Mar-2004	36.206806	-109.097583			9.93
T304-2006	Wheatfields Lake	unfiltered grab water	30-Mar-2004	36.206806	-109.097583			9.93
T304-2007	Wheatfields Lake	unfiltered grab water	30-Mar-2004	36.206806	-109.097583			9.93
T304-2008	Wheatfields Lake	unfiltered grab water	30-Mar-2004	36.206806	-109.097583			9.93
T304-2009	Red Lake	unfiltered grab water	29-Mar-2004	35.919417	-109.040661			10.39
T304-2010	Red Lake	unfiltered grab water	29-Mar-2004	35.919417	-109.040661			10.39
T304-2011	Red Lake	unfiltered grab water	29-Mar-2004	35.919417	-109.040661			10.39
T304-2012		unfiltered Blank meHg water	29-Mar-2004	35.919417	-109.040661			
T304-2013	Morgan Lake	unfiltered grab water	1-Apr-2004	36.702417	-108.474167			23.77
T304-2014	Morgan Lake	unfiltered grab water	1-Apr-2004	36.702417	-108.474167			23.77
T304-2015	Morgan Lake	unfiltered grab water	1-Apr-2004	36.702417	-108.474167			23.77
T304-2016	Morgan Lake	unfiltered grab water	1-Apr-2004	36.702417	-108.474167			23.77
T304-2017	Asaayi Lake	filtered, composited water	31-Mar-2004	35.980917	-108.930194			9.73
T304-2018	Asaayi Lake	filtered, composited water	31-Mar-2004	35.980917	-108.930194			9.73
T304-2019	Asaayi Lake	filtered, Blank Deionized water	31-Mar-2004	35.980917	-108.930194			
T304-2020	Wheatfields Lake	filtered, composited water	30-Mar-2004	36.206806	-109.097583			9.93
T304-2021	Wheatfields Lake	filtered, composited water	30-Mar-2004	36.206806	-109.097583			9.93
T304-2022	Wheatfields Lake	filtered, Blank Deionized water	30-Mar-2004	36.206806	-109.097583			
T304-2023	Red Lake	filtered, composited water	29-Mar-2004	35.919417	-109.040661			10.39
T304-2024	Red Lake	filtered, composited water	29-Mar-2004	35.919417	-109.040661			10.39
T304-2025	Red Lake	filtered, Blank Deionized water	29-Mar-2004	35.919417	-109.040661			
T304-2026	Morgan Lake	filtered, composited water	1-Apr-2004	36.702417	-108.474167			23.77
T304-2027	Morgan Lake	filtered, composited water	1-Apr-2004	36.702417	-108.474167			23.77
T304-2028	Morgan Lake	filtered, Blank Deionized water	1-Apr-2004	36.702417	-108.474167			

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Collection Date	Latitude (decimal degrees)	Longitude (decimal degrees)	Average Fish Length (mm)	Average Fish Weight (grams)	Average Surface Water Temperature (Celsius)
T304-2029	Asaayi Lake	composited skinless trout fillet	31-Mar-2004	35.980917	-108.930194	365.4	588.2	9.73
T304-2030	Asaayi Lake	composited skinless trout fillet	31-Mar-2004	35.980917	-108.930194	309.2	359.8	9.73
T304-2031	Asaayi Lake	composited skinless trout fillet	31-Mar-2004	35.980917	-108.930194	368.8	597.0	9.73
T304-2032	Asaayi Lake	composited skinless trout fillet	31-Mar-2004	35.980917	-108.930194	277.6	255.4	9.73
T304-2033	Wheatfields Lake	composited skinless trout fillet	30-Mar-2004	36.206806	-109.097583	247.4	163.9	9.93
T304-2034	Wheatfields Lake	composited skinless trout fillet	30-Mar-2004	36.206806	-109.097583	231.6	170.2	9.93
T304-2035	Wheatfields Lake	composited skinless trout fillet	30-Mar-2004	36.206806	-109.097583	188.8	83.0	9.93
T304-2036	Wheatfields Lake	composited skinless trout fillet	30-Mar-2004	36.206806	-109.097583	186.6	69.7	9.93
T304-2037	Red Lake	composited skinless catfish fillet	29-Mar-2004	35.919417	-109.040661	452.2	836.2	10.39
T304-2038	Red Lake	composited skinless catfish fillet	29-Mar-2004	35.919417	-109.040661	450.6	790.0	10.39
T304-2039	Red Lake	composited skinless catfish fillet	29-Mar-2004	35.919417	-109.040661	468.0	969.8	10.39
T304-2040	Red Lake	composited skinless catfish fillet	29-Mar-2004	35.919417	-109.040661	454.2	836.2	10.39
T304-2041	Morgan Lake	composited skinless catfish fillet	2-Apr-2004	36.702417	-108.474167	418.4	697.4	23.77
T304-2042	Morgan Lake	composited skinless catfish fillet	2-Apr-2004	36.702417	-108.474167	365.4	405.2	23.77
T304-2043	Morgan Lake	composited skinless bass fillet	1-Apr-2004	36.702417	-108.474167	349.0	734.2	23.77
T304-2044	Morgan Lake	composited skinless catfish fillet	2-Apr-2004	36.702417	-108.474167	407.6	504.8	23.77
T304-2045	Asaayi Lake	composited trout offal	31-Mar-2004	35.980917	-108.930194	365.4	588.2	9.73
T304-2046	Asaayi Lake	composited trout offal	31-Mar-2004	35.980917	-108.930194	309.2	359.8	9.73
T304-2047	Asaayi Lake	composited trout offal	31-Mar-2004	35.980917	-108.930194	368.8	597.0	9.73
T304-2048	Asaayi Lake	composited trout offal	31-Mar-2004	35.980917	-108.930194	277.6	255.4	9.73
T304-2049	Wheatfields Lake	composited trout offal	30-Mar-2004	36.206806	-109.097583	247.4	163.9	9.93
T304-2050	Wheatfields Lake	composited trout offal	30-Mar-2004	36.206806	-109.097583	231.6	170.2	9.93
T304-2051	Wheatfields Lake	composited trout offal	30-Mar-2004	36.206806	-109.097583	188.8	83.0	9.93
T304-2052	Wheatfields Lake	composited trout offal	30-Mar-2004	36.206806	-109.097583	186.6	69.7	9.93
T304-2053	Red Lake	composited catfish offal	29-Mar-2004	35.919417	-109.040661	452.2	836.2	10.39
T304-2054	Red Lake	composited catfish offal	29-Mar-2004	35.919417	-109.040661	450.6	790.0	10.39
T304-2055	Red Lake	composited catfish offal	29-Mar-2004	35.919417	-109.040661	468.0	969.8	10.39
T304-2056	Red Lake	composited catfish offal	29-Mar-2004	35.919417	-109.040661	454.2	836.2	10.39
T304-2057	Morgan Lake	composited catfish offal	2-Apr-2004	36.702417	-108.474167	418.4	697.4	23.77
T304-2058	Morgan Lake	composited catfish offal	2-Apr-2004	36.702417	-108.474167	365.4	405.2	23.77
T304-2059	Morgan Lake	composited bass offal	1-Apr-2004	36.702417	-108.474167	349.0	734.2	23.77
T304-2060	Morgan Lake	composited catfish offal	2-Apr-2004	36.702417	-108.474167	407.6	504.8	23.77

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Collection Date	Latitude (decimal degrees)	Longitude (decimal degrees)	Average Fish Length (mm)	Average Fish Weight (grams)	Average Surface Water Temperature (Celsius)
2945	Asaayi Lake	Re-integrated whole fish composite	31-Mar-2004	35.980917	-108.930194	365.4	588.2	9.73
3046	Asaayi Lake	Re-integrated whole fish composite	31-Mar-2004	35.980917	-108.930194	309.2	359.8	9.73
3147	Asaayi Lake	Re-integrated whole fish composite	31-Mar-2004	35.980917	-108.930194	368.8	597.0	9.73
3248	Asaayi Lake	Re-integrated whole fish composite	31-Mar-2004	35.980917	-108.930194	277.6	255.4	9.73
3349	Wheatfields Lake	Re-integrated whole fish composite	30-Mar-2004	36.206806	-109.097583	247.4	163.9	9.93
3450	Wheatfields Lake	Re-integrated whole fish composite	30-Mar-2004	36.206806	-109.097583	231.6	170.2	9.93
3551	Wheatfields Lake	Re-integrated whole fish composite	30-Mar-2004	36.206806	-109.097583	188.8	83.0	9.93
3652	Wheatfields Lake	Re-integrated whole fish composite	30-Mar-2004	36.206806	-109.097583	186.6	69.7	9.93
3753	Red Lake	Re-integrated whole fish composite	29-Mar-2004	35.919417	-109.040661	452.2	836.2	10.39
3854	Red Lake	Re-integrated whole fish composite	29-Mar-2004	35.919417	-109.040661	450.6	790.0	10.39
3955	Red Lake	Re-integrated whole fish composite	29-Mar-2004	35.919417	-109.040661	468.0	969.8	10.39
4056	Red Lake	Re-integrated whole fish composite	29-Mar-2004	35.919417	-109.040661	454.2	836.2	10.39
4157	Morgan Lake	Re-integrated whole fish composite	2-Apr-2004	36.702417	-108.474167	418.4	697.4	23.77
4258	Morgan Lake	Re-integrated whole fish composite	2-Apr-2004	36.702417	-108.474167	365.4	405.2	23.77
4359	Morgan Lake	Re-integrated whole fish composite	1-Apr-2004	36.702417	-108.474167	349.0	734.2	23.77
4460	Morgan Lake	Re-integrated whole fish composite	2-Apr-2004	36.702417	-108.474167	407.6	504.8	23.77

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Average Surface pH (Standard units)	Average Surface Dissolved Oxygen (mg/L)	Average Surface Specific Conductivity (uS/cm)	Secchi Depth (inches)	Average Surface Turbidity (NTU)	Average Bottom Temperature (Celsius)
T304-2001	Asaayi Lake	unfiltered grab water	8.48	8.58	108.5	34.4	9.6	8.00
T304-2002	Asaayi Lake	unfiltered grab water	8.48	8.58	108.5	34.4	9.6	8.00
T304-2003	Asaayi Lake	unfiltered grab water	8.48	8.58	108.5	34.4	9.6	8.00
T304-2004	Asaayi Lake	unfiltered grab water	8.48	8.58	108.5	34.4	9.6	8.00
T304-2005	Wheatfields Lake	unfiltered grab water	8.37	7.73	282.3	6.4	83.8	9.06
T304-2006	Wheatfields Lake	unfiltered grab water	8.37	7.73	282.3	6.4	83.8	9.06
T304-2007	Wheatfields Lake	unfiltered grab water	8.37	7.73	282.3	6.4	83.8	9.06
T304-2008	Wheatfields Lake	unfiltered grab water	8.37	7.73	282.3	6.4	83.8	9.06
T304-2009	Red Lake	unfiltered grab water	8.76	6.76	3.6	2.1	120.7	9.08
T304-2010	Red Lake	unfiltered grab water	8.76	6.76	3.6	2.1	120.7	9.08
T304-2011	Red Lake	unfiltered grab water	8.76	6.76	3.6	2.1	120.7	9.08
T304-2012		unfiltered Blank meHg water						
T304-2013	Morgan Lake	unfiltered grab water	8.83	7.85	854.8	34.4	5.5	21.19
T304-2014	Morgan Lake	unfiltered grab water	8.83	7.85	854.8	34.4	5.5	21.19
T304-2015	Morgan Lake	unfiltered grab water	8.83	7.85	854.8	34.4	5.5	21.19
T304-2016	Morgan Lake	unfiltered grab water	8.83	7.85	854.8	34.4	5.5	21.19
T304-2017	Asaayi Lake	filtered, composited water	8.48	8.58	108.5	34.4	9.6	8.00
T304-2018	Asaayi Lake	filtered, composited water	8.48	8.58	108.5	34.4	9.6	8.00
T304-2019	Asaayi Lake	filtered, Blank Deionized water						
T304-2020	Wheatfields Lake	filtered, composited water	8.37	7.73	282.3	6.4	83.8	9.06
T304-2021	Wheatfields Lake	filtered, composited water	8.37	7.73	282.3	6.4	83.8	9.06
T304-2022	Wheatfields Lake	filtered, Blank Deionized water						
T304-2023	Red Lake	filtered, composited water	8.76	6.76	3.6	2.1	120.7	9.08
T304-2024	Red Lake	filtered, composited water	8.76	6.76	3.6	2.1	120.7	9.08
T304-2025	Red Lake	filtered, Blank Deionized water						
T304-2026	Morgan Lake	filtered, composited water	8.83	7.85	854.8	34.4	5.5	21.19
T304-2027	Morgan Lake	filtered, composited water	8.83	7.85	854.8	34.4	5.5	21.19
T304-2028	Morgan Lake	filtered, Blank Deionized water						



Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Average Surface pH (Standard units)	Average Surface Dissolved Oxygen (mg/L)	Average Surface Specific Conductivity (uS/cm)	Secchi Depth (inches)	Average Surface Turbidity (NTU)	Average Bottom Temperature (Celsius)
T304-2029	Asaayi Lake	composited skinless trout fillet	8.48	8.58	108.5	34.4	9.6	8.00
T304-2030	Asaayi Lake	composited skinless trout fillet	8.48	8.58	108.5	34.4	9.6	8.00
T304-2031	Asaayi Lake	composited skinless trout fillet	8.48	8.58	108.5	34.4	9.6	8.00
T304-2032	Asaayi Lake	composited skinless trout fillet	8.48	8.58	108.5	34.4	9.6	8.00
T304-2033	Wheatfields Lake	composited skinless trout fillet	8.37	7.73	282.3	6.4	83.8	9.06
T304-2034	Wheatfields Lake	composited skinless trout fillet	8.37	7.73	282.3	6.4	83.8	9.06
T304-2035	Wheatfields Lake	composited skinless trout fillet	8.37	7.73	282.3	6.4	83.8	9.06
T304-2036	Wheatfields Lake	composited skinless trout fillet	8.37	7.73	282.3	6.4	83.8	9.06
T304-2037	Red Lake	composited skinless catfish fillet	8.76	6.76	3.6	2.1	120.7	9.08
T304-2038	Red Lake	composited skinless catfish fillet	8.76	6.76	3.6	2.1	120.7	9.08
T304-2039	Red Lake	composited skinless catfish fillet	8.76	6.76	3.6	2.1	120.7	9.08
T304-2040	Red Lake	composited skinless catfish fillet	8.76	6.76	3.6	2.1	120.7	9.08
T304-2041	Morgan Lake	composited skinless catfish fillet	8.83	7.85	854.8	34.4	5.5	21.19
T304-2042	Morgan Lake	composited skinless catfish fillet	8.83	7.85	854.8	34.4	5.5	21.19
T304-2043	Morgan Lake	composited skinless bass fillet	8.83	7.85	854.8	34.4	5.5	21.19
T304-2044	Morgan Lake	composited skinless catfish fillet	8.83	7.85	854.8	34.4	5.5	21.19
T304-2045	Asaayi Lake	composited trout offal	8.48	8.58	108.5	34.4	9.6	8.00
T304-2046	Asaayi Lake	composited trout offal	8.48	8.58	108.5	34.4	9.6	8.00
T304-2047	Asaayi Lake	composited trout offal	8.48	8.58	108.5	34.4	9.6	8.00
T304-2048	Asaayi Lake	composited trout offal	8.48	8.58	108.5	34.4	9.6	8.00
T304-2049	Wheatfields Lake	composited trout offal	8.37	7.73	282.3	6.4	83.8	9.06
T304-2050	Wheatfields Lake	composited trout offal	8.37	7.73	282.3	6.4	83.8	9.06
T304-2051	Wheatfields Lake	composited trout offal	8.37	7.73	282.3	6.4	83.8	9.06
T304-2052	Wheatfields Lake	composited trout offal	8.37	7.73	282.3	6.4	83.8	9.06
T304-2053	Red Lake	composited catfish offal	8.76	6.76	3.6	2.1	120.7	9.08
T304-2054	Red Lake	composited catfish offal	8.76	6.76	3.6	2.1	120.7	9.08
T304-2055	Red Lake	composited catfish offal	8.76	6.76	3.6	2.1	120.7	9.08
T304-2056	Red Lake	composited catfish offal	8.76	6.76	3.6	2.1	120.7	9.08
T304-2057	Morgan Lake	composited catfish offal	8.83	7.85	854.8	34.4	5.5	21.19
T304-2058	Morgan Lake	composited catfish offal	8.83	7.85	854.8	34.4	5.5	21.19
T304-2059	Morgan Lake	composited bass offal	8.83	7.85	854.8	34.4	5.5	21.19
T304-2060	Morgan Lake	composited catfish offal	8.83	7.85	854.8	34.4	5.5	21.19

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Average Surface pH (Standard units)	Average Surface Dissolved Oxygen (mg/L)	Average Surface Specific Conductivity (uS/cm)	Secchi Depth (inches)	Average Surface Turbidity (NTU)	Average Bottom Temperature (Celsius)
2945	Asaayi Lake	Re-integrated whole fish composite	8.48	8.58	108.5	34.4	9.6	8.00
3046	Asaayi Lake	Re-integrated whole fish composite	8.48	8.58	108.5	34.4	9.6	8.00
3147	Asaayi Lake	Re-integrated whole fish composite	8.48	8.58	108.5	34.4	9.6	8.00
3248	Asaayi Lake	Re-integrated whole fish composite	8.48	8.58	108.5	34.4	9.6	8.00
3349	Wheatfields Lake	Re-integrated whole fish composite	8.37	7.73	282.3	6.4	83.8	9.06
3450	Wheatfields Lake	Re-integrated whole fish composite	8.37	7.73	282.3	6.4	83.8	9.06
3551	Wheatfields Lake	Re-integrated whole fish composite	8.37	7.73	282.3	6.4	83.8	9.06
3652	Wheatfields Lake	Re-integrated whole fish composite	8.37	7.73	282.3	6.4	83.8	9.06
3753	Red Lake	Re-integrated whole fish composite	8.76	6.76	3.6	2.1	120.7	9.08
3854	Red Lake	Re-integrated whole fish composite	8.76	6.76	3.6	2.1	120.7	9.08
3955	Red Lake	Re-integrated whole fish composite	8.76	6.76	3.6	2.1	120.7	9.08
4056	Red Lake	Re-integrated whole fish composite	8.76	6.76	3.6	2.1	120.7	9.08
4157	Morgan Lake	Re-integrated whole fish composite	8.83	7.85	854.8	34.4	5.5	21.19
4258	Morgan Lake	Re-integrated whole fish composite	8.83	7.85	854.8	34.4	5.5	21.19
4359	Morgan Lake	Re-integrated whole fish composite	8.83	7.85	854.8	34.4	5.5	21.19
4460	Morgan Lake	Re-integrated whole fish composite	8.83	7.85	854.8	34.4	5.5	21.19

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Average Bottom pH (Standard Units)	Average Bottom Dissolved Oxygen (mg/L)	Sample Weight (grams)	Moisture Content (%)	Ag	Al	As	B	Ba
T304-2001	Asaayi Lake	unfiltered grab water	8.21	8.15	500	100	<0.01	0.143	0.0021	0.01	0.049
T304-2002	Asaayi Lake	unfiltered grab water	8.21	8.15	500	100	<0.01	0.153	0.0031	0.01	0.05
T304-2003	Asaayi Lake	unfiltered grab water	8.21	8.15	500	100	<0.01	<0.05	<0.0002	<0.01	<0.001
T304-2004	Asaayi Lake	unfiltered grab water	8.21	8.15	500	100	<0.01	0.127	0.0056	0.08	0.16
T304-2005	Wheatfields Lake	unfiltered grab water	8.37	7.39	500	100	<0.01	0.236	0.0056	0.08	0.168
T304-2006	Wheatfields Lake	unfiltered grab water	8.37	7.39	500	100	<0.01	<0.05	<0.0002	<0.01	<0.001
T304-2007	Wheatfields Lake	unfiltered grab water	8.37	7.39	500	100	<0.01	0.08	0.0082	0.13	0.09
T304-2008	Wheatfields Lake	unfiltered grab water	8.37	7.39	500	100	<0.01	0.166	0.0083	0.134	0.093
T304-2009	Red Lake	unfiltered grab water	8.79	6.68	500	100	<0.01	<0.05	<0.0002	<0.01	<0.001
T304-2010	Red Lake	unfiltered grab water	8.79	6.68	500	100	<0.01	<0.05	0.0063	0.634	0.132
T304-2011	Red Lake	unfiltered grab water	8.79	6.68	500	100	<0.01	<0.05	0.0059	0.652	0.136
T304-2012	Red Lake	unfiltered grab water	8.79	6.68	500	100	<0.01	<0.05	<0.0002	<0.01	<0.001
T304-2013	Morgan Lake	unfiltered Blank meHg water	8.51	5.83	500	100					
T304-2014	Morgan Lake	unfiltered grab water	8.51	5.83	500	100					
T304-2015	Morgan Lake	unfiltered grab water	8.51	5.83	500	100					
T304-2016	Morgan Lake	unfiltered grab water	8.51	5.83	500	100					
T304-2017	Asaayi Lake	filtered, composited water	8.21	8.15	500	100	<0.01	0.143	0.0021	0.01	0.049
T304-2018	Asaayi Lake	filtered, composited water	8.21	8.15	500	100	<0.01	0.153	0.0031	0.01	0.05
T304-2019	Asaayi Lake	filtered, Blank Deionized water	8.21	8.15	500	100	<0.01	<0.05	<0.0002	<0.01	<0.001
T304-2020	Wheatfields Lake	filtered, composited water	8.37	7.39	500	100	<0.01	0.127	0.0056	0.08	0.16
T304-2021	Wheatfields Lake	filtered, composited water	8.37	7.39	500	100	<0.01	0.236	0.0056	0.08	0.168
T304-2022	Wheatfields Lake	filtered, Blank Deionized water	8.37	7.39	500	100	<0.01	<0.05	<0.0002	<0.01	<0.001
T304-2023	Red Lake	filtered, composited water	8.79	6.68	500	100	<0.01	0.08	0.0082	0.13	0.09
T304-2024	Red Lake	filtered, composited water	8.79	6.68	500	100	<0.01	0.166	0.0083	0.134	0.093
T304-2025	Red Lake	filtered, Blank Deionized water	8.79	6.68	500	100	<0.01	<0.05	<0.0002	<0.01	<0.001
T304-2026	Morgan Lake	filtered, composited water	8.51	5.83	500	100	<0.01	<0.05	0.0063	0.634	0.132
T304-2027	Morgan Lake	filtered, composited water	8.51	5.83	500	100	<0.01	<0.05	0.0059	0.652	0.136
T304-2028	Morgan Lake	filtered, Blank Deionized water	8.51	5.83	500	100	<0.01	<0.05	<0.0002	<0.01	<0.001

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Average Bottom pH (Standard Units)	Average Bottom Dissolved Oxygen (mg/L)	Sample Weight (grams)	Moisture Content (%)	Ag	Al	As	B	Ba
T304-2029	Asaayi Lake	composited skinless trout fillet	8.21	8.15	310	77.3		<4.71	0.363	<0.942	<0.094
T304-2030	Asaayi Lake	composited skinless trout fillet	8.21	8.15	206	77.6		<4.97	0.483	<0.994	<0.099
T304-2031	Asaayi Lake	composited skinless trout fillet	8.21	8.15	276	77.9		<4.56	0.301	<0.912	<0.091
T304-2032	Asaayi Lake	composited skinless trout fillet	8.21	8.15	158	78.1		56.7	0.667	<0.978	0.176
T304-2033	Wheatfields Lake	composited skinless trout fillet	8.37	7.39	118	79.5		<4.85	1.01	<0.969	0.301
T304-2034	Wheatfields Lake	composited skinless trout fillet	8.37	7.39	134	78.4		<4.68	0.83	<0.937	0.806
T304-2035	Wheatfields Lake	composited skinless trout fillet	8.37	7.39	60	77.3		10.5	1.03	<0.959	0.269
T304-2036	Wheatfields Lake	composited skinless trout fillet	8.37	7.39	57	77.2		<4.80	0.511	<0.959	0.115
T304-2037	Red Lake	composited skinless catfish fillet	8.79	6.68	214	82.5		<4.68	0.295	<0.937	0.487
T304-2038	Red Lake	composited skinless catfish fillet	8.79	6.68	220	81.6		5.68	0.335	<0.947	0.275
T304-2039	Red Lake	composited skinless catfish fillet	8.79	6.68	314	81.3		21.7	0.279	<0.965	0.193
T304-2040	Red Lake	composited skinless catfish fillet	8.79	6.68	306	81.2		12.9	0.255	<0.955	0.344
T304-2041	Morgan Lake	composited skinless catfish fillet	8.51	5.83	308	81.4		7.9	<0.194	<0.968	0.135
T304-2042	Morgan Lake	composited skinless catfish fillet	8.51	5.83	202	82.9		5.62	0.216	1.01	0.112
T304-2043	Morgan Lake	composited skinless bass fillet	8.51	5.83	422	80.0		<4.83	0.408	<0.967	<0.097
T304-2044	Morgan Lake	composited skinless catfish fillet	8.51	5.83	238	83.3		19.7	0.246	<0.942	0.697
T304-2045	Asaayi Lake	composited trout offal	8.21	8.15	2652	73.5		26.8	<1.85	<0.924	9.24
T304-2046	Asaayi Lake	composited trout offal	8.21	8.15	1634	72.1		30.3	<1.82	<0.912	4.4
T304-2047	Asaayi Lake	composited trout offal	8.21	8.15	2793	74.7		33.4	<1.87	<0.935	14.2
T304-2048	Asaayi Lake	composited trout offal	8.21	8.15	1166	76.0		22.4	<1.89	<0.945	4.07
T304-2049	Wheatfields Lake	composited trout offal	8.37	7.39	726	77.7		22.5	<1.85	<0.924	6.92
T304-2050	Wheatfields Lake	composited trout offal	8.37	7.39	764	75.6		18.5	<1.85	<0.923	10.3
T304-2051	Wheatfields Lake	composited trout offal	8.37	7.39	382	73.9		24	1.49	<0.726	6.19
T304-2052	Wheatfields Lake	composited trout offal	8.37	7.39	320	74.2		12.5	<1.86	<0.929	4.49
T304-2053	Red Lake	composited catfish offal	8.79	6.68	4100	80.0		153	<1.81	3.770	129
T304-2054	Red Lake	composited catfish offal	8.79	6.68	4280	72.9		510	1.33	5.760	46.9
T304-2055	Red Lake	composited catfish offal	8.79	6.68	4860	77.5		65.1	<1.63	1.300	52.1
T304-2056	Red Lake	composited catfish offal	8.79	6.68	3996	76.0		70.1	<1.83	0.946	27.1
T304-2057	Morgan Lake	composited catfish offal	8.51	5.83	2904	70.7		17.4	<1.24	0.971	4.01
T304-2058	Morgan Lake	composited catfish offal	8.51	5.83	1632	76.8		49.7	<1.65	1.950	6.13
T304-2059	Morgan Lake	composited bass offal	8.51	5.83	3044	71.6		31	<1.34	<0.67	3.24
T304-2060	Morgan Lake	composited catfish offal	8.51	5.83	2114	75.1		49.7	<1.51	1.470	2.64

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Average Bottom pH (Standard Units)	Average Bottom Dissolved Oxygen (mg/L)	Sample Weight (grams)	Moisture Content (%)	Ag	Al	As	B	Ba
2945	Asaayi Lake	Re-integrated whole fish composite	8.21	8.15	2962	73.9		24.2	0.866	<0.46	8.278
3046	Asaayi Lake	Re-integrated whole fish composite	8.21	8.15	1840	72.7		27.2	0.862	<0.46	3.913
3147	Asaayi Lake	Re-integrated whole fish composite	8.21	8.15	3069	75.0		30.6	0.878	<0.46	12.927
3248	Asaayi Lake	Re-integrated whole fish composite	8.21	8.15	1324	76.3		26.5	0.912	<0.47	3.605
3349	Wheatfields Lake	Re-integrated whole fish composite	8.37	7.39	844	78.0		19.7	0.937	<0.47	5.995
3450	Wheatfields Lake	Re-integrated whole fish composite	8.37	7.39	898	76.0		16.1	0.911	<0.46	8.883
3551	Wheatfields Lake	Re-integrated whole fish composite	8.37	7.39	441.8	74.4		22.2	1.428	<0.38	5.389
3652	Wheatfields Lake	Re-integrated whole fish composite	8.37	7.39	377.0	74.7		11.0	0.867	<0.47	3.828
3753	Red Lake	Re-integrated whole fish composite	8.79	6.68	4314	80.1		145.5	0.875	3.606	122.625
3854	Red Lake	Re-integrated whole fish composite	8.79	6.68	4500	73.3		485.3	1.281	5.502	44.621
3955	Red Lake	Re-integrated whole fish composite	8.79	6.68	5174	77.7		62.5	0.782	1.250	48.950
4056	Red Lake	Re-integrated whole fish composite	8.79	6.68	4302	76.4		66.0	0.868	0.913	25.197
4157	Morgan Lake	Re-integrated whole fish composite	8.51	5.83	3212	71.7		16.5	0.570	0.920	3.638
4258	Morgan Lake	Re-integrated whole fish composite	8.51	5.83	1834	77.5		44.8	0.758	1.850	5.467
4359	Morgan Lake	Re-integrated whole fish composite	8.51	5.83	3466	72.6		27.5	0.638	<0.35	2.851
4460	Morgan Lake	Re-integrated whole fish composite	8.51	5.83	2352	75.9		46.7	0.703	1.370	2.443

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Be	Ca	Cd	Co	Cr	Cu	Fe	Hg	K	MeHg
T304-2001	Asaayi Lake	unfiltered grab water	<0.0005	23.8	<0.00005	<0.005	<0.005	<0.005	0.08	0.00000190	1.46	0.000000058
T304-2002	Asaayi Lake	unfiltered grab water	<0.0005	24.3	<0.00005	<0.005	<0.005	<0.005	0.08	0.00000189	1.48	0.000000071
T304-2003	Asaayi Lake	unfiltered grab water	<0.0005	0.04	<0.00005	<0.005	<0.005	<0.005	<0.01	<0.00000050	<0.10	0.000000077
T304-2004	Asaayi Lake	unfiltered grab water	<0.0005	29.6	<0.00005	<0.005	<0.005	<0.005	0.05	0.00000178	2.68	0.000000088
T304-2005	Wheatfields Lake	unfiltered grab water	<0.0005	30.9	<0.00005	<0.005	<0.005	<0.005	0.09	0.00000178	2.8	0.000000134
T304-2006	Wheatfields Lake	unfiltered grab water	<0.0005	0.08	<0.00005	<0.005	<0.005	<0.005	<0.01	0.00000114	<0.10	0.000000143
T304-2007	Wheatfields Lake	unfiltered grab water	<0.0005	22.7	<0.00005	<0.005	<0.005	<0.005	0.04	0.00000366	4.51	0.000000159
T304-2008	Wheatfields Lake	unfiltered grab water	<0.0005	23	<0.00005	<0.005	<0.005	<0.005	0.08	0.00000359	4.57	0.000000159
T304-2009	Red Lake	unfiltered grab water	<0.0005	0.03	<0.00005	<0.005	<0.005	<0.005	0.01	<0.00000050	<0.10	0.000000418
T304-2010	Red Lake	unfiltered grab water	<0.0005	98.4	<0.00005	<0.005	<0.005	<0.005	<0.01	<0.00000050	7.28	0.000000152
T304-2011	Red Lake	unfiltered grab water	<0.0005	101	<0.00005	<0.005	<0.005	<0.005	<0.01	<0.00000050	7.49	0.000000068
T304-2012	Red Lake	unfiltered grab water	<0.0005	0.07	<0.00005	<0.005	<0.005	<0.005	<0.01	<0.00000050	<0.10	<0.000000011
T304-2013	Morgan Lake	unfiltered Blank meHg water										0.000000037
T304-2014	Morgan Lake	unfiltered grab water										0.000000017
T304-2015	Morgan Lake	unfiltered grab water										0.000000023
T304-2016	Morgan Lake	unfiltered grab water										0.000000020
T304-2017	Asaayi Lake	filtered, composited water	<0.0005	23.8	<0.00005	<0.005	<0.005	<0.005	0.08	0.00000190	1.46	
T304-2018	Asaayi Lake	filtered, composited water	<0.0005	24.3	<0.00005	<0.005	<0.005	<0.005	0.08	0.00000189	1.48	
T304-2019	Asaayi Lake	filtered, Blank Deionized water	<0.0005	0.04	<0.00005	<0.005	<0.005	<0.005	<0.01	<0.00000050	<0.10	
T304-2020	Wheatfields Lake	filtered, composited water	<0.0005	29.6	<0.00005	<0.005	<0.005	<0.005	0.05	0.00000178	2.68	
T304-2021	Wheatfields Lake	filtered, composited water	<0.0005	30.9	<0.00005	<0.005	<0.005	<0.005	0.09	0.00000178	2.8	
T304-2022	Wheatfields Lake	filtered, Blank Deionized water	<0.0005	0.08	<0.00005	<0.005	<0.005	<0.005	<0.01	0.00000114	<0.10	
T304-2023	Red Lake	filtered, composited water	<0.0005	22.7	<0.00005	<0.005	<0.005	<0.005	0.04	0.00000366	4.51	
T304-2024	Red Lake	filtered, composited water	<0.0005	23	<0.00005	<0.005	<0.005	<0.005	0.08	0.00000359	4.57	
T304-2025	Red Lake	filtered, Blank Deionized water	<0.0005	0.03	<0.00005	<0.005	<0.005	<0.005	0.01	<0.00000050	<0.10	
T304-2026	Morgan Lake	filtered, composited water	<0.0005	98.4	<0.00005	<0.005	<0.005	<0.005	<0.01	<0.00000050	7.28	
T304-2027	Morgan Lake	filtered, composited water	<0.0005	101	<0.00005	<0.005	<0.005	<0.005	<0.01	<0.00000050	7.49	
T304-2028	Morgan Lake	filtered, Blank Deionized water	<0.0005	0.07	<0.00005	<0.005	<0.005	<0.005	<0.01	<0.00000050	<0.10	

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Be	Ca	Cd	Co	Cr	Cu	Fe	Hg	K	MeHg
T304-2029	Asaayi Lake	composited skinless trout fillet	<0.0471	679	<0.0471	<0.471	<0.471	1.5	14.6	0.489	15200	0.482000000
T304-2030	Asaayi Lake	composited skinless trout fillet	<0.0497	828	<0.0497	<0.497	<0.497	1.22	18.7	0.327	15900	0.293000000
T304-2031	Asaayi Lake	composited skinless trout fillet	<0.0456	588	<0.0456	<0.456	<0.456	1.56	17.2	0.593	15300	0.311000000
T304-2032	Asaayi Lake	composited skinless trout fillet	<0.0489	1450	<0.0489	<0.489	<0.489	1.57	16.7	0.184	15900	0.137000000
T304-2033	Wheatfields Lake	composited skinless trout fillet	<0.0485	1820	<0.0485	<0.485	<0.485	1.52	29.4	0.277	17400	0.292000000
T304-2034	Wheatfields Lake	composited skinless trout fillet	<0.0468	3090	<0.0468	<0.468	<0.468	1.72	15.1	0.297	16200	0.343000000
T304-2035	Wheatfields Lake	composited skinless trout fillet	<0.0480	1940	<0.0480	<0.480	<0.480	1.82	12.4	0.332	15600	0.380000000
T304-2036	Wheatfields Lake	composited skinless trout fillet	<0.0480	818	<0.0480	<0.480	<0.480	1.3	11.6	0.392	15600	0.417000000
T304-2037	Red Lake	composited skinless catfish fillet	<0.0468	373	<0.0468	<0.468	<0.468	1.1	20.8	2.53	16300	2.690000000
T304-2038	Red Lake	composited skinless catfish fillet	<0.0474	409	<0.0474	<0.474	<0.474	1.05	22.4	1.74	15100	1.640000000
T304-2039	Red Lake	composited skinless catfish fillet	<0.0483	358	<0.0483	<0.483	<0.483	1.17	16	2.12	15800	2.440000000
T304-2040	Red Lake	composited skinless catfish fillet	<0.0477	481	<0.0477	<0.477	<0.477	1.18	27.4	1.96	15800	1.870000000
T304-2041	Morgan Lake	composited skinless catfish fillet	<0.0484	559	<0.0484	<0.484	<0.484	1.25	16.5	0.0571	14900	0.059100000
T304-2042	Morgan Lake	composited skinless catfish fillet	<0.0467	467	<0.0467	<0.467	<0.467	1.33	15.4	0.0437	15900	0.034900000
T304-2043	Morgan Lake	composited skinless bass fillet	<0.0483	1210	<0.0483	<0.483	<0.483	1.08	8.49	0.0774	14700	0.081600000
T304-2044	Morgan Lake	composited skinless catfish fillet	<0.0471	844	<0.0471	<0.471	<0.471	1.34	37.6	0.0488	16900	0.044100000
T304-2045	Asaayi Lake	composited trout offal	<0.0462	22200	<0.0462	<0.462	0.709	2.87	117	0.343	10000	
T304-2046	Asaayi Lake	composited trout offal	<0.0456	22800	<0.0456	<0.456	0.819	3.86	93.9	0.243	9600	
T304-2047	Asaayi Lake	composited trout offal	<0.0467	28900	<0.0467	<0.467	<0.467	4.14	147	0.414	10300	
T304-2048	Asaayi Lake	composited trout offal	<0.0472	24500	<0.0472	<0.472	0.584	2.71	95.5	0.137	11200	
T304-2049	Wheatfields Lake	composited trout offal	<0.0462	31200	<0.0462	<0.462	<0.462	2.66	67.5	0.201	12100	
T304-2050	Wheatfields Lake	composited trout offal	<0.0461	25300	<0.0461	<0.461	0.48	3.6	62.9	0.213	11100	
T304-2051	Wheatfields Lake	composited trout offal	<0.0363	28300	<0.0363	<0.363	0.857	2.94	55.9	0.252	10100	
T304-2052	Wheatfields Lake	composited trout offal	<0.0465	22200	<0.0465	<0.465	1.16	2.28	46.7	0.262	10400	
T304-2053	Red Lake	composited catfish offal	<0.0453	51800	0.597	0.701	2.66	2.17	243	1.08	9440	
T304-2054	Red Lake	composited catfish offal	0.0387	47700	0.092	<0.327	2.18	2.92	466	0.783	7550	
T304-2055	Red Lake	composited catfish offal	0.0409	89500	0.048	<0.409	2.25	1.45	141	1.16	10000	
T304-2056	Red Lake	composited catfish offal	<0.0457	65500	0.054	<0.457	2.31	2.04	184	0.773	8030	
T304-2057	Morgan Lake	composited catfish offal	<0.0310	55700	<0.0310	<0.310	2.68	1.3	74	0.0241	7320	
T304-2058	Morgan Lake	composited catfish offal	<0.0413	65300	<0.0413	<0.413	1.77	2.14	139	0.0215	9830	
T304-2059	Morgan Lake	composited bass offal	0.0339	83800	0.037	<0.336	4.70	1.32	78.3	0.0358	8820	
T304-2060	Morgan Lake	composited catfish offal	<0.0377	34400	<0.0377	<0.377	1.54	1.8	113	0.0195	7500	

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Be	Ca	Cd	Co	Cr	Cu	Fe	Hg	K	MeHg
2945	Asaayi Lake	Re-integrated whole fish composite	<0.023	19948	<0.023	<0.23	0.659	2.7	106.3	0.358	10544	
3046	Asaayi Lake	Re-integrated whole fish composite	<0.023	20340	<0.023	<0.23	0.755	3.6	85.5	0.252	10305	
3147	Asaayi Lake	Re-integrated whole fish composite	<0.023	26354	<0.023	<0.23	<0.233	3.9	135.3	0.430	10750	
3248	Asaayi Lake	Re-integrated whole fish composite	<0.024	21749	<0.024	<0.23	0.543	2.6	86.1	0.143	11761	
3349	Wheatfields Lake	Re-integrated whole fish composite	<0.023	27092	<0.023	<0.23	<0.233	2.5	62.2	0.212	12841	
3450	Wheatfields Lake	Re-integrated whole fish composite	<0.023	21986	<0.023	<0.23	0.443	3.3	55.8	0.226	11861	
3551	Wheatfields Lake	Re-integrated whole fish composite	<0.019	24735	0.019	<0.19	0.774	2.8	50.0	0.263	10844	
3652	Wheatfields Lake	Re-integrated whole fish composite	<0.023	18965	<0.023	<0.23	1.021	2.1	41.4	0.282	11187	
3753	Red Lake	Re-integrated whole fish composite	<0.023	49249	0.569	0.678	2.540	2.1	232.0	1.152	9780	
3854	Red Lake	Re-integrated whole fish composite	0.038	45388	0.089	<0.17	2.085	2.8	444.3	0.830	7919	
3955	Red Lake	Re-integrated whole fish composite	0.040	84090	0.047	<0.21	2.128	1.4	133.4	1.218	10352	
4056	Red Lake	Re-integrated whole fish composite	<0.023	60875	0.052	<0.23	2.163	2.0	172.9	0.857	8583	
4157	Morgan Lake	Re-integrated whole fish composite	<0.016	50413	<0.016	<0.17	2.446	1.3	68.5	0.027	8047	
4258	Morgan Lake	Re-integrated whole fish composite	<0.021	58159	<0.021	<0.21	1.601	2.1	125.4	0.024	10499	
4359	Morgan Lake	Re-integrated whole fish composite	0.033	73744	<0.035	<0.19	4.157	1.3	69.8	0.041	9536	
4460	Morgan Lake	Re-integrated whole fish composite	<0.019	31004	<0.019	<0.19	1.408	1.8	105.4	0.022	8451	



Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Mg	Min	Mo	Na	Ni	P	Pb	S	Se	Sr	V	Zn
T304-2001	Asaayi Lake	unfiltered grab water	2.87	0.011	<0.01	3	<0.005	0.08	0.00018	1.5	0.00048	0.12	<0.01	<0.005
T304-2002	Asaayi Lake	unfiltered grab water	2.88	0.012	<0.01	4	<0.005	0.08	0.00014	1.6	0.00046	0.125	<0.01	<0.005
T304-2003	Asaayi Lake	unfiltered grab water	<0.01	<0.002	<0.01	<2	<0.005	<0.05	0.00007	<0.1	<0.00040	<0.00005	<0.01	0.026
T304-2004	Asaayi Lake	unfiltered grab water	13.6	0.003	<0.01	31.4	<0.005	0.125	0.00009	6.2	0.00045	0.414	<0.01	<0.005
T304-2005	Wheatfields Lake	unfiltered grab water	14.1	0.004	<0.01	33.6	<0.005	0.151	0.00010	6.5	0.00047	0.438	<0.01	<0.005
T304-2006	Wheatfields Lake	unfiltered grab water	0.01	<0.002	<0.01	<2	<0.005	<0.05	0.00005	<0.1	<0.00040	<0.00005	<0.01	<0.005
T304-2007	Wheatfields Lake	unfiltered grab water	9.29	<0.002	<0.01	65	<0.005	0.103	0.00007	10.7	0.00052	0.339	0.01	<0.005
T304-2008	Wheatfields Lake	unfiltered grab water	9.39	0.002	<0.01	66	<0.005	0.112	0.00009	11	0.00052	0.343	0.01	<0.005
T304-2009	Red Lake	unfiltered grab water	<0.01	<0.002	<0.01	<2	<0.005	<0.05	<0.00005	<0.1	<0.00040	<0.00005	<0.01	0.008
T304-2010	Red Lake	unfiltered grab water	35.8	<0.002	<0.01	99.9	<0.005	<0.05	0.00015	138	0.00128	1.66	<0.01	<0.005
T304-2011	Red Lake	unfiltered grab water	36.7	<0.002	0.01	106	<0.005	<0.05	0.00015	141	0.0013	1.76	<0.01	<0.005
T304-2012	Blank meHg water	unfiltered grab water	0.02	<0.002	<0.01	<2	<0.005	<0.05	<0.00005	<0.1	0.00042	0.0008	<0.01	0.021
T304-2013	Morgan Lake	unfiltered grab water												
T304-2014	Morgan Lake	unfiltered grab water												
T304-2015	Morgan Lake	unfiltered grab water												
T304-2016	Morgan Lake	unfiltered grab water												
T304-2017	Asaayi Lake	filtered, composited water												
T304-2018	Asaayi Lake	filtered, composited water												
T304-2019	Asaayi Lake	filtered, Blank Deionized water												
T304-2020	Wheatfields Lake	filtered, composited water												
T304-2021	Wheatfields Lake	filtered, composited water												
T304-2022	Wheatfields Lake	filtered, Blank Deionized water												
T304-2023	Red Lake	filtered, composited water												
T304-2024	Red Lake	filtered, composited water												
T304-2025	Red Lake	filtered, Blank Deionized water												
T304-2026	Morgan Lake	filtered, composited water												
T304-2027	Morgan Lake	filtered, composited water												
T304-2028	Morgan Lake	filtered, Blank Deionized water												

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Mg	Mn	Mo	Na	Ni	P	Pb	S	Se	Sr	V	Zn
T304-2029	Asaayi Lake	composited skinless trout fillet	1160	0.556	<0.942	925	<0.471	11000	0.234	8340	1.25	0.603	<0.942	13.3
T304-2030	Asaayi Lake	composited skinless trout fillet	1220	0.616	<0.994	1130	<0.497	11500	0.215	8430	1.14	0.945	<0.994	16.6
T304-2031	Asaayi Lake	composited skinless trout fillet	1130	0.511	<0.912	1070	<0.456	10900	0.182	8270	1.16	0.511	<0.912	14.8
T304-2032	Asaayi Lake	composited skinless trout fillet	1230	0.841	<0.978	1210	<0.489	11700	0.223	8160	1.23	1.56	<0.978	19.7
T304-2033	Wheatfields Lake	composited skinless trout fillet	1270	0.717	<0.969	1310	<0.485	12300	0.201	8350	1.22	3.57	<0.969	16.3
T304-2034	Wheatfields Lake	composited skinless trout fillet	1270	0.768	<0.937	1240	<0.468	12400	0.183	8400	1.09	7.64	<0.937	16
T304-2035	Wheatfields Lake	composited skinless trout fillet	1290	0.844	<0.959	1190	<0.480	11800	0.188	8220	1.06	3.04	<0.959	15.8
T304-2036	Wheatfields Lake	composited skinless trout fillet	1280	0.652	<0.959	1150	<0.480	11400	0.172	8470	1.17	1.06	<0.959	17.7
T304-2037	Red Lake	composited skinless catfish fillet	1050	0.778	<0.937	1210	<0.468	10500	0.152	9690	0.886	1.74	<0.937	21.9
T304-2038	Red Lake	composited skinless catfish fillet	1040	0.625	<0.947	1810	<0.474	9840	0.159	9320	0.849	1.74	<0.947	21.1
T304-2039	Red Lake	composited skinless catfish fillet	1060	0.531	<0.965	1790	<0.483	10400	0.153	9850	0.881	1.37	<0.965	20.2
T304-2040	Red Lake	composited skinless catfish fillet	1070	0.725	<0.955	1890	<0.477	10400	0.159	9530	0.848	1.82	<0.955	20.7
T304-2041	Morgan Lake	composited skinless catfish fillet	1090	0.735	<0.968	1720	<0.484	9980	0.144	10800	15.1	2.6	<0.968	22.5
T304-2042	Morgan Lake	composited skinless catfish fillet	1110	0.672	<0.933	2060	<0.467	10600	0.121	11600	21	2.38	<0.933	24.2
T304-2043	Morgan Lake	composited skinless bass fillet	1290	0.3	<0.967	1450	<0.483	9980	0.128	10800	17.2	5.63	<0.967	18.8
T304-2044	Morgan Lake	composited skinless catfish fillet	1210	1.47	<0.942	1950	<0.471	11200	0.234	12200	23.1	3.92	<0.942	24.9
T304-2045	Asaayi Lake	composited trout offal	1010	36.4	<0.924	3520	<0.462	17500	0.124	7320	1.39	29.4	<0.924	83
T304-2046	Asaayi Lake	composited trout offal	950	10	<0.912	3310	0.661	17800	0.138	6860	1.38	28.4	<0.912	86.6
T304-2047	Asaayi Lake	composited trout offal	1190	55.4	<0.935	3830	<0.467	20500	0.157	7500	1.28	41.1	<0.935	102
T304-2048	Asaayi Lake	composited trout offal	1110	10.2	<0.945	4060	<0.472	19700	0.11	7470	1.39	31.1	<0.945	102
T304-2049	Wheatfields Lake	composited trout offal	1220	5.47	<0.924	5040	<0.462	22900	0.113	8200	1.42	66.6	<0.924	131
T304-2050	Wheatfields Lake	composited trout offal	1150	5.01	<0.923	4290	<0.461	19700	0.113	7840	1.24	67.6	<0.923	103
T304-2051	Wheatfields Lake	composited trout offal	1240	5.24	<0.726	3520	0.497	22000	0.0841	7660	1.48	53.7	<0.726	87.8
T304-2052	Wheatfields Lake	composited trout offal	1040	3.79	<0.929	3420	0.737	18500	0.0767	7090	1.54	43.7	<0.929	74.9
T304-2053	Red Lake	composited catfish offal	1420	76.9	<0.905	7690	1.84	29500	0.388	6870	1.69	190	1.82	84.4
T304-2054	Red Lake	composited catfish offal	1430	46.7	<0.654	4930	1.37	26100	0.569	6280	1.63	185	2.34	66.7
T304-2055	Red Lake	composited catfish offal	1980	44.7	<0.817	6210	1.08	49100	0.315	6980	1.39	293	1.58	103
T304-2056	Red Lake	composited catfish offal	1510	30.2	<0.915	6300	1.36	36400	0.296	5680	1.6	231	1.48	90.1
T304-2057	Morgan Lake	composited catfish offal	1510	11.6	<0.620	4660	1.3	31900	0.306	6010	8.95	280	1.04	82.5
T304-2058	Morgan Lake	composited catfish offal	1910	26.4	<0.826	6020	1.15	38000	0.331	7880	12.5	338	0.90	127
T304-2059	Morgan Lake	composited bass offal	2270	6.93	<0.672	4820	2.33	45300	0.121	8080	12.1	488	<0.672	64
T304-2060	Morgan Lake	composited catfish offal	1180	5.77	<0.754	5120	0.97	21200	0.208	6780	12	179	<0.754	70.7

Table 4. Sample Information, Analytical Results, and Liminological Characteristics of Four Navajo Nation Lakes.  
 [See Table 3 for abbreviations; all water values are mg/L; all fish values are mg/kg dry weight].

Sample Number	Lake Name	Sample Type	Mg	Mn	Mo	Na	Ni	P	Pb	S	Se	Sr	V	Zn
2945	Asaayi Lake	Re-integrated whole fish composite	1026	32.65	<0.47	3248	<0.23	16820	0.14	7427	1.38	26.39	<0.46	75.71
3046	Asaayi Lake	Re-integrated whole fish composite	980	8.95	<0.47	3066	0.61	17095	0.15	7036	1.35	25.33	<0.46	78.76
3147	Asaayi Lake	Re-integrated whole fish composite	1185	50.46	<0.47	3582	<0.23	19637	0.16	7569	1.27	37.45	<0.47	94.16
3248	Asaayi Lake	Re-integrated whole fish composite	1124	9.08	<0.47	3720	<0.24	18745	0.12	7552	1.37	27.57	<0.47	92.18
3349	Wheatfields Lake	Re-integrated whole fish composite	1227	4.81	<0.47	4519	<0.23	21418	0.13	8221	1.39	57.79	<0.47	114.96
3450	Wheatfields Lake	Re-integrated whole fish composite	1168	4.38	<0.47	3835	<0.23	18611	0.12	7924	1.22	58.65	<0.46	90.02
3551	Wheatfields Lake	Re-integrated whole fish composite	1247	4.65	<0.38	3205	0.46	20620	0.10	7736	1.42	46.85	<0.38	78.06
3652	Wheatfields Lake	Re-integrated whole fish composite	1076	3.32	<0.47	3077	0.66	17426	0.09	7299	1.48	37.25	<0.47	66.25
3753	Red Lake	Re-integrated whole fish composite	1402	73.12	<0.47	7413	1.76	28557	0.38	7010	1.65	180.66	1.75	81.30
3854	Red Lake	Re-integrated whole fish composite	1411	44.45	<0.33	4777	1.31	25305	0.55	6429	1.59	176.04	2.25	64.47
3955	Red Lake	Re-integrated whole fish composite	1924	42.02	<0.41	5942	1.03	46751	0.31	7154	1.36	275.30	1.51	97.98
4056	Red Lake	Re-integrated whole fish composite	1479	28.10	<0.47	5986	1.28	34551	0.29	5954	1.55	214.70	1.41	85.16
4157	Morgan Lake	Re-integrated whole fish composite	1470	10.56	<0.33	4378	1.20	29798	0.29	6469	9.54	253.40	0.99	76.75
4258	Morgan Lake	Re-integrated whole fish composite	1822	23.57	<0.42	5584	1.05	34982	0.31	8290	13.44	301.03	0.85	115.68
4359	Morgan Lake	Re-integrated whole fish composite	2151	6.12	<0.35	4410	2.08	41000	0.12	8411	12.72	429.27	<0.35	58.50
4460	Morgan Lake	Re-integrated whole fish composite	1183	5.33	<0.39	4799	0.90	20188	0.21	7328	13.12	161.28	<0.39	66.07

Table 5. Average Concentration of Elements Dissolved in Lake Water Composites (N=2 from each lake) Compared with Selected Numeric Navajo Nation (2004) Water Quality Criteria for Various Designated Uses.

[Note: for the lead criteria that are dependent on hardness in the calculation, a value of 100 mg/L was used; all values are mg/L unless specified otherwise; **bolded** values may exceed water quality criteria within the same row as identified by *italics*; NNCNS = No Numeric Criteria, Narrative Standard applies.]

Dissolved Element (mg/L) <sup>a</sup>	Asaayi Lake	Wheat-fields Lake	Red Lake	Morgan Lake	Aquatic Habitat Acute	Aquatic Habitat Chronic	Agricultural Water Supply	Domestic Water Supply	Fish Consumption	Livestock & Wildlife Watering	Secondary Contact
Al	<b>0.15</b>	<b>0.18</b>	<b>0.12</b>	0.03	0.75	<i>0.087</i>	5.0	NNCNS	NNCNS	5.0	NNCNS
As	0.003	0.006	0.008	0.006	0.340	0.150	0.100	0.050	1.450	0.020	0.420
B	0.01	0.08	0.13	<b>0.64</b>	NNCNS	NNCNS	0.750	<i>0.630</i>	NNCNS	10.000	126.000
Ba	0.05	0.09	0.09	0.11	NNCNS	NNCNS	NNCNS	1.000	NNCNS	NNCNS	98.000
Ca	24.1	30.3	22.9	99.7	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS
Fe	0.08	0.07	0.06	<0.01	NNCNS	1.000	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS
K	1.5	2.7	4.5	7.4	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS
Total Hg (µg/L)	0.00190	0.00178	0.00363	0.00025	2.4	0.012	NNCNS	2.0	0.15	10.0	420.0
MeHg (µg/L)	0.00007	0.00015	0.00016	0.00002	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS
Mg	2.9	13.9	9.3	36.3	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS
Mn	0.012	0.004	<0.002	<0.002	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS
Na	3.5	32.5	65.5	102.9	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS
P	0.08	0.14	0.11	0.03	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS
Pb (µg/L)	0.16	0.09	0.08	0.15	64.58	2.52	5000.0	15.0	NNCNS	100.0	15.0
S	1.6	6.4	10.9	139.5	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS
Se	<0.001	<0.001	<0.001	0.001	0.020	0.002	0.130	0.050	9.000	0.050	7.000
Sr	0.12	0.43	0.34	1.71	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS	NNCNS

<sup>a</sup> = See Table 3 for element abbreviations, method of analysis, and limits of detection.

of 87µg/L at Asaayi, Red, and Wheatfield Lakes. Hem (1985) reported that in most natural waters aluminum is rarely above a few tenths of a milligram per liter. However, water samples with dissolved aluminum concentrations frequently exceeding the chronic and even acute (750g/L) water quality criteria for aquatic life are frequently reported in the Rio Grande basin (NMWQCC 2000; Buhl 2002; Lusk *et al.* 2002). Hem (1985) reported that some water-borne colloids that are rich in aluminum have small size (<0.10 micrometers in diameter) and can pass through the 0.45 micrometer filter media (as used in this study). In development of the aquatic life criteria for aluminum, research was focused primarily on aquatic systems with low pH (USEPA 1988). However, there may have been an information gap regarding the chemical and biological effects of elevated aluminum to aquatic life similar to those alkaline waters found on the Navajo Nation in the USEPA (1988) criteria document.

Boron was found dissolved in ambient water samples from Morgan Lake (0.64 mg/L) above the boron criteria for a domestic water supply (0.63 mg/L). However, Morgan Lake is not designated as a domestic water supply by the Navajo Nation. Furthermore, the USEPA (1991) reported that boron as low as 1.52 mg/L is not likely to affect aquatic life. Because boron concentrations dissolved in Morgan Lake water are less than this value, boron toxicity would not be expected to pose a significant risk to aquatic life. However, elevated boron in Morgan Lake water may limit its value as a domestic water supply (Table 5).

#### *Trace Elements in Fish Fillet Tissues*

The geometric mean concentrations of trace elements found in fish fillets from lakes sampled are reported in Table 6. Several elements (As, Ca, Cu, P, Mg, K, and P) were found at higher geometric mean concentrations in trout from coldwater lakes than in warmwater fish from Morgan and Red Lakes. In addition, the elements Ba and Na were found at higher concentrations in catfish and bass from warmwater lakes than in coldwater trout from Asaayi and Wheatfields Lakes.

The geometric mean concentrations in fish fillets collected from the Navajo Nation lakes was compared to the geometric mean concentrations in fish fillets collected from the San Juan River by Blanchard *et al.* (1993), Wilson *et al.* (1995), as well as unreported data from Bristol *et al.* (1997) and Simpson and Lusk (2000). Navajo Nation fish fillets had higher geometric mean concentrations of aluminum, arsenic, and strontium than in fillets from the San Juan River but lower geometric mean concentrations of barium and iron.

Concentrations of selenium in fish fillets were elevated in Morgan Lake (3.4 mg/kg WW), and they were greater than the concentration (>1.5 mg/kg WW) associated with human health advisories for adults (USEPA 2000). Therefore, for children whose diet contains a large portion of fish, consuming more than 6, 2-lb catfish per month would likely exceed the USEPA (2000) health effects threshold for selenium. Typically, selenium concentrations in fish fillets are < 0.6 mg/kg WW. Reproductive failure in fisheries has been identified in some species with fillet concentrations as low as 1.7 mg/kg WW (Lemly 1996a, 1996b; USDOJ 1998). Selenium in fish fillets may pose a risk to human health and may be reducing the reproductive success of some Morgan Lake fishes as well as to piscivorous wildlife that resides there and consume a diet consisting mainly of whole fish.

### *Trace Elements in Re-Integrated Fish*

The geometric mean concentrations of trace elements found in re-integrated fish from lakes sampled are reported in Table 7. Several trace elements (Al, Ba, Be, Ca, Cd, Fe, Mn, Pb, and V) were found at higher geometric mean concentrations in catfish from Red Lake than fish from any other lake. Sediment chemistry may play a part in the accumulation of these elements in Red Lake catfish, as it is a shallower, more turbid lake. Simpson and Lusk (2000) reported that an organisms association with sediment, such as benthic algae, aquatic worms, and fish explain over 80% of the accumulation of these elements than found in less turbid river reaches with pelagic organisms. These elements are often associated with soil and sediment and therefore, concentrations in benthic biota may likely reflect the ambient geochemical environment.

The geometric mean concentrations in re-integrated fish collected from the Navajo Nation lakes were compared to the those reported in whole fish collected from the San Juan River (Simpson and Lusk 2000), and to the 85<sup>th</sup> percentile concentration in whole fish collected nationwide (Schmitt and Brumbaugh 1990). Only mercury and selenium in re-integrated fish from the study lakes were above concentrations of concern or were above concentrations typical in fish collected from the San Juan River or collected nationwide.

Catfish collected from Morgan Lake had elevated selenium concentrations (> 12 mg/kg DW). While selenium concentrations in whole fish collected from the San Juan River have ranged from 0.1 to 15.1 mg/kg DW, the composite sample of catfish from Morgan Lake had the highest selenium concentrations (13.4 mg/kg DW) ever reported in the San Juan River Basin. Occasionally, wild catfish captured from the San Juan River are stocked into Morgan Lake (J. Brooks, USFWS, written communication, 2005), so there may be multiple sources of selenium in the tissues of these catfish. However, largemouth bass that were collected were not recently stocked (largemouth bass were stocked in 2002; C. Kitcheyan, USFWS, oral communication, 2005) and these fish contained selenium concentrations as high as 12.7 mg/kg DW.

Nationally, selenium concentrations in whole fish are typically < 2 mg/kg DW (USDOJ 1998). Bluegill that contained selenium concentrations from 4 to 6 mg/kg DW have been reported to have a 10 percent reproductive impairment (Lemly 1996a, 1996b; USDOJ 1998). As selenium concentrations in fish from Morgan Lake fish are much greater than this threshold, Morgan Lake fish may experience periodic reproductive failures that may affect the fishery. Selenium concentrations in whole body fish above 4 mg/kg DW have also been associated reduced growth and higher mortality rates (Lemly 1996a, 1996b; USDOJ 1998). Lemly (1996a, 1996b) and the USDOJ (1998) also reported selenium concentrations greater than 3 mg/kg DW in the diets of predatory species pose reproductive risks to migratory and resident birds such as those that may feed extensively on fish from Morgan Lake. Selenium contamination in Morgan Lake fish may be at a level where it could affect the fishery, as well as pose a health risk to people and wildlife that consume a large amount of fish from Morgan Lake. Sources of selenium to Morgan Lake fish should be identified and reduced if population effects are identified in fish, people or wildlife or if health impacts are observed in people or wildlife that regularly eat fish from this lake.

Element (ug/g Wet Weight) <sup>a</sup>	Asaayi Lake Trout Fillets	Wheat- fields Lake Trout Fillets	Red Lake Catfish Fillets	Morgan Lake Cattfish & Bass Fillets	All Lake Fish Fillets Combined	San Juan River Fillets <sup>b</sup>	No Fish Consumption Recommended (Non-Cancer Endpoint) <sup>c</sup>	No Fish Consumption Recommended (Cancer Endpoint) <sup>c</sup>	General Dietary Level of Concern for Wildlife <sup>d</sup>
Al	1.2	0.8	1.4	1.2	1.1	0.50	---	---	>200
As	0.10	<b>0.18</b>	0.05	0.04	0.08	0.06	> 5.6 <sup>f</sup>	> 0.13 <sup>f</sup>	> 10
B	<0.2	<0.2	<0.2	<0.2	<0.2	<0.20	---	---	> 30
Ba	0.01	0.01	0.06	0.03	0.03	0.07	---	---	> 20
Be	<0.01	<0.01	<0.01	<0.01	<0.01	<0.20	---	---	>0.8
Ca	185.3	378.2	73.6	126.7	161.0	---	---	---	---
Cd	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	> 0.09	---	>0.1
Cr	<0.05	<0.05	<0.05	<0.05	<0.05	0.08	---	---	>5.1
Cu	0.3	0.4	0.2	0.2	0.3	0.37	---	---	> 40
Fe	3.7	3.4	3.9	3.0	3.5	6.42	---	---	>500
Total Hg	0.08	0.07	<b>0.40</b>	0.01	0.07	0.08	---	---	>0.10
MeHg	0.06	0.08	<b>0.39</b>	0.01	0.06	---	> 0.3	---	---
K	3,468	3,541	2,888	2,811	3,160	---	---	---	---
Mg	264	280	194	212	234	287	---	---	>3,000
Min	<0.2	<0.2	<0.2	<0.2	<0.2	<0.25	---	---	>400
Mo	<0.1	<0.1	<0.1	<0.1	<0.1	0.08	---	---	> 10
Na	240	267	348	321	291	---	---	---	---
Ni	<0.1	<0.1	<0.1	<0.1	<0.1	0.03	---	---	> 50
P	2,510	2,618	1,886	1,883	2,198	---	---	---	---
Pb	<0.1	<0.1	<0.1	<0.1	<0.1	<0.04	---	---	> 40
S	1,849	1,829	1,760	2,046	1,868	---	---	---	---
Se	0.27	0.25	0.16	<b>3.4</b>	0.43	0.60	>1.5	---	> 0.8
Sr	0.2	0.7	0.3	0.6	0.4	2.0	---	---	>2,000
V	<0.2	<0.2	<0.2	<0.2	<0.2	<0.1	---	---	> 10
Zn	3.6	3.6	3.9	4.1	3.8	7.4	---	---	> 45

<sup>a</sup> = See Table 3 for element abbreviations, methods of analysis, and limits of detection.

<sup>b</sup> = Based on Blanchard *et al.* 1993; Wilson *et al.* 1995; Thomas *et al.* 1997, 1998; Simpson and Lusk 2000; and on unreported data from Bristol *et al.* (1997).

<sup>c</sup> = Based on USEPA 2000; fish consumption not recommended based cancer or chronic systemic effects using 8 oz meal size and other assumptions.

<sup>d</sup> = Based on NRC 1980; Eisler 1985; Eisler 1986; Eisler 1987; Eisler 1993; Eisler 1994; Eisler 1998.

<sup>e</sup> = Data not available.

Table 7. Comparison of the Geometric Mean Concentrations of Trace Elements in Re-integrated Fish collected from Four Navajo Nation Recreational Lakes to Whole Fish Collected from the San Juan River Basin, Collected Nationwide and General Dietary Concentrations of Concern for Wildlife.

Element (ug/g Dry Weight) <sup>a</sup>	Asaayi Lake Trout	Morgan Lake Catfish and Bass	Red Lake Catfish	Wheatfields Lake Trout	All Catfish	All Trout	Fish From All Lakes	San Juan River Fish <sup>b</sup>	Fish Sampled Nationwide (85 <sup>th</sup> Percentile) <sup>c</sup>	General Dietary Level of Concern for Wildlife <sup>d</sup>
Al	27.0	31.2	130.6	16.7	72.0	21.2	36.8	171.8	---	>1,000
As	0.9	0.7	0.9	1.0	0.8	0.9	0.9	0.3	1.2	>30
B	0.5	1.0	2.1	0.4	1.8	0.5	0.8	1.1	---	>30
Ba	6.2	3.4	51.0	5.8	16.5	6.0	8.9	7.4	---	---
Be	0.02	0.02	0.03	0.02	0.02	0.02	0.02	0.05	---	>3
Ca	21,960	50,884	58,161	22,991	52,087	22,470	34,962	---	---	---
Cd	0.02	0.02	0.11	0.02	0.2	0.2	0.03	0.06	0.28	>0.5
Cr	0.5	2.2	2.2	0.5	2.0	0.5	1.1	0.7	---	>10
Cu	3.1	1.6	2.0	2.7	1.9	2.9	2.3	3.6	5.1	>120
Fe	101	89	221	52	155	73	101	210	---	>1,000
Hg	0.27	0.03	<b>1.00</b>	0.24	0.20	0.26	0.21	0.19	<i>0.81</i>	>0.3
K	10,826	9,084	9,108	11,659	9,035	11,235	10,109	---	---	---
Mg	1,076	1,616	1,540	1,178	1,509	1,126	1,332	1038	---	>10,000
Mn	19.1	9.5	44.3	4.2	24.4	9.0	13.6	16.4	---	>1,000
Mo	0.5	0.4	0.4	0.4	0.4	0.5	0.4	0.4	---	>30
Na	3,394	4,769	5,958	3,615	5,477	3,503	4,321	---	---	---
Ni	0.3	1.2	1.3	0.4	1.2	0.3	0.6	0.5	---	>300
P	18,037	30,471	32,870	19,454	30,502	18,732	24,349	---	---	---
Pb	0.1	0.2	0.4	0.1	0.3	0.1	0.2	0.4	---	>120
S	7,393	7,583	6,619	7,788	6,913	7,588	7,332	---	---	---
Se	1.3	<b>12.1</b>	1.5	1.4	<b>3.7</b>	1.4	2.4	2.6	<i>3.1</i>	>3
Sr	28.8	269.6	208.2	49.3	217.6	27.7	94.5	53.0	---	>5,000
V	0.5	0.6	1.7	0.4	1.2	0.5	0.7	0.6	---	>30
Zn	84.8	76.5	81.3	85.5	82.3	85.2	82.0	82.5	258	>180

<sup>a</sup> = See Table 3 for element abbreviations, methods of analysis, and limits of detection.

<sup>b</sup> = Based on Simpson and Lusk (2000).

<sup>c</sup> = Based on Schmitt and Brumbaugh (1990); 85<sup>th</sup> percentile concentration converted to dry weight using 76.6 % moisture.

<sup>d</sup> = Based on NRC 1980; Eisler 1985; Eisler 1986; Eisler 1987; Eisler 1988; Eisler 1993; Eisler 1994; Eisler 1997; and USDOI 1998.

<sup>e</sup> = Data not available.



## Mercury and Methylmercury in Water and Fish

### *Mercury and Methylmercury Dissolved in Water*

The average concentrations of mercury dissolved in ambient lake water samples are reported in Tables 4 and 5. The highest dissolved mercury concentrations were found at Red Lake (~3.63 ng/L) and was below our detection limit at Morgan Lake (<0.5 ng/L). Dissolved mercury concentrations were significantly correlated with water temperature ( $r^2 = -0.77$ ), specific conductivity ( $r^2 = -0.91$ ), secchi disk visibility ( $r^2 = -0.73$ ), turbidity ( $r^2 = 0.82$ ), maximum lake depth ( $r^2 = -0.88$ ) and surface-to-depth ratio ( $r^2 = -0.81$ ). That is, in the four lakes studied, more dissolved mercury was found in those lakes that were warmer, more turbid, and shallow (*e.g.*, Red Lake) and less in those lakes that were cold, clear and deep (*e.g.* Asaayi Lake).

The average concentrations of total methylmercury in ambient lake water samples are reported in Table 5. As a percentage of dissolved mercury, methylmercury was approximately 4% in Asaayi Lake, 8% in Wheatfields Lake, and 6 % in Red Lake. The highest average methylmercury concentrations were found at Red Lake (0.21 ng/L; higher at the south end and decreasing northward) and the lowest were found at Morgan Lake (0.02 ng/L). Average methylmercury concentrations were 0.07 ng/L at Asaayi Lake and 0.15 ng/L at Wheatfields Lake. Harris and Snodgrass (1993) reported that ambient background methylmercury concentrations are approximately 0.05 ng/L. They cautioned however, that due to the long-range atmospheric transport of mercury and other atmospheric emissions, the current “background” mercury concentrations in water probably reflect an anthropogenic influence even in “pristine” lakes. Total methylmercury concentrations in water were significantly correlated with maximum lake depth ( $r^2 = -0.60$ ), specific conductivity ( $r^2 = -0.53$ ), secchi disk visibility ( $r^2 = -0.68$ ), turbidity ( $r^2 = 0.70$ ), maximum lake depth ( $r^2 = -0.88$ ) and surface-to-depth ratio ( $r^2 = -0.54$ ). That is, similar to dissolved mercury, more methylmercury was found in those lakes that were warmer, more turbid, and shallow (*e.g.*, Red Lake) and less in those lakes that were cold, clear and deep (*e.g.* Asaayi Lake) though these correlations were not as robust.

Mercury concentrations in water were below the Navajo Nation numeric criteria for mercury for all designated uses. However, methylmercury is generally more toxic than inorganic mercury to aquatic organisms. Methylmercury in these lakes would not be expected to exert aquatic plant toxicity (*i.e.*, > 800 ng/L; USEPA 1997). Concentrations of methylmercury that induce toxic effects in aquatic invertebrates are greater than 40 ng/L. Wiener and Spry (1996) suggested that histological changes and effects to fish behavior, reproduction, and development could occur at water concentrations as low as 100 ng/L. During the period sampled, the range of methylmercury concentrations (0.017 to 0.418 ng/L) measured in the lake water did not exceed these levels of concern for toxicity to aquatic life.

### *Bald Eagle Water Quality Criteria*

Calculation of protective numeric criteria to protect wildlife (WC) such as the bald eagle through the consumption of fish was based upon a reference dose approach, combined with the extent to which mercury becomes concentrated in the fish from the four lakes sampled (per Russell 2003). The BAFs were  $4.2 \times 10^{-6}$  for Asaayi,  $1.6 \times 10^{-6}$  for Wheatfields Lake,

$7.5 \times 10^{-6}$  for Red Lake, and  $1.3 \times 10^{-6}$  for Morgan Lake. Using these BAFs, the methylmercury wildlife criteria (WC) would be 0.11 ng/L for Asaayi Lake, 0.27 ng/L for Wheatfields Lake, 0.06 ng/L for Red Lake, and 0.35 ng/L for Morgan Lake. The WC for mercury was calculated using the estimate of methylmercury as a proportion of dissolved mercury in water for Asaayi, Wheatfields, and Red Lakes (as 0.039, 0.084, and 0.059, respectively; mercury was not found above the detection limit at Morgan Lake). Using these values, a methylmercury WC of 0.11 ng/L, 0.27 ng/L, and 0.35 ng/L corresponds to a mercury WC of 2.7 ng/L at Asaayi Lake, 3.2 ng/L at Wheatfields Lake, and 1.0 ng/L at Red Lake. Therefore, the current water quality standard to protect aquatic life (12 ng/L) may not be protective of bald eagles through the consumption of fish from these waters. The WC for mercury is the concentration in surface water that, if not exceeded, protects both bald eagles that use the water for drinking or as a foraging source. Thus, the WC is the highest aqueous concentration of mercury that causes no significant reduction in growth, reproduction, or viability of a population of animals exposed over multiple generations and may be an appropriate goal to protect bald eagles that forage on fish from these lakes (*i.e.*, 1.0 ng/L).

#### *Mercury and Methylmercury in Fish Tissues*

The average concentrations of methylmercury and mercury in fish fillets or in re-integrated fish samples are reported in Tables 4 and 6. Mercury concentrations in trout ranged from 0.03-0.11 mg/kg WW and from 0.01-0.27 mg/kg WW in catfish. Stafford and Haines (1997) and Walter *et al.* (1973) reported ambient concentrations of mercury in trout (0.1-0.4 mg/kg WW) and channel catfish (0.1-0.3 mg/kg WW). Simpson and Lusk (2000) reported that concentration of mercury in whole body fish collected from the San Juan River were <0.05 to 0.32 mg/kg WW. The USDOI (1998) reported that warmwater fish (bluegill) experienced toxic effects above 0.11 mg/kg WW. All catfish samples from Red Lake had mercury concentrations above this threshold of concern, depending on species sensitivity.

By excluding Morgan Lake data (which were below the detection limit), dissolved mercury concentrations in water were significantly correlated with mercury concentrations in fish fillets ( $r^2 = 0.96$ ) and in re-integrated fish ( $r^2 = 0.96$ ). By including Morgan lake data, methylmercury concentrations in water were more weakly correlated with methylmercury concentrations in fish fillets ( $r^2 = 0.56$ ) and in re-integrated fish ( $r^2 = 0.39$ ). By excluding Morgan Lake data, methylmercury concentrations in fish fillets were significantly correlated ( $r^2 = 0.96$ ) with mercury concentrations in re-integrated fish.

Methylmercury concentrations in catfish fillets were significantly correlated with average length ( $r^2 = 0.87$ ) and average weight ( $r^2 = 0.88$ ). There was no significant correlation found for the trout size and methylmercury in their fillets. For consideration of a consumption advisory that is based on the catfish, the relationship between methylmercury concentrations in catfish fillets and average catfish weight (in grams) can be described by the equation:

$$\text{MeHg in Fillets} = -0.4512 + 0.0095 \times \text{Average Total Weight (g)} \quad (\text{Equation 9})$$

Similar to water, mercury concentrations in re-integrated fish were significantly correlated with the specific conductivity ( $r^2 = -0.75$ ), secchi disc visibility ( $r^2 = -0.67$ ), and turbidity ( $r^2 = 0.78$ ) of the lake surface, as well as the maximum depth ( $r^2 = 0.71$ ) and surface-

to-depth ratio ( $r^2 = -0.90$ ) of the lakes. That is, more mercury was found in fish from lakes that had lower oxygen content, less salinity, and were more turbid and shallow in relation to their size. Mercury concentrations in whole fish also positively correlated with concentrations of Ba ( $r^2 = 0.81$ ), Mn ( $r^2 = 0.76$ ), S ( $r^2 = 0.70$ ), and V ( $r^2 = 0.72$ ) in the tissue.

### Human Health Risks

The goal of the Navajo Nation Lake Fish and Water Quality Investigation was to provide data that may be used to *estimate* mercury risks to human health (and to bald eagles) from the consumption of fish obtained from selected recreational lakes on the Navajo Nation. These data can be also used to develop site-specific bioaccumulation factors and to evaluate the need for management actions to limit fish consumption, reduce the process of methylation, or perhaps recommend reductions of local mercury emissions and discharges under various authorities of the Navajo Nation. This study was not designed to determine the sources of mercury found in water or fish. This study also did not determine the fish consumption patterns of the local community, which is the most critical factor in estimating human health exposures and therefore for the estimation of potential health risks.

Without these local estimates of fish consumption, national default consumption values were assumed for this study. However, although large-scale surveys of mercury contents of fish and fish consumption patterns have been conducted (USEPA 2000), these surveys have limitations for applications to a human health risk assessment on the Navajo Nation. Estimates of dietary intakes of mercury in food based on national or regional data cannot accurately reflect the intake of mercury from locally harvested foods, including vegetables, fish, and game. This can be a particularly important limitation when the people of concern are those whose diet contains a large portion of locally caught fish. National estimates do not address how many anglers are in an area, when and how often they catch fish, and to whom they may distribute fish to within a community, or any cultural practices that augment or limit fish consumption. To reduce these uncertainties, the contributions of mercury in local fish can be assessed by conducting consumption surveys for the local population of concern.

Every local population in the United States is exposed to a wide variety of metals in air, food, drinking water, and soils derived from both natural and anthropogenic sources. The distinction between natural and anthropogenic sources can sometimes be further blurred when a previously deposited metal is recycled by natural processes, thereby becoming an important source in some ecosystems. For many metals, present-day exposures are expected to be above the natural background doses and the biological implications of any increase in levels of historical human exposures are unknown. One collateral effect of background contribution to human metal intake would be to push the total exposure over the toxicity threshold depending on a person's relative amount of fish consumption. This is likely the case for mercury.

Using default assumptions for fish consumption, body weight, days of exposure, and the geometric mean of methylmercury concentrations in fish fillets from the four lakes sampled, those men, women, and children that consume fish on a *recreational* basis are not at significant health risks (Table 9). However, children that eat fish on a frequent basis (>150

days per year), or any adult whose diet contains a large portion of catfish from Red Lake may be at risk of mercury toxicity. Using consumption scenarios and Red Lake fish fillet data, consumption of more than 2 fish per month would not be recommended (Table 10). In the context of consumption of contaminated fish, risk managers may need to consider the fish consumption patterns around Red Lake and, if necessary, seek ways to minimize the health risk to people that eat lots of fish, depending on local conditions including social and cultural factors and the benefits of eating fish. However, a long-term goal may be to reduce the mercury contamination of the water body or reduce the rate of mercury accumulation in the fish tissue through a variety of means, including source control and management.

### Bald Eagle Health Risks

Using default assumptions for fish consumption, body weight, days of exposure, and averaging time, and using the geometric mean of methylmercury concentrations in re-integrated fish from the four lakes sampled, bald eagles that migrate through the Navajo Nation and consume fish equally from the all four lakes sampled are not at significant health risks (Table 10). However, bald eagles that consume catfish from Red Lake on a frequent basis (>30 days per year), have the potential to experience mercury toxicity. Bald eagles that may feed on catfish from Red Lake for over 60 days are at a significant health risk to mercury. Bald eagles that forage exclusively from Morgan Lake have the lowest exposure to mercury but the highest exposure to selenium. Bald eagles that attempt to establish nesting on the Navajo Nation may need to be monitored for their long-term mercury exposure and effects. There are few management options to affect bald eagle consumption patterns around Red Lake, and therefore additional information would be needed to identify if extensive use is occurring and to determine if there are additional sources of mercury to Red Lake. However, a long-term goal to reduce the mercury contamination of the water body or reduce the rate of mercury accumulation in the fish tissue through a variety of means would be recommended for bald eagles that feed at Red Lake.

Table 8. Human Health Risk Quotients for Children, Women, and Men Using Various Fish Exposure Scenarios for Each Lake and for All Lakes Combined. [Note: exposure duration of 365 days and body mass data not shown]

<i>Lake Scenario</i>	<i>Consumer Scenario</i>	<i>MeHg in Fillets (mg/kg WW)</i>	<i>Fish Consumption (kg/day)</i>	<i>Exposure to Fish (days/yr)</i>	<i>MeHg Dose (mg/kg-day)</i>	<i>USEPA Reference Dose (RfD)</i>	<i>Risk Quotient &gt; 1 ?</i>
Asaayi Lake	<i>Child - Recreation</i>	0.06	0.0175	14	2.78E-06	0.0001	0.03
<b>Asaayi Lake</b>	<b>Child - Subsistence</b>	0.06	0.1424	156	2.52E-04	0.0001	<b>2.52</b>
Wheatfields Lake	<i>Child - Recreation</i>	0.08	0.0175	14	3.70E-06	0.0001	0.04
<b>Wheatfields Lake</b>	<b>Child - Subsistence</b>	0.08	0.1424	156	3.36E-04	0.0001	<b>3.36</b>
Morgan Lake	<i>Child - Recreation</i>	0.01	0.0175	14	4.63E-07	0.0001	0.00
Morgan Lake	<i>Child - Subsistence</i>	0.01	0.1424	156	4.20E-05	0.0001	0.42
Red Lake	<i>Child - Recreation</i>	0.39	0.0175	14	1.81E-05	0.0001	0.18
<b>Red Lake</b>	<b>Child - Subsistence</b>	0.39	0.1424	156	1.64E-03	0.0001	<b>16.37</b>
All lakes combined	<i>Child - Recreation</i>	0.07	0.0175	14	3.01E-06	0.0001	0.03
<b>All lakes combined</b>	<b>Child - Subsistence</b>	0.07	0.1424	156	2.73E-04	0.0001	<b>2.73</b>
Asaayi Lake	<i>Woman - Recreation</i>	0.06	0.0175	14	6.20E-07	0.0001	0.01
Asaayi Lake	<i>Woman - Subsistence</i>	0.06	0.1424	156	5.62E-05	0.0001	0.56
Wheatfields Lake	<i>Woman - Recreation</i>	0.08	0.0175	14	8.26E-07	0.0001	0.01
Wheatfields Lake	<i>Woman - Subsistence</i>	0.08	0.1424	156	7.49E-05	0.0001	0.75
Morgan Lake	<i>Woman - Recreation</i>	0.01	0.0175	14	1.03E-07	0.0001	0.00
Morgan Lake	<i>Woman - Subsistence</i>	0.01	0.1424	156	9.36E-06	0.0001	0.09
Red Lake	<i>Woman - Recreation</i>	0.39	0.0175	14	4.03E-06	0.0001	0.04
<b>Red Lake</b>	<b>Woman - Subsistence</b>	0.39	0.1424	156	3.65E-04	0.0001	<b>3.65</b>
All lakes combined	<i>Woman - Recreation</i>	0.07	0.0175	14	6.71E-07	0.0001	0.01
All lakes combined	<i>Woman - Subsistence</i>	0.07	0.1424	156	6.09E-05	0.0001	0.61
Red Lake	<i>Man - Recreation</i>	0.39	0.0175	14	3.36E-06	0.0001	0.03
<b>Red Lake</b>	<b>Man - Subsistence</b>	0.39	0.1424	156	3.04E-04	0.0001	<b>3.04</b>
All lakes combined	<i>Man - Recreation</i>	0.07	0.0175	14	5.59E-07	0.0001	0.01
All lakes combined	<i>Man - Subsistence</i>	0.07	0.1424	156	5.07E-05	0.0001	0.51

**Table 9. Estimation of the Maximum Allowable Fillet Consumption Rate and the Maximum Allowable Fish Consumption Rates for each Lake Scenario that has a Risk Quotient > 1. [Note: RfD of 0.0001 mg/kg-bw/day not shown]**

<b>Consumer Scenario</b>	<b>Lake Scenario</b>	<b>Consumer Body mass (kg)</b>	<b>Geometric Mean Fillet MeHg (mg/kg WW)</b>	<b>Maximum Allowable Daily Fillet Consumption (ounces/day)</b>	<b>Maximum Allowable Weekly Fillet Consumption (8-oz. fillets)</b>	<b>Maximum Allowable Number of Fish per Week (1-lb. fish)</b>	<b>Maximum Allowable Number of Fish per Month (1-lb. fish)</b>
Children - Subsistence	All lakes	14.5	0.07	0.8	6.3	3	13
Children - Subsistence	Asaayi Lake	14.5	0.06	0.9	6.8	3	14
Children - Subsistence	Red Lake	14.5	0.39	0.1	1.0	0.5	2
Children - Subsistence	Wheatfields Lake	14.5	0.08	0.6	5.1	3	10
Women - Subsistence	Red Lake	65.0	0.39	0.6	4.7	2	9
Men - Subsistence	Red Lake	78.0	0.39	0.7	5.6	3	11

Table 10. Bald Eagle Health Risk Quotients for Various Exposure Scenarios for Each Lake, for various Types of Lakes, and for All Lakes Combined. [Note: averaging time of 10950 days not shown]

Bald Eagle Exposure Scenario	Geometric Mean Mercury Concentration in Whole Fish (mg/kg WW)	Fraction of Fish in the Total Diet	Food Ingestion Rate (kg/day)	Exposure Frequency (days an eagle spends in an area)	Exposure Duration (estimated bald eagle lifetime - years)	Bald Eagle Body Mass (kg)	Mercury Ingestion Rate (mg/kg/day)	Risk Quotient > 1?
Migrant ~ All lakes combined	0.05	0.71	0.583	30	30	0.525	0.0073	0.3
Migrant ~ Asaayi Lake	0.07	0.71	0.583	30	30	0.525	0.0067	0.3
Migrant ~ Coldwater lakes	0.06	0.71	0.583	30	30	0.525	0.0078	0.4
Migrant ~ Morgan Lake	0.01	0.71	0.583	30	30	0.525	0.0011	0.1
<b>Migrant ~ Red Lake</b>	0.23	0.71	0.583	30	30	0.525	0.0436	<b>2.1</b>
Migrant ~ Warmwater lakes	0.05	0.71	0.583	30	30	0.525	0.0067	0.3
Migrant ~ Wheatfields Lake	0.06	0.71	0.583	30	30	0.525	0.0089	0.4
<b>Wintering ~ All lakes combined</b>	0.05	0.71	0.583	182	30	0.525	0.0441	<b>2.1</b>
<b>Wintering ~ Asaayi Lake</b>	0.07	0.71	0.583	182	30	0.525	0.0407	<b>1.9</b>
<b>Wintering ~ Coldwater lakes</b>	0.06	0.71	0.583	182	30	0.525	0.0474	<b>2.3</b>
Wintering ~ Morgan Lake	0.01	0.71	0.583	182	30	0.525	0.0068	0.3
<b>Wintering ~ Red Lake</b>	0.23	0.71	0.583	182	30	0.525	0.2643	<b>12.6</b>
<b>Wintering ~ Warmwater lakes</b>	0.05	0.71	0.583	182	30	0.525	0.0407	<b>1.9</b>
<b>Wintering ~ Wheatfields Lake</b>	0.06	0.71	0.583	182	30	0.525	0.0542	<b>2.6</b>
<b>Nesting ~ All lakes combined</b>	0.05	0.71	0.583	365	30	0.525	0.0883	<b>4.2</b>
<b>Nesting ~ Asaayi Lake</b>	0.07	0.71	0.583	365	30	0.525	0.0815	<b>3.9</b>
<b>Nesting ~ Coldwater lakes</b>	0.06	0.71	0.583	365	30	0.525	0.0951	<b>4.5</b>
Nesting ~ Morgan Lake	0.01	0.71	0.583	365	30	0.525	0.0136	0.6
<b>Nesting ~ Red Lake</b>	0.23	0.71	0.583	365	30	0.525	0.5301	<b>25.2</b>
<b>Nesting ~ Warmwater lakes</b>	0.05	0.71	0.583	365	30	0.525	0.0815	<b>3.9</b>
<b>Nesting ~ Wheatfields Lake</b>	0.06	0.71	0.583	365	30	0.525	0.1087	<b>5.2</b>