



Figure 2. Alluvial ferricrete along Mineral Creek between Middle and South Forks of Mineral Creek (view to west). Outcrop is about 30 m thick and rests on granulated porphyry bedrock. Ferricrete forms in paleo-alluvial terrace deposits where terrace sands and gravels are cemented by iron oxyhydroxide minerals.




Figure 3. Alluvial ferricrete above mouth of Cement Creek (view to south). Kendall Mountain in distance.



Figure 4. Log in alluvial ferricrete along Middle Fork Mineral Creek is below and about 400 m downstream from Bonner mine. Note the cemented in place by iron oxyhydroxide cement. The radiocarbon age of this log is 700 ± 80 yr B.P. Logs encased in ferricrete throughout the upper Animas River watershed range in age from modern to 5,150 yr B.P.





Figure 5. Alluvial ferricrete at mouth of Middle Fork Kendall Creek.



Base from U.S. Geological Survey 1995, 3.3 ft (1 m) resolution digital orthoquadrants. Quadrangles and include Ironsides, Haines Peak, Ophir, Silverton, and Howardsville.




Figure 6. An active iron spring located about 200 m above the confluence of Mineral Creek and Middle Fork Mineral Creek (view to south). Wet outcrop is about 15 cm deep, and consists of both living and decaying sedges, grasses, mosses, and iron precipitates accumulate. Note acidophilic, dark-green sphagnum moss at base of spring.





Figure 7. Iron spring in lower Prospect Gulch near head of Cement Creek (view to south). Bog iron and colloidal ferricrete deposits are preserved adjacent to spring.



Figure 8. Alluvial ferricrete at base of excavated foundation (town of Silverton). Builders commonly encounter ferricrete "hardpan" when excavating foundations throughout the town.



CORRELATION OF MAP UNITS

ab	acp	ub	cf	cf	cf	cf	cf	cf	cf
ab	acp	ub	cf	cf	cf	cf	cf	cf	cf

DESCRIPTION OF MAP UNITS

ab Sedge bog (late Pleistocene to modern) air-saturated ground that is colonized by acid-adjacent sedges, grasses, mosses, and willows with pH typically ranging from 1.2 to 5.5. Sedge bogs generally form at the base of hillslopes, at sites where the water table is intersected by the ground surface, and develop from colloidal ferricrete deposited above, however, manganese is a sufficient abundance to cause encrust matrix and clay coating to have a brown to black color. The term transitional is used to describe those deposits that are observed to have a mixed manganese and ferricrete composition. Deposits are preserved in Placer Gulch. Thickness 0.5 to 3 m.

acp Iron spring (late Pleistocene to modern) air-saturated, predominantly brown to reddish-brown, but may be whitish-gray to yellow or orange, impure hydrous iron oxide deposits in water-saturated ground. Consist of nodules of iron, aluminum, and manganese compound precipitates. Precipitates form in acidic, poorly drained conditions by the oxidizing action of algae, iron (thio)halophilic ferromicrobials and sulfur (thio)halophilic (thio)acidobacteria existing beneath, or the atmosphere. Actinophyllite sponges and mosses may be present. Substrate consists of hydrous iron oxyhydroxide (schwertmannite), amorphous iron oxyhydroxide, and goethite, which have porous textures ranging from thinly layered to irregular aggregations. Algal mats trap freshly formed oxyhydroxide-oxyhydroxide precipitates. Clans, logs, and twigs are locally preserved. Observed thickness ranges from 0.1 to 0.5 m.

ub Undifferentiated bog (late Pleistocene to modern) Sedge bog or iron bog, as described above. Unit is undifferentiated because deposits were either identified from aerial photograph interpretation or were not accessible due to land ownership issues.

cf Alluvial ferricrete (late Pleistocene to modern) brown to yellowish-brown, iron oxyhydroxide-cemented sandstone or conglomerate; cement consists principally of goethite. Deposits are bedded to weakly stratified and consist mostly of heterogeneous subrounded to subangular pebbles and cobbles with occasional boiler-size clasts in an iron oxyhydroxide-cemented, clay-sand matrix. Pebbles and cobbles in some places are coated with a fine filamentous iron oxyhydroxide cement similar in appearance to algae. Exceptional (as much as 30%) alluvial ferricrete exposures preserved along the west side of Mineral Creek between South Fork and Middle Fork of Mineral Creeks, and near the mouth of Cement Creek. Silts, clays, and shales are common in some coarse-grained graded beds of gravel, which are indicators of high-energy stream transport. Alluvial ferricrete deposits are either wet or dry at present. Along active flood-plain channels, such as Cement and Mineral Creeks, seeps and springs flow from remaining ferricrete. Substrate consists of heterogeneous iron-rich zones, from 0.5 to 2 m from stream beds. Conifer logs are locally fossil within these deposits and the alluvium locally interfingers with peat. Alluvial ferricrete preserved several meters above the active channel is often dry and represents cemented alluvial fan remnants and stream terrace deposits. ¹⁴C ages from logs and twigs recovered from these deposits range in age from modern to 5,600 yr B.P. Thickness, 0.5 to 3 m.

cf Colloidal ferricrete (late Pleistocene to modern) iron oxyhydroxide cemented deposits, varicolored, brown (predominant), reddish-brown to brownish-yellow with dark brown stained clasts. Cement consists primarily of goethite. Deposits are massive to weakly stratified subparallel to the current slope or stage existing topography; consist of mostly homogeneous angular, subangular, or subrounded pebbles, cobbles, and boulders in an iron oxyhydroxide-cemented, fine-grained matrix, with variable clay matrix. Cobbles are weakly subrounded and dip downwind. Clans consist of subangular to subrounded pebbles in contact with silty and sand-size sediment. Pebbles and cobbles are locally coated with a fine filamentous iron oxyhydroxide cement similar in appearance to algae. Fine logs and twigs or other organic materials are preserved. Colloidal ferricrete deposits are either wet or dry, and are formed on hillslopes and in narrow debris channels where rock and soil clasts maintain its colluvial, alluvial, and alluvial fan deposits. Source materials were derived from weathering of local bedrock that was transported less than a few kilometers. ¹⁴C ages on wood fragments from colloidal ferricrete deposits have yielded radiocarbon ages ranging from 870 ± 80 yr B.P. whereas ¹⁴C ages of fully rippled casts of woody material and undifferentiated "organic carbon" have minimum ¹⁴C ages ranging from 1,170 to 1,600 yr B.P. Maximum thickness 2 to 7 m, maximum thickness unknown.

cf Undifferentiated ferricrete (late Pleistocene to modern) Colloidal or alluvial ferricrete, as described above. Colloidal ferricrete likely is preserved on hillslopes several meters above tributary streams, alluvial ferricrete forms in alluvial terraces and fans.

cf Colloidal manganese (late Pleistocene to modern) black to dark-gray, manganese-rich ferricrete. Scanning electron microscopy analyses indicate that matrix consists of Mn and Fe-rich material. Unit description is similar to colloidal ferricrete (cf), however, manganese has sufficient concentrations, about 2.4 to 4.8 weight percent, to impart a black to dark gray color. Some manganese outcrops are transitional toward more highly iron enriched ferricrete. The distribution of manganese is confined to areas that the manganese-rich ferricrete was deposited in the Placer Gulch, such as Placer Gulch, California Gulch, and the Eureka basin where the original, late 1800's Sumaside mine adit was excavated. Thickness 0.5 to 3 m.

cf Alluvial manganese (late Pleistocene to modern) black to dark-gray, manganese-rich ferricrete. Scanning electron microscopy analyses indicate that matrix consists of Mn and Fe-rich material. Unit description is similar to alluvial ferricrete (cf), however, manganese has sufficient concentrations, about 2.4 to 4.8 weight percent, to impart a black to dark gray color. Some manganese outcrops are transitional toward more highly iron enriched ferricrete. One manganese outcrop sampled in Eureka Gulch adjacent to the original Sumaside mine adit near the former Lake Emma consists of finely layered, silty to sand-size material at its base, which grades upward into a sequence of clay matrix supporting subangular to subrounded pebbles and cobbles and alternating interbedded coarse sands. This deposit is thought to represent an alluvial fan that was deposited into the former Lake Emma. A matrix radiocarbon date of this deposit yielded an age of 8,000 ± 80 yr B.P. The distribution of manganese is principally confined to areas that the manganese-rich ferricrete was deposited in the Placer Gulch, such as Placer Gulch, California Gulch, and the Eureka basin, where the original, late 1800's Sumaside mine adit was excavated. Alluvial manganese deposits in the former Eureka watershed, however, have been identified as far south as beyond Elbow Park, south of Silverton. Thickness 0.5 to 3 m.

cf Alluvial manganese (late Pleistocene to modern) black to dark-gray, manganese-rich ferricrete. Scanning electron microscopy analyses indicate that matrix consists of Mn and Fe-rich material. Unit description is similar to alluvial ferricrete (cf), however, manganese has sufficient concentrations, about 2.4 to 4.8 weight percent, to impart a black to dark gray color. Some manganese outcrops are transitional toward more highly iron enriched ferricrete. One manganese outcrop sampled in Eureka Gulch adjacent to the original Sumaside mine adit near the former Lake Emma consists of finely layered, silty to sand-size material at its base, which grades upward into a sequence of clay matrix supporting subangular to subrounded pebbles and cobbles and alternating interbedded coarse sands. This deposit is thought to represent an alluvial fan that was deposited into the former Lake Emma. A matrix radiocarbon date of this deposit yielded an age of 8,000 ± 80 yr B.P. The distribution of manganese is principally confined to areas that the manganese-rich ferricrete was deposited in the Placer Gulch, such as Placer Gulch, California Gulch, and the Eureka basin, where the original, late 1800's Sumaside mine adit was excavated. Alluvial manganese deposits in the former Eureka watershed, however, have been identified as far south as beyond Elbow Park, south of Silverton. Thickness 0.5 to 3 m.

cf Alluvial manganese (late Pleistocene to modern) black to dark-gray, manganese-rich ferricrete. Scanning electron microscopy analyses indicate that matrix consists of Mn and Fe-rich material. Unit description is similar to alluvial ferricrete (cf), however, manganese has sufficient concentrations, about 2.4 to 4.8 weight percent, to impart a black to dark gray color. Some manganese outcrops are transitional toward more highly iron enriched ferricrete. The distribution of manganese is confined to areas that the manganese-rich ferricrete was deposited in the Placer Gulch, such as Placer Gulch, California Gulch, and the Eureka basin where the original, late 1800's Sumaside mine adit was excavated. Thickness 0.5 to 3 m.

cf Alluvial manganese (late Pleistocene to modern) black to dark-gray, manganese-rich ferricrete. Scanning electron microscopy analyses indicate that matrix consists of Mn and Fe-rich material. Unit description is similar to alluvial ferricrete (cf), however, manganese has sufficient concentrations, about 2.4 to 4.8 weight percent, to impart a black to dark gray color. Some manganese outcrops are transitional toward more highly iron enriched ferricrete. One manganese outcrop sampled in Eureka Gulch adjacent to the original Sumaside mine adit near the former Lake Emma consists of finely layered, silty to sand-size material at its base, which grades upward into a sequence of clay matrix supporting subangular to subrounded pebbles and cobbles and alternating interbedded coarse sands. This deposit is thought to represent an alluvial fan that was deposited into the former Lake Emma. A matrix radiocarbon date of this deposit yielded an age of 8,000 ± 80 yr B.P. The distribution of manganese is principally confined to areas that the manganese-rich ferricrete was deposited in the Placer Gulch, such as Placer Gulch, California Gulch, and the Eureka basin, where the original, late 1800's Sumaside mine adit was excavated. Alluvial manganese deposits in the former Eureka watershed, however, have been identified as far south as beyond Elbow Park, south of Silverton. Thickness 0.5 to 3 m.

Figure 12. Colloidal ferricrete above South Fork Cement Creek, deposited on a hillslope topographically below altered lava flows (upper right). View to east.

Figure 13. Bog iron outcrop near headwater region of Eureka Gulch. Deposit is essentially clay free and consists of fine, horizontally laminated iron oxyhydroxide (goethite). Vertically laminated bog iron deposits also crop out in upper Animas River watershed (Fig. 11). Bog iron deposits are inactive, and are likely the remnants of once active iron springs and iron bogs.

Figure 14. Iron bog located east of Dry Gulch along Cement Creek. Iron bogs are shallow mixing zones, having at this stage of pH (3.2–5.7) and conductivity (900–1,700 microsiemens per centimeter) where Oxidation of ferrous iron and mixing of ground water result in precipitation of iron trapped in pools of organic material. Iron bogs are often transitional to sedge bogs.

Figure 15. Sedge bog in South Fork Mineral Creek subbasin. Sedges, grasses, and willows are prevalent along bog margins. Dark area of photograph is water saturated, about 15 cm deep, and consists of both living and decaying sedges, grasses, mosses, and shrubs (photograph by M.R. Stanton, U.S. Geological Survey).




Figure 16. Inset map of Animas River watershed.

PROJECT DESCRIPTION AND STUDY AREA

During 1996–2006, the Bureau of Land Management, National Park Service, Environmental Protection Agency, United States Department of Agriculture (USDA) Forest Service, and the U.S. Geological Survey (USGS) developed a coordinated strategy to (1) study the environmental effects of historical mining on Federal lands, and (2) monitor contaminated sites that have the greatest impact on water quality and ecosystem health. The focus of our involvement in this study was to develop a methodology to identify and characterize watersheds that are most at risk for degradation caused by historical mining. A watershed scale of observation was utilized because of the riparian ecosystem of the upper Animas River watershed was chosen for study in large part because of the hundreds of inactive mines and prospects scattered throughout the watershed.

One important objective of our USGS Abandoned Mine Lands Initiative was to establish the preliminary geochemical baseline conditions of the Animas River watershed to aid in understanding of this characteristic, it is needed to establish achievable restoration goals. Ferricrete (trifoliate iron- and manganese-oxyhydroxide-cemented secondary deposits) are one indicator of the geochemical baseline conditions as well as the effect that weathering of mineralized rocks had on water quality in the Animas River watershed prior to mining. The term ferricrete was first used by Lamplugh (1902) to describe iron-cemented surficial sands and gravels formed by precipitation of infiltrating solutions of "iron salts." Ferricretes occur in several mining districts throughout the western United States. They have been used as an exploratory tool to map and assess iron-rich ferricrete. One high trace element abundance (Pb, Verplank, D.B. Yager, and S.E. Church, work in progress) and as an indicator of paleo-weathering conditions (Harris-Quinn, 1969; Harris and others, 1999). Lead and wood fragments preserved in several ferricrete at the upper Animas River watershed, collected only along the river, yield radiocarbon ages of modern 5,500 years B.P. (P.L. Verplank, D.B. Yager, and S.E. Church, work in progress). Radiocarbon ages of rings of wood collected from ferricrete in the study area overlap the range of post-deglaciation ages determined for wood fragments collected from iron deposits at the base of Eureka Gulch, northeast of Silverton, Colo. (Carrara and others, 1991; Carrara and others, 1991; Elias and others, 1991). The presence of ferricrete deposits along the current stream course indicates that climate and physiography of the Animas River watershed have been relatively constant throughout the Holocene and that weathering processes have been ongoing for thousands of years prior to mining activities. Thus, by knowing where ferricrete is preserved in the watershed today, land-management agencies have an indication of (1) where metal precipitation from weathering of altered rocks has occurred in the past and (2) where this process is ongoing and may confound remediation efforts.

We mapped the distribution of ferricrete and determined their physical properties as part of the Animas River watershed study to build a spatial framework for observing the processes responsible for their formation and preservation, and to document the stability of the current weathering surface throughout the Holocene. The Animas River watershed study area, as defined for this study, is the drainage of three tributaries (Mineral and Cement Creeks, and the Animas River upstream from their confluence near Silverton, Colo.) is ideally suited for study of detailed ferricrete occurrences because (1) the combined weathering, glacial, hydrologic, erosion, and deposition processes have preserved and exposed numerous ferricrete outcrops, (2) iron- and manganese-rich springs and seeps are abundant, (3) bedrock and surficial deposit exposures are excellent, and (4) ferricrete deposits occur in both inactive mines and naturally occurring alteration zones found throughout the watershed.

MAP SUMMARY

This map shows the distribution of ferricrete, manganese, and iron bog and iron spring in the upper Animas River watershed. The Mineral and Cement Creeks basins were mapped in detail to delineate extent and variation of ferricrete occurrences. However, the Animas River basin upstream of the town of Silverton was mapped at a reconnaissance level of detail. Field data were compiled on the Haines Peak, Ophir, Silverton, and Howardsville 1:24,000 topographic maps and on 1:63,000 scale aerial photos of Cement Creek. All data were digitized with ARCTIC (ARCTIC ERAS/BAKING) software. Two coverages were created, including an active bog coverage and a ferricrete coverage. Digital orthoquadrants (DOQ), 1997, Unavailable for release under the provisions of the National Information Security Act (50 USC 3605), were used to create a background map. The background map was created with a 1 m resolution where data were compiled and digitized from the topographic maps and aerial photos (DOQ) have a resolution of 30 m. Data were compiled and digitized from the ferricrete coverage in stream terraces between Mineral and Cement Creeks. MAPublisher was used to import the ArcView coverage and digital orthoquadrants into Adobe Illustrator to produce the final map.

FIELD METHODS

Physical properties were recorded at each outcrop to create a classification scheme (P.L. Verplank, D.B. Yager, and S.E. Church, work in progress). Important observations include clast presence or absence, nature and degree of induration, grain matrix, matrix type, porosity, induration, and occurrence of small-scale structures, orientation of layers, and dimensions of outcrop. We also noted whether the outcrop was wet or dry to determine if the deposit was active or ancient, observing that wetness or temporal variation in ground-water flow is not always a definitive indicator of active or ancient. Five principal classes of these non-cemented deposits were mapped: 1) bog iron, thinly bedded deposits with essentially bedded deposits with associated with active or paleo-springs; 2) colloidal ferricrete, massive to finely bedded deposits with angular clasts that are primarily monolithologic; 3) alluvial ferricrete, massive to finely bedded deposits with laminated and commonly indurated clasts; 4) alluvial and colloidal manganese, deposits within the alluvial and colloidal class types that are very dark brown to black in outcrop owing to the presence of highly elevated concentrations of manganese and iron matrix cement; 5) transitional ferricrete and manganese, compositionally transitional between manganese and ferricrete.

FERRICRETE, MANGANOCRETE, AND BOG IRON OCCURRENCES WITH SELECTED SEDGE BOGS AND ACTIVE IRON BOGS AND SPRINGS IN PART OF THE ANIMAS RIVER WATERSHED, SAN JUAN COUNTY, COLORADO

By Douglas B. Yager, Stanley E. Church, Philip L. Verplank, and Laurie Wirt

Manuscript approved for publication on February 19, 2005. Any use of trade names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Geological Survey. This map was produced on request, directly from digital files, electronic photos. For sale by the U.S. Geological Survey Information Services, 486 1/2 Federal Center, Denver, CO 80225. 1-888-ASK-USGS. ARCIINFO coverages and a PDF map are available at http://pubs.usgs.gov/