

USAID Country Health Statistical Report Bulgaria

1. About the Report

2. Country Profile

I. Statistical Overview

- Demographic Indicators
- Socioeconomic Indicators
- Family Planning Indicators
- Maternal Health Indicators
- Child Survival Indicators
- HIV/AIDS Prevention Indicators
- Infectious Diseases Control Indicators

II. Trends in Health Indicators

- Current and Projected Population by Age and Sex (Figure 1 and Table 1.1)
- Population Estimates and Projections (Table 1.2)
- Estimated Total Fertility Rate per Woman (Figure 2 and Table 2)
- Infant Mortality Rates / Under-5 Mortality Rates (Figure 3 and Table 3)
- Vaccination Coverage (DPT3) Trends (Figure 4 and Table 4)
- Contraceptive Prevalence Rates Married Women (Figure 5 and Table 5)

3. Data Notes

4. Sources

About the Report

Released May 2008

This report is one of a series of Country Health Statistical Reports produced on behalf of the United States Agency for International Development (USAID) by the Analysis, Information Management and Communication Activity (AIM) Project. Each profile contains statistical data on current health conditions, population dynamics, health and family planning behavior, and health and population trends in a given developing country. Information is compiled from AIM's health statistics database, which draws data from a diverse range of sources listed at the end of this profile.

Hard copy editions of USAID's profiles are available from AIM'S publications department. Reports are available on the USAID Global Health Web site in Portable Document Format (pdf). Any feedback you have on the content or presentation of this report would be greatly appreciated. We would also appreciate receiving any more recent, more accurate, or more representative information. Contact us at the following:

> Analysis, Information Management & Communications Activity (AIM) 655 15th Street, NW, Suite 450 Washington, DC 20005

Phone: 202-661-8600 Fax: 202-661-8601 E-mail: dolphn@aimglobalhealth.org

The Analysis, Information Management & Communications (AIM) Activity, a USAID contract managed by MasiMax Resources, Inc, with John Snow, Inc., ORC Macro, and Insight Systems Corporation prepared this document under task order number GPO-M-00-05-00043-00, with the Bureau for Global Health, USAID.

I. Statistical Overview

*see data notes

Demographic Indicators				
Indicator	Value	Data Unit	Year	Source
Total Population	7,262,675		2008	BUCEN-IDB-2008
Population Growth Rate	<0.01	%	2008	BUCEN-IDB-2008
Percent Urban	70	%	2004	World Bank/WDI-2006
Women, 15-19	207,721		2008	BUCEN-IDB-2008
Women, 15-49	1,728,484		2008	BUCEN-IDB-2008
Life Expectancy at Birth	72.8		2008	BUCEN-IDB-2008
Crude Birth Rate	9.6	per 1,000	2008	BUCEN-IDB-2008
Crude Death Rate	14.3	per 1,000	2008	BUCEN-IDB-2008
Number of Live Births	69,576		2008	BUCEN-IDB-2008
Healthy Life Expectancy: Female	67.1		2002	WHO World Health Report-2004
Healthy Life Expectancy: Male	62.6		2002	WHO World Health Report-2004
Population: Percent < Age 15	13.8	%	2008	BUCEN-IDB-2008

Socioeconomic Indicators				
Indicator	Value	Data Unit	Year	Source
GNI per Capita (PPP)	8,630	\$	2005	World Bank/WDI-2006
Health Expenditure as Percentage of GDP	7.5	%	2003	World Bank/WDI-2006
Physicians per 1,000 People	3.6	per 1,000	1997-2004	World Bank/WDI-2006
Adult Literacy Rate	98.2	%	2004	World Bank/WDI Database-2007
Adult Literacy Rate, Female	98 1	%	2002	World Bank/WDI-2006
Adult Literacy Rate, Male	99 ²	%	2002	World Bank/WDI-2006
Gross Enrollment Ratio - Primary School	102.2	%	2005	UNESCO EFA Global Monitoring Report- 2008
Gender Parity Index - Net Enrollment Ratio	0.99		2005	UNESCO EFA Global Monitoring Report- 2008
Access to an Improved Water Source (Rural)	100	%	2002	World Bank/WDI-2005
Access to an Improved Water Source (Urban)	100	%	2002	World Bank/WDI-2005
Access to Improved Sanitation Facilities (Rural)	100	%	2002	World Bank/WDI-2005
Access to Improved Sanitation Facilities (Urban)	100	%	2002	World Bank/WDI-2005
Population Living Below \$1 a Day	2	%	1990-2004	Human Development Report-2006
Human Development Index	0.82		2005	Human Development Report - 2007/2008- 2008
Population Below Poverty Line	14.1 ³	%	2003	CIA World Factbook-2008
Real GDP (Growth) Rate	6.1 ⁴	%	2007	CIA World Factbook-2008
Income Inequality (Gini index)	31.6		2005	CIA World Factbook-2008
Unemployment Rate	8 5	%	2007	CIA World Factbook-2008
Inflation Rate	7.8 ⁶	%	2007	CIA World Factbook-2008
Airports	214		2007	CIA World Factbook-2008
Airports with paved runways	131		2007	CIA World Factbook-2008

¹ Based on census data

² Based on census data

³ Estimate

⁴ estimate

⁵ estimate

⁶ Estimate

Family Planning Indicators				
Indicator	Value	Data Unit	Year	Source
Total Fertility Rate (BUCEN)	1.4		2008	BUCEN-IDB-2008
Total Fertility Rate (DHS)		Data I	Not Ava	ilable
Contraceptive Prevalence Rate, Modern Methods, All Women		Data I	Not Ava	ilable
Contraceptive Prevalence Rate, Modern Methods, Married Women		Data I	Not Ava	ilable
Median Age of Sexual Debut Among Women, Ages 25-49		Data I	Not Ava	ilable
Mean Ideal Family Size		Data I	Not Ava	ilable
Women 20-24 Who Gave Birth Before Age 20		Data I	Not Ava	ilable
Total Fertility Rate, Rural		Data I	Not Ava	ilable
Unmet Need for Family Planning: Rural		Data I	Not Ava	ilable
Unmet Need for Family Planning: Urban		Data I	Not Ava	ilable
Contraceptive Prevalence Rate, All Methods		Data I	Not Ava	ilable
Equity in use of modern contraception		Data I	Not Ava	ilable
Percent of need satisfied by modern methods of family planning		Data I	Not Ava	ilable

Maternal Health Indicators					
Indicator	Value	Data Unit	Year	Source	
Maternal Mortality Ratio (WHO/Hill)	11 Per 100,000 live births 2005 WHO/Hi				
Maternal Mortality Ratio (DHS)		Data Not Ava	ilable		
Antenatal Care (at least 1 visit)		Data Not Ava	ilable		
Antenatal Care (2+ visits)	Data Not Available				
Antenatal Care (4+ visits)		Data Not Ava	ilable		
Assisted Delivery by a Health Professional		Data Not Ava	ilable		
Assisted Delivery by Doctor		Data Not Ava	ilable		
Assisted Delivery by Other Health Professional		Data Not Ava	ilable		
Anemia prevalence among women of reproductive age	Data Not Available				
Equity in skilled attendance at delivery		Data Not Ava	ilable		

Child Survival Indicators								
Indicator	Value	Data Unit	Year	Source				
Acute Respiratory Infection (ARI) and Oral Rehydrati	ion Thera	apy (ORT)						
ARI Care Seeking - Children Under 5	Data Not Available							
ORT Use Rate (ORS, RHS, or Increased Fluids)		Da	ta Not Av	vailable				
Mortality Indicators								
Infant Mortality Rate (BUCEN)	18.5	per 1,000 live births	2008	BUCEN-IDB-2008				
Infant Mortality Rate (UNICEF)	12		2006	UNICEF-2008				
Infant Mortality Rate (DHS)		Da	ta Not Av	vailable				
Infant Mortality Rate, Females (BUCEN)	14.8	per 1,000 live births	2008	BUCEN-IDB-2008				
Infant Mortality Rate, Males (BUCEN)	22	per 1,000 live births	2008	BUCEN-IDB-2008				
Under-5 Mortality Rate (BUCEN)	22.8	per 1,000 live births	2006	BUCEN-IDB-2006				
Neonatal Mortality Rate		Da	ta Not Av	vailable				
Under-5 Mortality Rate (UNICEF)	14	per 1,000 live births	2006	UNICEF-2008				
Under-5 Mortality Rate (DHS)	Data Not Available							
Nutrition Indicators								
Exclusive Breastfeeding (under 4 mos.)		Da	ta Not Av	vailable				
Exclusive Breastfeeding (under 6 mos.)		Da	ta Not Av	vailable				
Stunted (height-for-age)		Da	ta Not Av	vailable				
Underweight (weight-for-age)		Da	ta Not Av	vailable				
Wasted (weight-for-height)		Da	ta Not Av	vailable				
Vaccination Coverage								
DPT3 Vaccination Rate (DHS)		Da	ta Not Av	vailable				
DPT3 Vaccination Rate (WHO)	95	%	2006	WHO/Global Summary-2008				
Measles Vaccination Rate (DHS)		Da	ta Not Av	vailable				
Measles Vaccination Rate (WHO)	96	%	2006	WHO/Global Summary-2008				
Polio Vaccination Rate (DHS)		Da	ta Not Av	vailable				
Polio Vaccination Rate (WHO)	96	%	2006	WHO/Global Summary-2008				
Tetanus Toxoid Vaccination		Da	ta Not Av	vailable				
Equity in DPT3 coverage (DHS)		Da	ta Not Av	vailable				
Improved sanitation/hygiene practices		Da	ta Not Av	vailable				
Vitamin A supplementation coverage among children 6- 59 months		Da	ta Not Av	/ailable				

HIV/AIDS Prevention Indicators						
Indicator	Value	Data Unit	Year	Source		
HIV Prevalence proportion: Adults (15–49 years)	Data Not Available					
Estimated number of people living with HIV: Adults and Children	ated number of people living with HIV: Adults and Children Data Not Available					
Estimated number of people living with HIV: Women (15+ years)		Data Not Av	vailable			
Estimated number of people living with HIV: Children (0–14 years)		Data Not Av	vailable			
ales Reporting Condom Use With Last Non-Regular Partner Data Not Available						
Females Reporting Condom Use With Last Non-Regular Partner Data Not Available						

Infectious Diseases Control Indicators				
Indicator	Value	Data Unit	Year	Source
TB Estimated Number of Cases	3,101		2006	WHO Global Health Atlas-2008
TB Case Detection Rate	94	%	2006	WHO Global Health Atlas-2008
TB (DOTS) Treatment Success Rate	86	%	2005	WHO Global Health Atlas-2008
Malaria Cases per 100,000				Data Not Available
DOTS Coverage	100	100 % 2006 V		WHO Global Health Atlas-2008
HIV-Infected with Active TB	<0.01	%	2006	WHO Global Health Atlas-2008
Avian Influenza: Cumulative Number of Confirmed Human Cases			C	Data Not Available
Avian Influenza: Cumulative Number of Confirmed Human Deaths			C	Data Not Available
Child Sleeps Under Insecticide-Treated Net			C	Data Not Available
Equity in Bednet (ITN) Ownership			C	Data Not Available

II. Trends in Health Indicators




Figure 1: Current and Projected Population by Age and Sex (in thousands)

Table 1.1: Current and Projected Population by Age and Sex Source: BUCEN-IDB

Age Group	Male - 2000	Source: BUCE Male - 2020	Female - 2000	Female - 2020
0-4	171,096	140,884	161,911	133,459
5-9	201,063	155,484	191,391	147,281
10-14	256,509	166,879	245,183	158,225
15-19	279,596	169,467	263,212	160,522
20-24	306,319	153,052	289,403	143,825
25-29	283,162	180,396	276,453	171,725
30-34	256,542	229,207	256,523	221,382
35-39	251,710	243,347	258,277	235,113
40-44	262,773	263,762	274,171	260,745
45-49	271,159	240,565	287,115	250,134
50-54	276,955	214,311	299,012	231,464
55-59	220,568	204,719	245,359	231,276
60-64	201,673	202,914	235,184	240,651
65-69	203,842	190,502	248,615	240,987
70-74	165,381	167,004	217,275	229,966
75-79	115,444	102,906	169,837	158,884
80+	67,582	106,708	108,200	221,211

TABLE 1.2: Population Estimates and Projections

1960	1970	1980	1990	2000	2010	2020	2030	2040	2050					
7,867,374	8,489,574	8,843,528	8,894,028	7,818,495	7,148,785	6,568,957	5,940,822	5,305,010	4,651,477					




Figure 2: Estimated Total Fertility Rate per Woman

 Table 2: Estimated Total Fertility Rate per Woman

 Source: BUCEN-IDB

1960	1970	1980	2000	2010	2020	2030	2040	2050
2.3	2.2	2.1	1.3	1.4	1.5	1.6	1.6	1.7

Figure 3: Infant Mortality Rates / Under-5 Mortality Rates




Table 3: Infant Mortality Rates / Under-5 Mortality Rates

Source:	BUCEN-IDB 2008
---------	----------------

Indicator	1950	1960	1970	1980	2000	2010	2020	2030	2040	2050
Infant Mortality Rate (BUCEN)	94.5	45.1	27.3	20.2	24.6	17.3	12.4	9.3	7.2	5.9
Under-5 Mortality Rate (BUCEN)					28.4	19.8	14.2	10.5	8.2	6.6




Figure 4: Vaccination Coverage (DPT3) Trends



 Source: WHO/Global Summary

 1985
 1990
 1995
 2000
 2005
 2006

99.0 99.0 96.0 93.0 96.0 95.0

Figure 5: Contraceptive Prevalence Rates, Modern Methods, Married Women Data Not Available

 Table 5: Contraceptive Prevalence Rates, Modern Methods, Married Women

 Data Not Available

Data Notes

*Indicator definitions that differ from those in the data notes section are identified by footnotes accompanying the indicator tables.

Demographic Indicators	
Total Population	The number of people in a given area (i.e., country) in a particular time period (usually a midyear estimate).
Population Growth Rate	The average annual growth rate is the rate of natural increase in a population plus the net migration rate. The rate of natural increase is the difference between the birth rate and the death rate, but it is conventionally measured in percentage terms (per hundred rather than per thousand).
Percent Urban	The percentage of the midyear population living in areas defined as urban in each country and reported to the United Nations.
Women, 15-19	The number of women between the ages of 15-19 in the midyear population.
Women, 15-49	The number of women between the ages of 15-49 in the midyear population.
Life Expectancy at Birth	The average number of years that a person at age 0 will live if age-specific death rates remain constant. Life expectancy at birth is highly affected by rates of infant and child death.
Crude Birth Rate	The number of births per thousand of the population. The product of the number of live births, divided by the midpoint population, multiplied by 1,000.
Crude Death Rate	The number of deaths per thousand of the population. The product of the number of deaths divided by the midpoint population, multiplied by 1,000.
Number of Live Births	The number of live births, annually, within a country.
Healthy Life Expectancy: Female	The number of years of in full health that a female newborn can expect to live based on current rates of ill-health and mortality. HALE is based on life expectancy at birth but includes an adjustment for time spent in poor health.
Healthy Life Expectancy: Male	The number of years of in full health that a male newborn can expect to live based on current rates of ill- health and mortality. HALE is based on life expectancy at birth but includes an adjustment for time spent in poor health.
Population: Percent < Age 15	Percent of population less than 15 years of age.
Socioeconomic Indicators	
GNI per Capita (PPP)	The gross national income (GNI) converted to international dollars using purchasing power parity (PPP) rates. An international dollar has the same purchasing power over GNI as a U.S. dollar has in the United States.
Health Expenditure as Percentage of GDP	Total health expenditures is the sum of public and private health expenditures figured as a percentage of a country's gross domestic product (GDP). It covers the provision of health services (preventive and curative), family planning activities, nutrition activities, and emergency aid designated for health but does not include provision of water and sanitation.
Physicians per 1,000 People	Physicians are defined as graduates of any faculty or school of medicine who are working in the country in any medical field (practice, teaching, research).
Adult Literacy Rate	The percentage of people ages 15 and over who can, with understanding, read and write a short, simple statement about their everyday life.
Adult Literacy Rate, Female	The percentage of women ages 15 and over who can, with understanding, read and write a short, simple statement about their everyday life.

Adult Literacy Rate, Male	The percentage of men ages 15 and over who can, with understanding, read and write a short, simple
-	statement about their everyday life.
Gross Enrollment Ratio - Primary School	Total enrolment in primary level of education, regardless of age, expressed as a percentage of the population in the official age group corresponding to primary level of education. The GER can exceed 100% due to late entry or/and repetition.
Gender Parity Index - Net Enrollment Ratio	The ratio of the female-to-male values (or male to female, in certain cases) of net primary school enrollment rates (NER). NER measures the number of pupils in the official age group for a given level of ecudation, expressed as a percentage of the population in that age group.
Access to an Improved Water Source (Rural)	Access to an improved water source in rural areas. Refers to the percentage of the rural population with reasonable access to an adequate amount of water from an improved source, such as a household connection, public standpipe, borehole, protected well or spring, or rainwater collection. Unimproved sources include vendors, tanker trucks, and unprotected wells and springs.
Access to an Improved Water Source (Urban)	Access to an improved water source in urban areas. Refers to the percentage of the urban population with reasonable access to an adequate amount of water from an improved source, such as a household connection, public standpipe, borehole, protected well or spring, or rainwater collection. Unimproved sources include vendors, tanker trucks, and unprotected wells and springs.
Access to Improved Sanitation Facilities (Rural)	Access to improved sanitation facilities in rural areas. Refers to the percentage of the rural population with access to at least adequate excreta disposal facilities (private or shared, but not public) that can effectively prevent human, animal, and insect contact with excreta.
Access to Improved Sanitation Facilities (Urban)	Access to improved sanitation facilities in urban areas. Refers to the percentage of the urban population with access to at least adequate excreta disposal facilities (private or shared, but not public) that can effectively prevent human, animal, and insect contact with excreta.
Population Living Below \$1 a Day	The percentage of the population living below the specified poverty line of \$1 a day at 1985 international prices (equivalent to \$1.08 at 1993 international prices), adjusted for purchasing power parity.
Human Development Index	The HDI is a composite index that measures the average achievements in a country in three basic dimensions of human development: a long and healthy life, as measured by life expectancy at birth; knowledge, as measured by the adult literacy rate and the combined gross enrollment ratio for primary, secondary, and tertiary schools; and a decent standard of living, as measured by gross domestic product (GDP) per capita in purchasing power parity (PPP) US dollars.
Population Below Poverty Line	National estimates of the percentage of the population falling below the poverty line are based on surveys of sub-groups, with the results weighted by the number of people in each group. Definitions of poverty vary considerably among nations. For example, rich nations generally employ more generous standards of poverty than poor nations.
Real GDP (Growth) Rate	This entry gives the gross domestic product (GDP) growth on an annual basis adjusted for inflation and expressed as a percent.
Income Inequality (Gini index)	This index measures the degree of inequality in the distribution of family income in a country. It is calculated from the Lorenz curve, in which cumulative family income is plotted against the number of families arranged from the poorest to the richest. The index is the ratio of (a) the area between a country's Lorenz curve and the 45 degree helping line to (b) the entire triangular area under the 45 degree line. The more nearly equal a country's income distribution, the closer its Lorenz curve to the 45 degree line and the lower its Gini index, e.g., a Scandinavian country with an index of 25. The more unequal a country's income distributed with perfect equality, the index would be zero; if income were distributed with perfect equality, the index would be zero; if income were distributed with perfect equality.
Unemployment Rate	This entry contains the percent of the labor force that is without jobs. Substantial underemployment might be noted.
Inflation Rate	This entry furnishes the annual percent change in consumer prices compared with the previous year's consumer prices.

Airports	This entry gives the total number of airports or airfields recognizable from the air. The runway(s) may be paved (concrete or asphalt surfaces) or unpaved (grass, earth, sand, or gravel surfaces) but may include closed or abandoned installations. Airports or airfields that are no longer recognizable (overgrown, no facilities, etc.) are not included. Note that not all airports have accomodations for refueling, maintenance, or air traffic control.
Airports with paved runways	This entry gives the total number of airports with paved runways (concrete or asphalt surfaces) by length. For airports with more than one runway, only the longest runway is included according to the following five groups - (1) over 3,047 m, (2) 2,438 to 3,047 m, (3) 1,524 to 2,437 m, (4) 914 to 1,523 m, and (5) under 914 m. Only airports with usable runways are included in this listing. Not all airports have facilities for refueling, maintenance, or air traffic control.
Family Planning Indicators	5
Total Fertility Rate (BUCEN)	The number of children a woman between ages 15-49 would have during her reproductive life, if, for all of her childbearing years she were to experience the age-specific birth rates for that given year.
Total Fertility Rate (DHS)	The number of children a woman between ages 15-49 would have during her lifetime if she were to bear children at the currently observed rates.
Contraceptive Prevalence Rate, Modern Methods, All Women	Percentage of all women ages 15-49 currently using a modern method of contraception. Modern methods include oral contraceptives, IUDs, injectables, female and male sterilization, all emergency contraception, and barrier methods (diaphragm, foam, jelly, male and female condom).
Contraceptive Prevalence Rate, Modern Methods, Married Women	Percent of currently married women ages 15-49 currently using a modern method of contraception. Modern methods include oral contraceptives, IUDs, injectables, female and male sterilization, all emergency contraception, and barrier methods (diaphragm, foam, jelly, male and female condom).
Median Age of Sexual Debut Among Women, Ages 25-49	Median age of first sexual intercourse for women ages 25-49.
Mean Ideal Family Size	Mean ideal number of children for all women, according to number of living children.
Women 20-24 Who Gave Birth Before Age 20	Percentage of women ages 20-24 who have given birth before age 20.
Total Fertility Rate, Rural	In rural areas only, the number of children a woman between ages 15-49 would have during her lifetime if she were to bear children at the currently observed rates.
Unmet Need for Family Planning: Rural	In rural areas, percentage of married women ages 15-49 with unmet need for family planning. Unmet need for spacing includes pregnant women whose pregnancy was mistimed, amenorrheic women who are not using family planning and whose last birth was mistimed, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and say they want to wait 2 or more years for their next birth. Also included are fecund women who are not using any method of family planning and say they are unsure whether they want another child or who want another child but are unsure when to have the birth unless they say it would not be a problem if they discovered they were pregnant in the next few weeks. Unmet need for limiting refers to pregnant women whose pregnancy was unwanted, amenorrheic women whose last child was unwanted, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and say they are unsure who are not using refers to pregnant women whose pregnancy was unwanted, amenorrheic women whose last child was unwanted, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and want no more children.
Unmet Need for Family Planning: Urban	In urban areas, percentage of married women ages 15-49 with unmet need for family planning. Unmet need for spacing includes pregnant women whose pregnancy was mistimed, amenorrheic women who are not using family planning and whose last birth was mistimed, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and say they want to wait 2 or more years for their next birth. Also included are fecund women who are not using any method of family planning and say they are unsure whether they want another child or who want another child but are unsure when to have the birth unless they say it would not be a problem if they discovered they were pregnant in the next few weeks. Unmet need for limiting refers to pregnant women whose pregnancy was unwanted, amenorrheic and who are not using any method of family planning and say they are unsure whose last child was unwanted, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and say they are weeks. Unmet need for limiting refers to pregnant women whose pregnancy was unwanted, amenorrheic and who are not using any method of family planning and say they are unsure whose last child was unwanted, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and want no more children.

Methods	
Equity in use of modern contraception	Contraceptive prevalence rate in the lowest wealth quintile/contraceptive prevalence rate in the highest wealth quintile.
Percent of need satisfied by modern methods of family planning	Number of women in union who are fecund and are currently using modern contraceptive methods to stop or postpone the next pregnancy/number of women in union who are fecund and who desire to either terminate childbearing or to postpone their next birth for 2 years or more.
Maternal Health Indicators	
Maternal Mortality Ratio (WHO/Hill)	The estimated number of women who die as a result of pregnancy or childbirth per 100,000 live births arrived through Hill et al., adjustment procedure depending on the nature of data used. The margins of uncertainty associated with the estimated maternal mortality ratios are very large and the estimates should not be used to monitor trends in the short term (including comparisons between 1995 and 2000 estimates). In addition, cross-country comparisons should be treated with considerable circumspection because different strategies are used to derive the estimates for different countries rendering comparisons fraught with difficulty. For further information please refer to the source.
Maternal Mortality Ratio (DHS)	The estimated number of women who die as a result of pregnancy or childbirth per 100,000 live births, arrived mostly through the "sisterhood method". The data are aggregated based on different time periods ranging from four to ten years preceding the survey. Thus, the data may not be suitable for trend analysis due to over-lapping year ranges of estimates. For further information on methodology please refer to: Stanton, Cynthia, Noureddine Abderrahim, and Kenneth Hill. 1997. DHS Maternal Mortality Indicators: An assessment of Data Quality and Implications for Data Use. Calverton: Macro International Inc., or individual country DHS reports.
Antenatal Care (at least 1 visit)	Percentage of women of reproductive age (15-49) who receive at least one antenatal care visit during pregnancy (in the three-year period preceding the survey).
Antenatal Care (2+ visits)	Percentage of women of reproductive age (15-49) who receive at least two antenatal care visits during pregnancy (in the three-year period preceding the survey).
Antenatal Care (4+ visits)	Percentage of women of reproductive age (15-49) who receive four or more antenatal care visits during pregnancy (in the three-year period preceding the survey).
Assisted Delivery by a Health Professional	The percentage of births/deliveries that occur with the assistance of any trained health professional during the five-year period preceding the survey. May include doctors, nurses, or midwives.
Assisted Delivery by Doctor	The percentage of live births/deliveries that occur with the assistance of a doctor during the five-year period preceding the survey.
Assisted Delivery by Other Health Professional	The percentage of live births/deliveries that occur with the assistance of a nurse or midwife during the five-year period preceding the survey.
Anemia prevalence among women of reproductive age	Number of women of reproductive age (15-49 years) with anemia/number of women of reproductive age (15-49 years).
Equity in skilled attendance at delivery	Percentage of births in the lowest wealth quintile attended by a doctor, nurse, or trained midwife/ percentage of births in the highest wealth quintile attended by a doctor, nurse, or trained midwife.
Child Survival Indicators	
ARI Care Seeking - Children Under 5	Percentage of children under five years who were ill with an acute respiratory infection (ARI), which is associated with cough, rapid breathing and a high fever, during the two weeks preceding the survey, for whom care was sought from a health facility or provider.
ORT Use Rate (ORS, RHS, or Increased Fluids)	Oral Rehydration Therapy (ORT). Percentage of children under age 5 with diarrhea two weeks prior to survey who received increased fluids, oral rehydration solution (ORS) or recommended home solution (RHS).
Infant Mortality Rate (BUCEN)	The estimated annual number of deaths of infants under 12 months in a given year per 1,000 live births in that same year.

Percentage of currently married women ages 15-49 currently using ANY method of contraception.

Contraceptive Prevalence Rate, All

Infant Morality Rate (UNICEF)The under five morality rate is the probability (as expressed as a rate per 1,000 live births) and hidd born in a specified year dying before reaching the age of one if subject to current age specific mortality rates.Infant Morality Rate (DISS)The estimated annual number of deaths of infants under 12 months in a given year per 1,000 live births in that same year.Infant Morality Rate, Females (BUCEN)The estimated annual number of deaths of female infants under 12 months in a given year per 1,000 live births in that same year.Infant Morality Rate (BuCEN)The estimated annual number of male infants under 12 months in a given year per 1,000 live births in that same year.Under 5 Morality Rate (BUCEN)The estimated annual number of male infant sunder 12 months in a given year per 1,000 live births in the five-year period preceding the survey.Under 5 Morality Rate (UNICEF)Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey).Under 5 Morality Rate (UNICEF)Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey).Under 5 Morality Rate (UNICEF)Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey).Under 5 Morality Rate (UNICEF)Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey).Under 5 Morality Rate (UNICEF)Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey).Under 5 Morality Rate (UNICEF)Annual number of deaths that occur in		
(DHS) in that same year (five-year period preceding survey). Infant Mortality Rate, Freenales (BUCCEN) The estimated annual number of deaths of female infants under 12 months in a given year per 1,000 Infant Mortality Rate, Males (BUCEN) The estimated annual number of male infants under 12 months in a given year per 1,000 live births in that same year. Under-S Mortality Rate (BUCEN) Annual number of deaths that occur in children 0-4 years old per 1,000 live births in the five-year period preceding the survey. Under-S Mortality Rate (BUCEN) Annual number of deaths that occur in children 0-4 years old per 1,000 live births in the five-year period preceding survey). Under-S Mortality Rate (UNICEF) Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey). Exclusive Breastfeeding Percentage of children under 4 months exclusively breastfed. Exclusive breastfeeding is defined as growing no food or liquid other than breast milk to the child during the 24-hour period before the survey. Stunted (height-for-age) Percentage of children under age five whose height-for-age is below minus-two standard deviations from the median of the reference population. Under-S Marchality Rate (BUS) Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population. Underweight (weight-for- ge) Percentage of children under age five whose weight-for-leght is below minus-two standard deviations from the m		born in a specified year dying before reaching the age of one if subject to current age-specific mortality
Females (BUCEN) live biths in that same year. Infant Mortaitiy Rate, Males (BUCEN) The estimated annual number of male infants under 12 months in a given year per 1,000 live biths in that same year. Under-5 Mortaity Rate (BUCEN) Annual number of deaths that occur in children 0-4 years old per 1,000 live biths in the five-year period preceding the survey. Under-5 Mortaity Rate (BUCEN) The estimated number of deaths that occur in children 0-4 years old per 1,000 biths (five-year period preceding survey). Under-5 Mortaity Rate (DHS) Annual number of deaths that occur in children 0-4 years old per 1,000 biths (five-year period preceding survey). Exclusive Breastfeeding (under 4 mos.) Annual number of deaths that occur in children 0-4 years old per 1,000 biths (five-year period preceding survey). Exclusive Breastfeeding (under 4 mos.) Percentage of children under 4 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey. Sturted (height-for-age) Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population. Underweight (weight-for- ge) Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population. Underweight (weight-for- ge) Percentage of children under age five whose weight-for-height is below minus-two standard deviations from the	-	
Males (BUCEN) that same year. Under-5 Mortality Rate (BUCEN) Annual number of deaths that occur in children 0-4 years old per 1,000 births. Neonatal Mortality Rate (BUCEN) The estimated number of infant deaths in the first month of life per 1,000 births (five-year period preceding survey). Under-5 Mortality Rate (UNICEF) Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey). Under-5 Mortality Rate (DHS) Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey). Exclusive Breastfeeding (under 4 mos.) Percentage of children under 4 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey. Stunted (height-for-age) Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population. Wasted (weight-for- age) Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population. DPT3 Vaccination Rate (PHS) Proportion of living children 12-23 months old who have received three complete doses of vaccines aginst diphtheria, pertusis, and tetanus. DPT3 Vaccination Rate (PHS) Proportion of living children 12-23 months old who have received three doses of polo vaccine. Polo Vaccination Rate (PHS)		
(EUCEN) Neonatal Mortality Rate (UNICEF) The estimated number of infant deaths in the first month of life per 1,000 live births in the five-year period preceding the survey. Under-5 Mortality Rate (UNICEF) Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey). Exclusive Breastfeeding (under 4 mos.) Percentage of children under 4 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey. Exclusive Breastfeeding (under 6 mos.) Percentage of children under 6 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey. Stunted (height-for-age) Percentage of children under age five whose height-for-age is below minus-two standard deviations from the median of the reference population. Underveight (weight-for- height) Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population. VPT3 Vaccination Rate (DHS) Proportion of living children 12-23 months old who have received three complete doses of vaccines against diphtheria, pertussis, and tetanus. DPT3 Vaccination Rate (DHS) Proportion of living children 12-23 months old who have received one dose of Measles Containing Vaccinee. Proportion of living children 12-23 months who have received three doses of polio vaccine. Proportion of living		
period preceding the survey.Under-5 Mortality Rate (UNICEF)Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey).Under-5 Mortality Rate (UNICEF)Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey).Exclusive Breastleeding (under 4 mos.)Percentage of children under 4 months exclusively breastled. Exclusive breastleeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey.Exclusive Breastleeding (under 6 mos.)Percentage of children under 6 months exclusively breastled. Exclusive breastleeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey.Stunted (height-for-age)Percentage of children under age five whose height-for-age is below minus-two standard deviations from the median of the reference population.Underweight (weight-for- age)Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population.Underweight (weight-for- ageis)Percentage of children under age five whose weight-for-height is below minus-two standard deviations from the median of the reference population.DPT3 vaccination Rate (WHO)Proportion of living children 12-23 months old who have received three complete doses of vaccines against diphtheria, pertussis, and tetanus.Measles Vaccination Rate (WHO)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (PHS)Proportion of living childre	-	Annual number of deaths that occur in children 0-4 years old per 1,000 births.
(UNICEF) preceding survey). Under-S Mortality Rate (DHS) Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey). Exclusive Breastfeeding Percentage of children under 4 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey. Exclusive Breastfeeding Percentage of children under 6 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey. Stunted (height-for-age) Percentage of children under age five whose height-for-age is below minus-two standard deviations from the median of the reference population. Underweight (weight-for- age) Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population. DPT3 Vaccination Rate (WHO) Proportion of living children 12-23 months old who have received three complete doses of vaccines aginst diphtheria, pertussis, and tetanus. Measles Vaccination Rate (WHO) Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince. Measles Vaccination Rate (WHO) Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince. Measles Vaccination Rate (WHO) Percentage of living children 12-23 months who have received three doses of po	Neonatal Mortality Rate	
(DHS)preceding survey).Exclusive Breastfeeding (under 4 mos.)Percentage of children under 4 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey.Exclusive Breastfeeding (under 6 mos.)Percentage of children under 6 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey.Exclusive Breastfeeding (under 6 mos.)Percentage of children under age five whose height-for-age is below minus-two standard deviations from the median of the reference population.Underweight (weight-for- age)Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population.Underweight (weight-for- age)Percentage of children under age five whose weight-for-height is below minus-two standard deviations from the median of the reference population.DPT3 Vaccination Rate (DHS)Proportion of living children 12-23 months old who have received three complete doses of vaccines aginst diphtheria, pertussis, and tetanus.Meastes Vaccination Rate (DHS)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Meastes Vaccination Rate (DHG)Percentage of living children 12-23 months old who have received three doses of polio vaccine.(DHS)Percentage of living children 12-23 months old who have received three doses of Measles Containing Vaccince.Meastes Vaccination Rate (UHG)Percentage of pregnant women receiving two (or more) doses of tetanu	-	
(under 4 mos.)providing no food or liquid other than breast milk to the child during the 24-hour period before the survey.Exclusive Breastfeeding (under 6 mos.)Percentage of children under 6 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey.Stunted (height-for-age)Percentage of children under age five whose height-for-age is below minus-two standard deviations from the median of the reference population.Underweight (weight-for- age)Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population.Wasted (weight-for- height)Percentage of children under age five whose weight-for-height is below minus-two standard deviations from the median of the reference population.DPT3 Vaccination Rate (DHS)Proportion of living children 12-23 months old who have received three complete doses of vaccines against diphtheria, pertussis, and tetanus.DPT3 Vaccination Rate (DHS)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Measles Vaccination Rate (DHS)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vac	-	
(under 6 mos.)providing no food or liquid other than breast milk to the child during the 24-hour period before the survey.Stunted (height-for-age)Percentage of children under age five whose height-for-age is below minus-two standard deviations from the median of the reference population.Underweight (weight-for- age)Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population.Wasted (weight-for- height)Percentage of children under age five whose weight-for-height is below minus-two standard deviations from the median of the reference population.DPT3 Vaccination Rate (DHS)Proportion of living children 12-23 months old who have received three complete doses of vaccines against diphtheria, pertussis, and tetanus.DPT3 Vaccination Rate (WHO)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Measles Vaccination Rate (DHS)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (OHS)Proportion of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (DHS)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (OHS)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (OHS)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (OHS)Proportion of living children 12-23 months who have		providing no food or liquid other than breast milk to the child during the 24-hour period before the
Indexweight (weight-for- age)For the median of the reference population.Underweight (weight-for- age)Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population.Wasted (weight-for- height)Percentage of children under age five whose weight-for-height is below minus-two standard deviations from the median of the reference population.DPT3 Vaccination Rate (DHS)Percentage of children 12-23 months old who have received three complete doses of vaccines against diphtheria, pertussis, and tetanus.DPT3 Vaccination Rate (WHO)Proportion of living children 12-23 months old who have received three complete doses of vaccines aginst diphtheria, pertussis, and tetanus.Measles Vaccination Rate (DHS)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Measles Vaccination Rate (UHO)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (DHS)Proportion of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (DHS)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Percentage of pregnant women receiving two (or more) doses of tetanus toxid (three-year period percentage of pregnant women receiving two (or more) doses of tetanus toxid (three-year period percentage of pregnant women receiving two (or more) doses of tetanus toxid (three-year period percentage of yaccinationPortacination / (DHS)DPT3 coverage in the lowest wealth quintile/		providing no food or liquid other than breast milk to the child during the 24-hour period before the
age)from the median of the reference population.Wasted (weight-for- height)Percentage of children under age five whose weight-for-height is below minus-two standard deviations from the median of the reference population.DPT3 Vaccination Rate (DHS)Proportion of living children 12-23 months old who have received three complete doses of vaccines against diphtheria, pertussis, and tetanus.DPT3 Vaccination Rate (WHO)Proportion of living children 12-23 months old who have received three complete doses of vaccines aginst diphtheria, pertussis, and tetanus.Measles Vaccination Rate (DHS)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Measles Vaccination Rate (WHO)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (DHS)Proportion of living children 12-23 months who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (DHS)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination (HS)Percentage of pregnant women receiving two (Stunted (height-for-age)	
height)from the median of the reference population.DPT3 Vaccination Rate (DHS)Proportion of living children 12-23 months old who have received three complete doses of vaccines against diphtheria, pertussis, and tetanus.DPT3 Vaccination Rate (WHO)Proportion of living children 12-23 months old who have received three complete doses of vaccines aginst diphtheria, pertussis, and tetanus.Measles Vaccination Rate (DHS)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Measles Vaccination Rate (DHS)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Measles Vaccination Rate (WHO)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (DHS)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (DHS)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (DHS)Proportion of living children 12-23 months who have received three doses of polio vaccine. <t< td=""><td></td><td></td></t<>		
(DHS)against diphtheria, pertussis, and tetanus.DPT3 Vaccination Rate (WHO)Proportion of living children 12-23 months old who have received three complete doses of vaccines aginst diphtheria, pertussis, and tetanus.Measles Vaccination Rate (DHS)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Measles Vaccination Rate (WHO)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (DHS)Percentage of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Porportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Porportion of living children 12-23 months who have received three doses of polio vaccine.Dette Social Context Context Context Context Context Context Context Con	· •	
(WHO)alignst diphtheria, pertussis, and tetanus.Measles Vaccination Rate (DHS)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Measles Vaccination Rate (WHO)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (DHS)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination (WHO)Percentage of pregnant women receiving two (or more) doses of tetanus toxoid (three-year period preceding survey).Equity in DPT3 coverage (DHS)DPT3 coverage in the lowest wealth quintile/DPT3 coverage in the highest wealth quintile.Improved sanitation/ hygiene practicesNumber of households with child under 5 years of age whose youngest child's feces was safely disposed of the last time he/she passed stool/total number of households with a child under 5 years of<		
Rate (DHS)Vaccince.Measles Vaccination Rate (WHO)Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.Polio Vaccination Rate (DHS)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination (WHO)Percentage of pregnant women receiving two (or more) doses of tetanus toxoid (three-year period preceding survey).Equity in DPT3 coverage (DHS)DPT3 coverage in the lowest wealth quintile/DPT3 coverage in the highest wealth quintile.Improved sanitation/ hygiene practicesNumber of households with child under 5 years of age whose youngest child's feces was safely disposed of the last time he/she passed stool/total number of households with a child under 5 years of		
Rate (WHO)Vaccination (WHO)Polio Vaccination Rate (DHS)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Polio Vaccination Rate (WHO)Percentage of pregnant women receiving two (or more) doses of tetanus toxoid (three-year period preceding survey).Equity in DPT3 coverage (DHS)DPT3 coverage in the lowest wealth quintile/DPT3 coverage in the highest wealth quintile.Improved sanitation/ hygiene practicesNumber of households with child under 5 years of age whose youngest child's feces was safely disposed of the last time he/she passed stool/total number of households with a child under 5 years of		
(DHS)Polio Vaccination Rate (WHO)Proportion of living children 12-23 months who have received three doses of polio vaccine.Tetanus Toxoid VaccinationPercentage of pregnant women receiving two (or more) doses of tetanus toxoid (three-year period preceding survey).Equity in DPT3 coverage (DHS)DPT3 coverage in the lowest wealth quintile/DPT3 coverage in the highest wealth quintile.Improved sanitation/ hygiene practicesNumber of households with child under 5 years of age whose youngest child's feces was safely disposed of the last time he/she passed stool/total number of households with a child under 5 years of		
(WHO)Tetanus Toxoid VaccinationPercentage of pregnant women receiving two (or more) doses of tetanus toxoid (three-year period preceding survey).Equity in DPT3 coverage (DHS)DPT3 coverage in the lowest wealth quintile/DPT3 coverage in the highest wealth quintile.Improved sanitation/ hygiene practicesNumber of households with child under 5 years of age whose youngest child's feces was safely disposed of the last time he/she passed stool/total number of households with a child under 5 years of		Proportion of living children 12-23 months who have received three doses of polio vaccine.
Vaccination preceding survey). Equity in DPT3 coverage (DHS) DPT3 coverage in the lowest wealth quintile/DPT3 coverage in the highest wealth quintile. Improved sanitation/ hygiene practices Number of households with child under 5 years of age whose youngest child's feces was safely disposed of the last time he/she passed stool/total number of households with a child under 5 years of		Proportion of living children 12-23 months who have received three doses of polio vaccine.
(DHS) Improved sanitation/ hygiene practices Number of households with child under 5 years of age whose youngest child's feces was safely disposed of the last time he/she passed stool/total number of households with a child under 5 years of		
hygiene practices disposed of the last time he/she passed stool/total number of households with a child under 5 years of		DPT3 coverage in the lowest wealth quintile/DPT3 coverage in the highest wealth quintile.
	-	disposed of the last time he/she passed stool/total number of households with a child under 5 years of

Vitamin A supplementation coverage among children 6-59 months

Number of children aged 6-59 months who received a vitamin A supplement in the last 6 months/total number of children aged 6-59 months surveyed.

HIV/AIDS Prevention Indicators

HIV Prevalence proportion: Adults (15–49 years)	The adult HIV prevalence proportion is the estimated number of adults living with HIV divided by the adult population (aged 15–49).
Estimated number of people living with HIV: Adults and Children	Estimated number of adults and children living with HIV. Adults are 15 years and over. Children are defined as those aged 0–14 years.
Estimated number of people living with HIV: Women (15+ years)	Estimated number of women (aged 15 and over) living with HIV.
Estimated number of people living with HIV: Children (0–14 years)	Estimated number of children under age 15 living with HIV.
Males Reporting Condom Use With Last Non- Regular Partner	Among men who have had high risk sex in the last year, percentage who used condoms during their last sexual intercourse with a non-regular partner/non-cohabitating partner.
Females Reporting Condom Use With Last Non-Regular Partner	Among women who have had high risk sex in the last year, percentage who used condoms during their last sexual intercourse with a non-regular partner/non-cohabitating partner.
Infectious Diseases Contro	ol Indicators
TB Estimated Number of Cases	The estimated number of cases (all forms).
TB Case Detection Rate	TB Case Detection Rate is defined as the percentage of the annual new smear-positive notifications of the estimated annual new smear-positive incidence.
TB (DOTS) Treatment Success Rate	The proportion of smear-positive patients who were cured plus the proportion who completed treatment

Malaria Cases per Number of malaria cases per 100,000 of the population during the year for which data was reported.

DOTS Coverage The percentage of the national population living in areas where health services have adopted DOTS. "Areas" are the lowest administrative or management units in the country (townships, districts, counties, etc). If an area is considered a DOTS area, then all cases registered and reported in that area are considered DOTS cases and the population living within the boundaries of that area counts toward national DOTS coverage. As a measure of patient access to diagnosis and treatment under DOTS, coverage is an approximation, and usually an overestimate.

HIV-Infected with Active
TBThe TB incidence rate in HIV-infected people 15-49 divided by the incidence rate in HIV-uninfected
people 15-49.

Avian Influenza: Cumulative Number of Confirmed Human Cases

Avian Influenza: Cumulative Number of Confirmed Human Deaths

100,000

Child Sleeps Under Insecticide-Treated Net The cumulative number of confirmed human cases of Avian Influenza A/(H5N1) reported to WHO. The total number of cases includes number of deaths. WHO reports only laboratory-confirmed cases.

The cumulative number of confirmed human deaths from Avian Influenza A/(H5N1) reported to WHO. WHO reports only laboratory-confirmed cases and deaths.

s Under Treated NetPercentage of children under five years of age who slept under an Insecticde Treated Net (ITN) the night before the survey. An ITN is (1) a permanent net that does not require any treatment, or (2) a pretreated net obtained within the last six months, or (3) a net that has been soaked with insecticide within the past six months.

Equity in Bednet (ITN) P Ownership h

Proportion of households in the lowest income quintile with an ITN / proportion of households in the highest income quintile with an ITN.

Statistical Sources For Bulgaria

BUCEN-IDB 2006	United States Census Bureau (BUCEN), International Programs Center. International Database. August 2006.
BUCEN-IDB 2008	United States Census Bureau (BUCEN), International Programs Center. International Database. February 2008.
CIA World Factbook 2008	CIA World Factbook, 2008 https://cia.gov/cia//publications/factbook/docs/profileguide. html. Last accessed April, 2008.
Human Development Report 2006	Human Development Report 2006. New York, United Nations Development Programme, 2006. Available at: http://hdr.undp.org/hdr2006/. Last accessed February, 2007.
Human Development Report - 2007/2008 2008	Human Development Report 2007/2008. Fighting climate change: Human solidarity in a divided world. Published for the United Nations Development Programme (UNDP)
UNESCO EFA Global Monitoring Report 2008	EFA Global Monitoring Report 2008. Education for All by 2015 - Will we make it? United Nations Educational, Scientific and Cultural Organization, Paris.
UNICEF 2008	UNICEF. Progress since the World Summit for Children. http://www.childinfo.org/areas/ childmortality/u5data.php. Data accessed on web site in March, 2008
WHO Global Health Atlas 2008	Accessed April of 2008
WHO Global Health Atlas 2008	Accessed March of 2008
WHO World Health Report 2004	The World Health Report 2004: Changing history. Geneva, World Health Organization, 2004. Available at: http://www.who.int/whr/2004/en/index.html. Last accessed February, 2007.
WHO/Global Summary 2002	WHO Vaccine-Preventable Diseases Monitoring System. 2002 Global Summary. World Health Organization, Geneva.
WHO/Global Summary 2008	Immunization surveillance, assessment and monitoring section of the WHO website: http://www.who.int/immunization_monitoring/en/globalsummary/timeseries/ tscoveragedtp3.htm
WHO/Global Summary 2008	Immunization surveillance, assessment and monitoring section of the WHO website: http://www.who.int/immunization_monitoring/en/globalsummary/timeseries/ tscoveragemcv.htm
WHO/Global Summary 2008	Immunization surveillance, assessment and monitoring section of the WHO website: http://www.who.int/immunization_monitoring/en/globalsummary/timeseries/ tscoveragepol3.htm
WHO/Hill 2005	Maternal Mortality in 2005: Estimates developed by WHO, UNICEF, UNFPA and The World Bank
World Bank/WDI 2005	World Bank Development Indicators 2005, CD-Rom. Washington, DC: The World Bank, 2005.
World Bank/WDI 2006	World Bank Development Indicators 2006. Washington, DC: The World Bank, 2005. Available at: http://devdata.worldbank.org/wdi2006/contents/index2.htm. Last accessed April, 2007.

World Bank Development Indicators Database 2007. Washington, DC: The World Bank, 2007. Available at: http://devdata.worldbank.org/data-query/. Last accessed April, 2007.