

The Concentration–Response Relation between PM_{2.5} and Daily Deaths

Joel Schwartz,^{1,2,3} Francine Laden,^{1,3} and Antonella Zanobetti¹

¹Environmental Epidemiology Program, Department of Environmental Health, Harvard School of Public Health, Boston, Massachusetts, USA; ²Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts, USA; ³Channing Laboratory, Brigham and Women's Hospital, and Department of Medicine, Harvard Medical School, Boston, Massachusetts, USA

Particulate air pollution at commonly occurring concentrations is associated with daily deaths. Recent attention has focused on the shape of the concentration–response curve, particularly at low doses. Several recent articles have reported that particulate matter with aerodynamic diameter $\leq 10 \mu\text{m}$ (PM₁₀) was associated with daily deaths with no evidence of a threshold. These reports have used smoothing or spline methods in individual cities and pooled the results across multiple cities to obtain estimates that are more robust. To date, fine particulate matter (aerodynamic diameter $\leq 2.5 \mu\text{m}$; PM_{2.5}), a component of PM₁₀, has not been examined in this regard. We examined this association in a hierarchical model in six U.S. cities. In the first stage, we fit log-linear models including smooth functions of PM_{2.5} in each city, controlling for season, weather, and day of the week. These smooth functions allowed for nonlinearities in the city-specific associations. We combined the estimated curves across cities using a hierarchical model that allows for heterogeneity. We found an essentially linear relationship down to $2 \mu\text{g}/\text{m}^3$. The same approach was applied to examine the concentration response to traffic particles, controlling for particles from other sources. Once again, the association showed no sign of a threshold. The magnitude of the association suggests that controlling fine particle pollution would result in thousands fewer early deaths per year. **Key words:** meta-analysis, mortality, particulate air pollution, smoothing, time series, traffic. *Environ Health Perspect* 110:1025–1029 (2002). [Online 27 August 2002] <http://ehpnet1.niehs.nih.gov/docs/2002/110p1025-1029schwartz/abstract.html>

In the last decade, a series of studies reported associations between daily concentrations of airborne particles and daily deaths (1–3). The magnitude of the regression coefficients in those studies indicated that particulate air pollution was associated with between 50 and 100,000 early deaths per year in the United States, and similar numbers were found in Europe. More recently, a number of large, multicity studies (4–7) have reported associations between airborne particles, measured in various ways, and daily deaths. The largest study demonstrated that gaseous air pollutants did not confound the association, and that none of the gaseous air pollutants showed an independent effect on daily deaths (7). These studies assumed a linear concentration–response relation between airborne particles and daily deaths and did not address the question of what the association looked like for particle constituents, characterized by size, physiochemical composition, or source.

In a recent study of six U.S. cities (5), we demonstrated that daily mortality was associated with fine particulate matter (aerodynamic diameter $\leq 2.5 \mu\text{m}$; PM_{2.5}) and not with coarse particulate matter (aerodynamic diameter between 2.5 and $10 \mu\text{m}$; PM_{2.5–10}). Each $10 \mu\text{g}/\text{m}^3$ increase in the 2-day mean concentration of PM_{2.5} was associated with a 1.5% (95% confidence interval, 1.1–1.9%) increase in daily mortality.

Ambient PM_{2.5} consists mainly of combustion particles from motor vehicles and the burning of coal, fuel oil, and wood, but also

contains some crustal particles from finely pulverized road dust and soils. These sources produce particles with different characteristics, and the relative toxicity of those sources and characteristics is an area of relative recent but intense interest. In a follow-up study (8), we used the elemental composition of size-fractionated particles to identify several distinct source-related fractions of fine particles. We then examined the association of these fractions with daily mortality in each of the six cities and combined the city-specific results in a meta-analysis to derive overall relative risks for each fraction. We found positive associations with particles from traffic, particles from coal, and particles from residual oil combustion when included jointly in the model predicting daily deaths (8). The largest effect size was for residual oil particles, followed by traffic particles and then coal particles. Only the latter two associations were statistically significant, however. Again, as traditional, these analyses assumed a linear association between the various particle constituents and daily deaths.

The shape of the concentration–response relationship is critical for public health assessment, and in particular, some have speculated that thresholds might exist.

Recently, three reports have explored this question for particulate air pollution, using multicity studies in the United States. In one study, Daniels et al. (9) used data from 20 U.S. cities, five of which had daily measurements of PM₁₀, with the rest having measurements only one day in six. They used regression

splines to model the concentration–response curve in each city and combined the results across cities. They found no evidence for a threshold. In fact, the concentration–response relation was quite linear across the entire range of exposure. In another report, Schwartz and Zanobetti (10) used data from 10 cities, all of which had daily measurements of PM₁₀, resulting in slightly more days of study than in the first report. They used non-parametric smoothing to model the concentration–response curve between air pollution and daily deaths in each city and combined the results across cities. Again, a linear, no-threshold relationship was seen. Schwartz and Zanobetti also performed simulations to confirm the ability of this approach to detect thresholds and other types of nonlinearity (10). Schwartz et al. (11), using data from eight Spanish cities, similarly reported a linear association between daily deaths and black smoke, an optical measure of black particles. These results held after adjusting for SO₂. To date, no similar examination of the concentration–response curve has been done for PM_{2.5}, or for any source components. Because PM_{2.5} is now the regulated form of particulate air pollution in the United States, we here report results of such an analysis.

Materials and Methods

Air pollution data. As part of the Harvard Six Cities studies (12), dichotomous virtual impactor samplers were placed at a central residential monitoring site in six U.S. metropolitan areas: Boston, Massachusetts; Knoxville, Tennessee; St. Louis, Missouri; Stuebenville, Ohio; Madison, Wisconsin; and Topeka, Kansas. Separate filter samples were collected of fine particles (PM_{2.5}) and of the coarse mass (PM_{2.5–10}) fraction. Integrated 24-hr samples were collected at least every other day from 1979 until the late 1980s, with daily sampling during health survey periods. For fine and coarse particle samples, mass concentration was determined separately by beta-attenuation (13). Except for a period

Address correspondence to J. Schwartz, Environmental Epidemiology Program, Harvard School of Public Health, 665 Huntington Ave., Boston, MA 02115 USA. Telephone: (617) 384-8752. Fax: (617) 384-8745. E-mail: jschwartz@hsph.harvard.edu

This work was supported by U.S. EPA Grant R827353 and NIEHS Grant ES 00002.

Received 21 December 2001; accepted 20 March 2002.

between October 1981 and January 1984 in all cities, elemental composition of fine and coarse mass was determined by X-ray fluorescence (14). Elemental composition was available on 97% of these samples. In the fine fraction, 15 elements were routinely found above the limit of detection: silicon, sulfur, chlorine, potassium, calcium, vanadium, manganese, aluminum, nickel, zinc, selenium, bromine, lead, copper, and iron.

Source identification. In separate analyses for each city, we used specific rotation factor analysis to identify up to five common factors from the 15 specified elements. We specified a single element as the tracer for each factor and maximized the projection of these elements using the Procrustes rotation, a variant of the oblique rotation method (15). The Procrustes method allows us to use known tracers for different sources as targets for the different factors and to maximize their loadings on those factors instead of having factors defined in an entirely data-driven manner. To rescale the factor scores from the normalized scale to the mass scale (in micrograms per cubic meter), we regressed the total daily fine particle concentrations on the daily factor scores for all of the factors in separate regression models for each city and took the product of each factor score with its regression coefficient (16). Only sources that were significant predictors of total fine particle mass ($p < 0.10$) were considered in the mortality analyses. Further details have been published previously (8).

Meteorologic data. We obtained meteorologic data from the National Center for Atmospheric Research, including hourly measures of temperature, dew point temperature, and precipitation from the National Oceanographic and Atmospheric Administration weather station nearest to each city (17). We calculated 24-hr mean values for temperature and dew point temperature.

Mortality data. We defined the six metropolitan areas in this study as the county containing the air pollution monitor and contiguous counties (5). We extracted daily deaths from annual detail mortality tapes (National Center for Health Statistics) (18) for people who lived and died in the selected counties for the time periods with fine particulate measurements. After excluding all deaths caused by accidents and other external causes [International Classification of Diseases, 9th Revision (ICD-9) (19), clinical modification codes 800–999], we analyzed the remaining total daily deaths.

Poisson regression of mortality. We investigated the association of daily deaths with sources of fine particles separately for each city using Poisson regression in a generalized additive model (GAM) (20,21). That is, in each city we assumed

$$\text{Log}[E(Y_t)] = \beta_0 + \sum S_i(X_{it}), \quad [1]$$

where Y_t is the number of deaths in the city on day t and X_{it} is the value of covariate i on day t . GAMs are distinguished by allowing us to use smooth functions S_i instead of linear terms to control for covariates, such as temperature, that may affect daily deaths in a nonlinear way. Linear functions may be used where appropriate. This approach was introduced for time series of counts in 1994 (22) and is now standard (23,24).

To control for trend and season, we used a locally weighted linear regression (LOESS) smooth function of date with a span of 0.05 (25). For the smooth functions of temperature and dew point temperature, we used LOESS functions with spans of 0.80. Indicator variables for day of the week also were included in the models. This is the identical model used by Schwartz et al. (5) and Laden et al. (8), and more details are provided there. To these models we added a smooth function of the mean $\text{PM}_{2.5}$ concentration on the day of death and

the previous day, instead of the linear term previously used by Schwartz et al. (5). The smoothing window included 50% of the data, which corresponds to between four and five degrees of freedom for the air pollution relation in each city. Alternatively, we added the estimated mass for each of the source factor scores (in micrograms per cubic meter) simultaneously in the model. That is, the estimate of the mobile source factor is in a model controlling for coal-derived particles, crustal particles, and the other source factors, and vice versa. Because only the particles from traffic showed a strong linear association, and because the exposure ranges for the exposures to coal particles did not overlap sufficiently, we only used a smooth function for the traffic particles and followed Laden et al. (8) in treating the particle mass from the other sources as linear terms.

Hierarchical model. To combine the smooth curves across cities, we applied the approach of Schwartz and Zanobetti (10), as modified by Schwartz et al. (11). In each city, the predicted log relative risk and its pointwise standard error was computed for each 2

Table 1. Mean daily deaths in six U.S. cities and mean concentrations of $\text{PM}_{2.5}$ overall, and from the three source categories showing evidence of an association with daily deaths in Laden et al. (8).

City	Deaths	$\text{PM}_{2.5}$ ($\mu\text{g}/\text{m}^3$)	Traffic ($\mu\text{g}/\text{m}^3$)	Coal ($\mu\text{g}/\text{m}^3$)	Residual oil ($\mu\text{g}/\text{m}^3$)	Dates (month/year)
Boston	59	16.5	4.8	8.3	0.5	5/79–1/86
Knoxville	12	21.1	4.4	6.8	—	1/80–12/87
St. Louis	55	19.2	2.9	5.6	—	9/79–1/87
Steubenville	3	30.5	1.5	19.2	0.9	4/79–9/87
Madison	11	11.3	3.1	4.9	—	3/79–12/97
Topeka	3	12.2	2.1	7.0	—	9/79–10/88




Figure 1. Overall estimated dose–response relation between total $\text{PM}_{2.5}$ and daily deaths in six U.S. cities. The estimate is obtained by combining the estimated smoothed curves in each of the cities, after controlling for weather, season, and day of the week. The shaded area indicates the pointwise 95% confidence intervals at each point. The line shown is a least-squares regression line through the estimated points.

$\mu\text{g}/\text{m}^3$ increment in exposure. These estimates are provided by the GAM function in S-plus (MathSoft, Inc., Seattle, WA). To successfully combine data across cities, we need to use a range of exposures that is common to all cities. Because high concentrations of PM_{2.5} were rare, the curves were combined only in the range of 0–35 $\mu\text{g}/\text{m}^3$. The first phase of the analysis produced estimated effect sizes (log relative risks) \hat{Y}_{ij} in each city i for each exposure category j . A pointwise standard error of the estimate is also estimated by GAM. To produce the combined curve, we regressed these estimates against indicator variables for each level, using inverse variance weighting and allowing for a random variance component to capture heterogeneity in the association across cities. That is, we assumed

$$\hat{Y}_{ij} \sim N(\beta_1 d_1 + \beta_2 d_2 + \dots + \beta_k d_k, V_{ij} + \delta), \quad [2]$$

where d_j are dummy variables for the j exposure levels, V_{ij} is the estimated variance in city i at level j , and δ is the estimated random variance component.

We used the iterative meta-regression approach of Berkey et al. (26) to obtain a maximum likelihood estimate of the random variance component.

The nonparametric smooth functions we use to estimate the shape of the concentration response relation use four to five degrees of freedom, and it is not clear that the source-specific relations can support so many degrees

of freedom, which would entail a total of 20 degrees of freedom for all the PM_{2.5} sources. In our previous report (8), the relation between PM_{2.5} from traffic and daily deaths was estimated with considerably greater precision than for particles from other sources, most of which were not significant. Further, the range of overlap in exposures across cities was lower for coal, crustal, and residual oil factors. Therefore, in our source-specific models, we only modeled the traffic source particles using a nonparametric smooth, while controlling for PM_{2.5} from the other sources using linear terms, as in Laden et al. (8). We then combined the estimated concentration–response relations for traffic particles similarly to what we did for PM_{2.5} from the other sources.

Results

Table 1 shows the daily deaths, PM_{2.5} levels, and estimated concentrations of PM_{2.5} from each source. Figure 1 shows the meta-smooth dose–response relation between PM_{2.5} and daily deaths in the six cities. There is no evidence of a threshold, and the relation occurs well below the U.S. Environmental Protection Agency standard of 65 $\mu\text{g}/\text{m}^3$ (27). The line shows the least-squares fit of a linear relation through the estimated points.

The next results come from the source component models. These models had a smooth function of PM_{2.5} from traffic and linear functions of PM_{2.5} from the other sources in each city. Figure 2 shows the results when we combined the estimated

dose–response curves for traffic particles across the six cities. Again, there is no evidence of a threshold, and the association is essentially linear. If anything, the slope is steeper at lower concentrations. To test the robustness of the association with traffic particles to our method of controlling for particles from other sources, we re-estimated the relationship controlling for smooth functions of the estimated particle mass from other sources, rather than the linear terms. This association is shown in Figure 3 and differs little from that shown in Figure 2. We also fit linear regressions through the points shown on Figures 1 and 2. We obtained a slope of 1.5% increase in deaths per 10 $\mu\text{g}/\text{m}^3$ increase in PM_{2.5} and 3% increase in deaths per 10 $\mu\text{g}/\text{m}^3$ increase in particles from traffic, which is the same as the results reported by Laden et al. (8). These lines are shown on the figures. This supports the assumption of a linear relationship.

Discussion

We have explored the concentration–response relation between PM_{2.5} and daily deaths in six U.S. cities and combined the results to obtain greater stability, while accounting for heterogeneity in response. The population mean curve shows no evidence of a threshold down to the lowest levels of PM_{2.5}. In fact, the curve is quite linear over the exposure range from 0 to 35 $\mu\text{g}/\text{m}^3$. This is consistent with previous results using a similar methodology but with PM₁₀ (10) and black smoke (11) as the exposure metric. In addition, a different methodology, using regression splines, was applied by Daniels et al. (9) to PM₁₀ data in different cities. They combined these spline models across 20 cities. Again, the association appeared to be quite linear without any evidence of a threshold. A spline model had previously been applied by Schwartz (22) to the PM_{2.5} data from Boston, with a similar finding. Indeed, the original study of these data by Schwartz, Dockery, and Neas (5) found a significant association when limited to days below 30 $\mu\text{g}/\text{m}^3$, with a slightly larger slope. The consistency of the results on two continents, and using different techniques, suggests that this finding is robust. The concentration–response curve seen here for PM_{2.5} is steeper than that previously reported (per $\mu\text{g}/\text{m}^3$) for PM₁₀ (10). This is consistent with the previous report from this study (5) that coarse mass (the difference between PM₁₀ and PM_{2.5}) is not associated with daily deaths. We note that Schwartz and Zanobetti (10) demonstrated in simulation studies that measurement error was not likely to distort the shape of the association. Similarly, recent studies of “harvesting” have shown that effect sizes increase rather than decrease when longer lags are taken into account; for example, high




Figure 2. Overall estimated dose–response relation between PM_{2.5} from traffic and daily deaths in six U.S. cities. The estimate is obtained by combining the estimated smoothed curve in each of the cities, after controlling for weather, season, and day of the week and for PM_{2.5} from crustal sources, coal combustion, residual oil, salt, and metal processes as linear terms. The line shown is a least-squares regression line through the estimated points.

days producing harvesting that mutes the effect on the next high day is unlikely to have distorted the shape of the association.

These results are also biologically plausible. Schwartz (28) pointed out that if thresholds exist in individuals, but there is a distribution of those thresholds among individuals, and if multiple genetic and predisposing illnesses each contributed to the distribution of those thresholds, then by the central limit theorem, the distribution of thresholds should approach a normal distribution. Hence, the population concentration–response curve should approach a cumulative normal curve. But the low-dose end of the cumulative normal curve is linear. To see this, consider that typical death rates in U.S. cities are 8/1,000 per year, or 2×10^{-6} per day. The normal range of variation in daily deaths in U.S. cities is a factor of two or less. Hence, the normal range of daily death probabilities in response to all risk factors is from 1 to 3×10^{-6} . Figure 4 shows the cumulative normal curve in that range of probabilities, which is quite linear. Because we are clearly in the low-dose regime, in the sense that the exposures to particles are well below the threshold for mortality for most people, this linearity is exactly what would be expected.

Figure 1 also indicates that the association reported here has public health significance. The difference between mean $PM_{2.5}$ concentrations of $10 \mu\text{g}/\text{m}^3$ and $20 \mu\text{g}/\text{m}^3$, which is a difference found between U.S. cities, is associated with about a 1.5% increase in deaths. In a metropolitan area of a million

inhabitants, this would amount to about 130 additional early deaths per year, and in the country as a whole, these results indicate that a reduction of $10 \mu\text{g}/\text{m}^3$ would be expected to result in about 36,000 fewer early deaths per year. Although this study does not indicate the extent to which these deaths are brought forward, other studies of the harvesting issue (29–32) suggest that they are considerable.

The association of daily deaths with traffic particles also has no threshold and is somewhat steeper than the association with all $PM_{2.5}$. This is consistent with the results of Laden et al. (8), except that they used linear terms instead of smooth functions. This study confirms that this association extends to low levels. This result has considerable public policy relevance. Recently, automotive companies have proposed using diesel engines

to achieve higher fuel economy in the future. However, diesel engines produce substantially greater emissions of particles and particle precursors such as NO_x . The present results indicate that such an expansion of diesel engine use in the United States before diesel engines can meet the same particle emission levels as gasoline engines may result in important public health problems. A $1 \mu\text{g}/\text{m}^3$ increase in the concentration of traffic particles in the United States, for example, could be associated with about 7,000 additional early deaths per year in the United States.

REFERENCES AND NOTES

- Schwartz J, Dockery DW. Increased mortality in Philadelphia associated with daily air pollution concentrations. *Am Rev Respir Dis* 145:600–604 (1992).
- Pope CA, Dockery DW, Schwartz J. Review of epidemiologic evidence of health effects of particulate air pollution. *Inhal Toxicol* 7:1–18 (1995).
- Touloumi G, Pocock SJ, Katsouyanni K, Trichopoulos D. Short-term effects of air pollution on daily mortality in Athens: a time-series analysis. *Int J Epidemiol* 23:957–967 (1994).
- Katsouyanni K, Touloumi G, Spix C, Schwartz J, Balducci F, Medina S, Rossi G, Wojtyniak D, Sunyer J, Bacharova L, et al. Short term effects of ambient sulphur dioxide and particulate matter on mortality in 12 European cities: results from time series data from the APHEA project. *Br Med J* 314:1658–1663 (1997).
- Schwartz J, Dockery DW, Neas LM. Is daily mortality associated specifically with fine particles? *J Air Waste Manag Assoc* 46:2–14 (1996).
- Schwartz J. Assessing confounding, effect modification, and thresholds in the association between ambient particles and daily deaths. *Environ Health Perspect* 108:563–568 (2000).
- Samet JM, Zeger SL, Dominici F, Currier F, Coursac I, Dockery DW, Schwartz J, Zanobetti A. The National Morbidity, Mortality, and Air Pollution Study Part II: Morbidity, Mortality, and Air Pollution in the United States. Report 94. Boston, MA:Health Effects Institute, 2000.
- Laden F, Neas LM, Dockery DW, Schwartz J. Association of fine particulate matter from different sources with daily mortality in six U.S. cities. *Environ Health Perspect* 108(10):941–947 (2000).
- Daniels MJ, Dominici F, Samet JM, Zeger SL. Estimating particulate matter mortality dose response curves and threshold levels: an analysis of daily time series for the 20 largest U.S. cities. *Am J Epidemiol* 152:397–406 (2000).
- Schwartz J, Zanobetti A. Using meta-smoothing to estimate dose-response trends across multiple studies, with application to air pollution and daily death. *Epidemiology* 11(6):666–672 (2000).
- Schwartz J, Ballester F, Saez M, Pérez-Hoyos S, Bellido J, Cambra K, Arribas F, Cañada A, Pérez-Boillos M.J, Jordi Sunyer J. The concentration–response between air pollution and daily deaths. *Environ Health Perspect* 109:1001–1006 (2001).
- Ferris BG Jr, Speizer FE, Spengler JD, Dockery DW, Bishop YMM, Wolfson M, Humble C. Effects of sulfur oxides and respirable particulates on human health: methodology and demography of populations in study. *Am Rev Respir Dis* 120:767–779 (1979).
- Courtney WJ, Shaw RW, Dzuby TD. Precision and accuracy of a beta-gauge for aerosol mass determinations. *Environ Sci Technol* 16:236–239 (1982).
- Spengler JD, Thurston G. Mass and elemental composition of fine and coarse particles in six U.S. cities. *J Air Pollut Control Assoc* 33:1162–1171 (1983).
- Browne MW. On oblique Procrustes rotation. *Psychometrika* 32:125–132 (1967).
- Thurston GD, Spengler JD. A quantitative assessment of source contributions to inhalable particulate matter pollution in metropolitan Boston. *Atmos Environ* 19:9–25 (1985).
- U.S. National Climate Data Center. EarthInfo CD. NCDC Surface Airways. Boulder, CO:EarthInfo Inc.
- U.S. Department of Health and Human Services, Public




Figure 4. Cumulative normal curve versus a standardized predictor (the sum of the effects of all risk factors) over the range of exposures that correspond to daily death rates of between 1 and 3 per million, which is the observed range of variation in U.S. cities. It is quite linear in the predictor.




Figure 3. Overall estimated dose–response relation between $PM_{2.5}$ from traffic and daily deaths in six U.S. cities. The estimate is obtained by combining the estimated smoothed curve in each of the cities, after controlling for weather, season, and day of the week. Instead of linear terms for particles from other sources, in this analysis we controlled for smoothed terms for $PM_{2.5}$ from crustal sources, coal combustion, residual oil, and salt. The line shown is a least-squares regression line through the estimated points.

- Health Service, Centers for Disease Control and Prevention, National Center for Health Statistics. Public Use Data Tape Documentation. Hyattsville, MD:U.S. Department of Health and Human Services, 1993.
19. International Classification of Diseases, 9th revision, Vol 1. Commission on Professional and Hospital Activities. Ann Arbor, MI:Edward Brothel Inc, 1979.
 20. Hastie T, Tibshirani R. *Generalized Additive Models*. London:Chapman and Hall, 1990.
 21. Schwartz J. Generalized additive models in epidemiology. *Invited Papers: Proceedings of the 17th International Biometric Conference*. Hamilton, Ontario, Canada:International Biometric Society, 1994:55–80.
 22. Schwartz J. Air pollution and daily mortality: a review and meta analysis. *Environ Res* 64:36–52 (1994).
 23. Hoek G, Schwartz J, Groot B, Eilers P. Effects of ambient particulate matter and ozone on daily mortality in Rotterdam, The Netherlands. *J Arch Environ Health* 52:455–463 (1997).
 24. Michelozzi P, Forastiere F, Fusco D, Perucci CA, Ostro B, Ancona C, Pallotti G. Air pollution and daily mortality in Rome, Italy. *Occup Environ Med* 55(9):605–610 (1998).
 25. Cleveland WS, Devlin SJ. Robust locally-weighted regression and smoothing scatterplots. *J Am Stat Assoc* 74:829–836 (1988).
 26. Berkey CS, Hoaglin DC, Mosteller F, Colditz GA. A random-effects regression model for meta-analysis. *Stat Med* 14:395–411 (1995).
 27. U.S. EPA. National Ambient Air Quality Standards for Ozone and Particulate Matter, Proposed Decision. Fed Reg 62:7743–7744 (1997). Available: <http://www.epa.gov/fedrgstr/EPA-AIR/1997/February/Day-20/a4330.htm> [cited 31 July 2002].
 28. Schwartz J. Health effects of particulate air pollution: is there a threshold? In: *Relationship between Respiratory Disease and Exposure to Air Pollution* (Mohr U, ed). Washington, DC:ILSI Press, 1998:195–206.
 29. Zeger SL, Dominici F, Samet JM. Harvesting resistant estimates of air pollution effects on mortality. *Epidemiology* 10:171–175 (1999).
 30. Zanobetti A, Wand MP, Schwartz J, Ryan L. Generalized additive distributed lag models: quantifying mortality displacement. *Biostatistics* 1:279–292 (2000).
 31. Schwartz J. Is there harvesting in the association of airborne particles with daily deaths and hospital admissions? *Epidemiology* 12:55–61 (2001).
 32. Zanobetti A, Schwartz J, Samoli E, Gryparis A, Touloumi G, Atkinson R, Le Tertre A, Bobros J, Celko M, Goren A, et al. The temporal pattern of mortality responses to air pollution. *Epidemiology* 13(1):87–93 (2002).