



Risk and Exposure Assessment for Review of the Secondary National Ambient Air Quality Standards for Oxides of Nitrogen and Oxides of Sulfur

First Draft, Chapters 7-8

[This page intentionally left blank.]

EPA-452/P-08-005b
August 2008

**RISK AND EXPOSURE ASSESSMENT FOR REVIEW OF THE SECONDARY NATIONAL AMBIENT
AIR QUALITY STANDARDS FOR OXIDES OF NITROGEN AND OXIDES OF SULFUR**

FIRST DRAFT, CHAPTERS 7-8

U.S. Environmental Protection Agency
Office of Air Quality Planning and Standards
Research Triangle Park, NC

DISCLAIMER

This draft document has been prepared by staff from the Health and Environmental Impacts and Air Quality Analysis Divisions of the Office of Air Quality Planning and Standards, and the Clean Air Markets Division, Office of Air Programs, U.S. Environmental Protection Agency. Any opinions, findings, conclusions, or recommendations are those of the authors and do not necessarily reflect the views of EPA. This document is being circulated to obtain review and comment from the Clean Air Scientific Advisory Committee (CASAC) and the general public. Comments on this draft document should be addressed to Dr. Anne Rea, U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards, C539-02, Research Triangle Park, North Carolina 27711 (email: rea.anna@epa.gov).

7. SYNTHESIS AND INTEGRATION OF CASE STUDY RESULTS

The final version of this chapter will synthesize the results of the current conditions assessments for each of the targeted ecosystem effect areas (i.e., aquatic acidification, terrestrial acidification, aquatic nutrient enrichment, and terrestrial nutrient enrichment). Further, we will use these summaries of ecological effects and their associated ecological indicators to characterize the relationship between levels of an ecological indicator and the associated degree of ecologically adverse effects. This characterization of adversity can then be used to inform policy judgments on public welfare. Once analyses are completed, the next draft of this chapter will describe the range of effects associated with the current deposition of nitrogen and sulfur and demonstrate the links between adverse ecological effects and the level of associated ecological indicators. These relationships will then be used to inform alternative standards by translating adverse effects levels into levels of ecological indicators that can be linked through atmospheric deposition to ambient air concentrations of NO_x and SO_x.

Conducting a joint risk assessment for two criteria pollutants that impact multiple (and in some cases overlapping) ecological processes means that disparities exist in the level of information available for the targeted ecosystem effect areas highlighted in this review. This is reflected in the nature of the science, data, models, and time available for the case study analyses. However, a concise synthesis of the information from the case studies, including an assessment of the similarities and differences in the level of scientific data available to support the risk assessments for each effect area, will be useful to inform the standard setting process. The case study synthesis is intended to provide an integrated, weight-of-evidence approach that will consider the combined information from both evidence-based (i.e., relying primarily on reference to the ISA review of published literature on the effects of the deposition of NO_x and SO_x) and empirical approaches (based on results of applications of data, models, and information from the published literature), and their associated uncertainties, to explain how the risk and exposure assessment can inform decision makers about what levels of protection might be appropriate to protect public welfare from adverse impacts on ecosystems. This synthesis will look at the ecological indicators selected for each of the targeted effect area and examine the commonalities and differences among them. This information can then be used to inform a policy judgment on a known or anticipated adverse public welfare effect.

1 This chapter summarizes the current progress of the case studies (Section 7.1), begins to
2 discuss how the results of the case studies can be used to inform potential levels of protection
3 against ecosystem effects (Section 7.2), and preliminarily points to how this might be used in
4 designing a meaningful secondary standard or standards for ambient concentrations of NO_x and
5 SO_x (Section 7.3). In this draft, we focus on the ecological effects of aquatic acidification, using
6 surface water ANC as an ecological indicator. As our analyses progress, we will include
7 additional ecological indicators and explain how they may be used to characterize adverse
8 ecological effects.

9 **7.1 SUMMARY OF CASE STUDY ANALYSES**

10 **Table 7.1-1** summarizes the ecosystem characteristics, ecological indicators, ecological
11 effects, and ecosystem services evaluated in the case study analyses presented in Chapters 4 and
12 5. *Note: currently these results reflect the present status of the first draft risk and exposure*
13 *assessment and will be updated for the second draft as the analyses progress.*

14 **7.1.1 Aquatic Acidification**

15 Acidic deposition has altered major biogeochemical processes in the United States by
16 increasing the sulfur and nitrogen content of soils. This increase accelerates SO₄²⁻ and NO₃⁻
17 leaching from soil to drainage water, depletes base cations (namely Ca²⁺ and Mg²⁺) from soils,
18 and increases the mobility of toxic aluminum (Al) (see ISA Section 3.2) (U.S. EPA, 2008). Some
19 ecosystems and some areas of the United States are more sensitive than others to the effects of
20 acidic nitrogen and sulfur deposition, particularly streams and lakes with naturally low ANC.
21 The impact of acidic deposition on aquatic systems is controlled by several environmental
22 factors, such as geology, surface water flow, soil depth, and weathering rates, all of which
23 contribute to the ability of an ecosystem to neutralize the additional acidic deposition and prevent
24 the lowering of ANC. The Adirondack Mountains of New York and the Blue Ridge Mountains
25 in the Shenandoah National Park of Virginia were selected for the evaluation of impacts from
26 acidic deposition of nitrogen and sulfur. These regions receive relatively high levels of acidic
27 deposition and are known to be predisposed to acidification, and extensive research, data, and
28 models exist for these areas. In addition, the results will likely be relevant in characterizing risks
29 to larger areas of the Appalachian Mountains and the Northeast.

1 ANC is a key indicator of acidification with relevance to terrestrial and aquatic
2 ecosystems. ANC is useful because it integrates the overall acid-base status of a lake or stream
3 and reflects how aquatic ecosystems respond to acidic deposition over time. There is also a
4 relationship between ANC and the surface water constituents that directly contribute to or
5 ameliorate acidity-related stress, in particular, concentrations of hydrogen ion (as pH), Ca^{2+} , and
6 aluminum. Moreover, low pH surface waters leach aluminum from soils, which is quite lethal to
7 fish and other aquatic organisms. In aquatic systems, there is a direct relationship between ANC
8 and fish and phyto-zooplankton diversity and abundance (see Section 7.3). These relationships
9 also impact several ecosystem services, such as fishing, recreation, food, natural habitat, and
10 tourism.

11 **7.1.2 Terrestrial Acidification**

12 Terrestrial acidification occurs where inorganic and mineral acids are added to the soil.
13 This occurs as a result of both natural biogeochemical processes and acidic deposition of
14 nitrogen and sulfur. Biological effects of acidification on terrestrial ecosystems are generally
15 attributable to aluminum toxicity and decreased ability of plant roots to take up base cations.
16 Acidic deposition increases concentrations of sulfur and nitrogen in soil, which accelerates
17 leaching of SO_4^{2-} and NO_3^- from soil to drainage water. In the absence of anthropogenic sulfur
18 and nitrogen deposition, the limited mobility of anions in the soil controls the rate of base cation
19 leaching. Atmospheric deposition of sulfur and nitrogen provides anions that are more mobile in the
20 soil environment than naturally occurring anions in the soil; these mineral acid anions can accelerate
21 natural rates of base-cation leaching, particularly Ca^{2+} and Mg^{2+} , leading to mobilization of inorganic
22 aluminum, which is toxic to tree roots. Acidic deposition can also affect terrestrial ecosystems by
23 causing direct impacts on plant foliage.

24 The regions considered to be most sensitive to terrestrial acidification effects due to
25 acidic deposition are forested ecosystems of the Adirondack Mountains of New York, Green
26 Mountains of Vermont, White Mountains of New Hampshire, Allegheny Plateau of
27 Pennsylvania, and high-elevation forests in the southern Appalachians. The Kane Forest in the
28 Allegheny Plateau of Pennsylvania and the Hubbard Brook Experimental Forest in the White
29 Mountains of New Hampshire were selected to evaluate the effects of acidic deposition on the
30 sugar maple and red spruce due to known sensitivity of these species in these areas and the
31 availability of data. The ecosystem services affected include food, natural habitat, and tourism.

1 For our case study areas, we anticipate that sugar maple and red spruce abundance and growth
2 (i.e., crown vigor, biomass, and geographic extent) will be quantitatively linked to acidification
3 symptoms (specifically the calcium to aluminum ratio in the soil) through U.S. Forest Service
4 Forest Inventory and Analysis (FIA) database analyses and the analysis of estimated sales of
5 maple sugar products for the second draft risk and exposure assessment.¹

6 **7.1.3 Aquatic Nutrient Enrichment**

7 Nitrogen is an essential nutrient for estuarine and marine ecosystem fertility and is often
8 the limiting nutrient in aquatic ecosystems. Excessive nitrogen contributions can cause habitat
9 degradation, algal blooms, toxicity, hypoxia (i.e., reduced dissolved oxygen), anoxia (i.e.,
10 absence of dissolved oxygen), fish kills, and decreases in biodiversity. A recent national
11 assessment of estuary trophic conditions used five biological indicators to assess nutrient
12 enrichment: chlorophyll *a*, macroalgae, dissolved oxygen, nuisance/toxic algal blooms, and
13 SAV. This assessment is discussed in detail in Attachment 5.

14 The Chesapeake Bay and Pamlico Sound were chosen as case study areas based on the
15 availability of atmospheric deposition data, water quality modeling data that accounted for
16 atmospheric deposition, a large main stem river, and scalability. Because the Chesapeake Bay
17 and Pamlico Sound are fed by multiple river systems, the case study areas were scaled to one
18 main stem river for each system: the Potomac River Basin for the Chesapeake Bay and the Neuse
19 River Basin for the Pamlico Sound. A eutrophication index, developed by National Oceanic and
20 Atmospheric Administration's (NOAA's) National Estuarine Eutrophication Assessment
21 (NEEA) was used in this analysis, which estimates the likelihood that the estuary is experiencing
22 eutrophication or will experience eutrophication in the future. The endpoints of interest are fish
23 populations; water quality; and habitat quality and their related ecosystem services, including
24 fisheries (e.g., closings, decreased species richness), recreation, and tourism.

25 **7.1.4 Terrestrial Nutrient Enrichment**

26 Terrestrial ecosystems typically respond to total reactive nitrogen loading and, with few
27 exceptions, have limited differential responses to the specific chemical species of deposited
28 nitrogen. Nitrogen compounds that are contained in atmospheric deposition, and that influence

¹ The FIA can be found at <http://fia.fs.fed.us/>.

1 nutrient dynamics, include multiple oxidized and reduced forms of inorganic nitrogen. Excess
2 nitrogen in ecosystems causes inadvertent fertilization of trees and grasslands, creating unnatural
3 growth rates in some species and changing competitive interactions among species, and nutrient
4 imbalances. These impacts ultimately decrease ecosystem health and biodiversity. Forest growth
5 enhancement can potentially exacerbate other nutrient deficiencies, such as Ca, Mg, or K,
6 thereby causing problems with forest health. Enhanced growth generally occurs above ground
7 level. This can cause changes in the shoot-to-root ratio, causing decreased resistance to
8 environmental stressors, such as drought. In conifer forests, multiple long-term experiments (see
9 Attachment 6) have demonstrated transient growth increases (generally at deposition rates lower
10 than 10 kg N/ha/yr) followed by increased mortality, especially at higher rates of fertilization. In
11 the western United States, atmospheric nitrogen deposition has been shown to cause increased
12 litter accumulation and carbon storage in aboveground woody biomass, which, in turn, may lead
13 to increased susceptibility to more severe fires. Grassland communities that are adapted to low
14 nutrient supply can exhibit substantial sensitivity to nutrient enrichment effects of nitrogen
15 deposition.

16 Two of the primary indicators of nitrogen enrichment in forested watersheds are the
17 leaching of NO_3^- in soil drainage waters and the export of NO_3^- in stream water, especially during
18 the growing season. Low C:N ratios in soils are also commonly related to increased nitrification,
19 potential increases in soil acidity, and releases of nitrate to receiving waters; however, these
20 measurements are not always widely available.

21 California's CSS and mixed conifer forest communities were selected for an initial case
22 study analysis. These areas were selected based on the availability of atmospheric ambient and
23 deposition data; the availability of digitized datasets of biotic communities; fire-prone areas;
24 sensitive, rare species; their representativeness of western United States ecosystems potentially
25 impacted by nitrogen deposition; and their scalability and generalization potential.

26 California's CSS and mixed conifer forests on the slopes of the San Bernardino and
27 Sierra Nevada Mountains have important recreational value, protect water resources, and provide
28 habitats for many other species. In the mixed conifer forest, lichen communities and nitrogen
29 saturation can provide a means to identify the effects of nitrogen loadings. Both habitats provide
30 a number of ecological services, including biodiversity, water quality, recreation, timber, and fire
31 hazard mitigation. Linkage methods from endpoint to services may include measurement of

1 changes in biodiversity and abundance and distribution of threatened and endangered species,
2 comparison of past and present photography, and measurement of the distribution of soil
3 moisture with depth and possible nitrate leaching.

4 **7.1.5 Sulfur and Mercury Methylation**

5 The biogeochemical cycle of mercury, a well-known neurotoxin, is closely tied to the
6 sulfur cycle. Mercury is taken up by living organisms in the methylated form, which is easily
7 bioaccumulated in the food web. Sulfate-reducing bacteria in wetland and lake sediments play a
8 key role in mercury methylation. Changes in sulfate² deposition have resulted in changes in both
9 the rate of mercury methylation and the corresponding mercury concentrations in fish.

10 Although sulfur deposition is important to mercury methylation, several other interrelated
11 factors seem to also be related to mercury uptake, including low lake water pH, dissolved
12 organic carbon, suspended particulate matter concentrations in the water column, temperature,
13 and dissolved oxygen. In addition, the proportion of upland to wetland land area within a
14 watershed, as well as wetland type and annual water yield, appear to be important.

15 The major ecosystem services potentially impacted by sulfur-mediated mercury
16 methylation are provisioning and cultural services. Fishing and shellfishing can involve both
17 commercial operations and sports fishing, which provide food for human populations. For some
18 socioeconomic groups (especially groups with low incomes), fishing is a subsistence activity that
19 makes a very significant contribution to household food intake. Sport fishing often involves
20 important recreational services, and for many groups (e.g., Native Americans, Alaska native
21 villagers), fishing and consuming local fish or shellfish is of cultural and spiritual significance.

² EPA notes that in its review of the primary NAAQS for sulfur oxides, it has made the decision to examine all sulfate health effects as part of the review of the PM NAAQS (sulfates being one of the chief types of secondary PM). EPA believes it appropriate; however, to consider secondary sulfate effects as part of this review, rather than in the PM NAAQS review, because the effect at issue is related specifically to chemical composition. The fact that sulfate is deposited as a particle is not relevant to its contribution to mercury methylation.

Table 7.1-1. Summary of Characteristics, Ecological Indicators, Effects, and Ecosystem Services Impacted for Each Case Study Evaluated in This Review

| Targeted Effect Area | Characteristics of Sensitivity (Variable Ecological Factors) | Biological/Chemical Characteristic | Ecological Indicator | Ecological Effects | Ecosystem Services Impacted | Case Study Locations | Deposition Levels (NO _x and/or SO _x) Associated with Observed Effects (from ISA) |
|------------------------------------|----------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------|------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|
| Aquatic Acidification | Geology, surface water flow, soil depth, weathering rates | [Al] pH ANC | Species richness, abundance, composition ANC | Species losses of fish, phytoplankton, zooplankton; changed community composition, ecosystem structure, and function | Fisheries, recreation, tourism | Adirondack Mountains, NY Blue Ridge Mountains, Shenandoah National Park, VA | 15 kg SO ₄ ⁻² /ha yr 10 kg NO ₃ /ha yr |
| Terrestrial Acidification | Geology, surface water flow, soil depth, weathering rates | Soil base saturation [Al] [Ca] C:N ratio Ca:Al ratio | Tree health Red spruce, sugar maple ANC Ca:Al ratio | Decreased tree growth Increased susceptibility to stress, episodic dieback; changed community composition, ecosystem structure, and function | Food, natural habitat, tourism | Kane Forest (Allegheny Plateau, PA) Hubbard Brook Experimental Forest (White Mountains, NH) | |
| Aquatic Nutrient Enrichment | N-limited systems, presence of nitrogen in surface water, eutrophication status, nutrient criteria | Chlorophyll <i>a</i> , macroalgae, dissolved oxygen, nuisance/toxic algal blooms, submerged aquatic vegetation (SAV) | Eutrophication Index (EI) | Habitat degradation, algal blooms, toxicity, hypoxia, anoxia, fish kills, decreases in biodiversity | Fish populations, water quality, habitat quality | Potomac River Basin, Chesapeake Bay Neuse River Basin, Pamlico Sound | |

| Targeted Effect Area | Characteristics of Sensitivity (Variable Ecological Factors) | Biological/Chemical Characteristic | Ecological Indicator | Ecological Effects | Ecosystem Services Impacted | Case Study Locations | Deposition Levels (NO _x and/or SO _x) Associated with Observed Effects (from ISA) |
|-------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------|---------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------|----------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|
| Terrestrial Nutrient Enrichment | Presence of acidophytic lichens, anthropogenic land cover | Cation Exchange Capacity, C:N ratios, Ca:Al ratios, NO ₃ ⁻ leaching and export | Species composition | Species changes, nitrogen enrichment of soil, changes in fire regime, changes in nutrient cycling | Loss of habitat, loss of biodiversity, recreation, water quality | Coastal sage scrub and mixed conifer forest (San Bernardino and Sierra Nevada Mountain Ranges, CA) | 10–15 kg N/ha yr 3–8 kg/ha yr for lichen effects |
| Sulfur and Mercury Methylation Potential | Wetland type, presence of sulfate-reducing bacteria, water pH, dissolved organic carbon, suspended particulate matter | Interaction among dissolved organic carbon, temperature, anoxia, and sulfide Land cover, precipitation response, and limnography | MeHg concentrations in fish and shellfish | Neurotoxic effects in fish and throughout food web | Fishing, shellfishing, sports fishing, food, recreation, biodiversity | Little Rock Lake, WI (ISA case study) | |

1 **7.2 INTEGRATING CASE STUDY RESULTS**

2 The ISA and case study analyses have shown that, in certain ecosystems, welfare effects
3 are occurring under current ambient concentrations of NO_x and SO_x, even though nationwide
4 (with only a few exceptions) we are in attainment of the current primary and secondary NO₂ and
5 SO₂ NAAQS. The next step in the risk and exposure assessment process is to determine what
6 alternative levels of ambient NO_x and SO_x might allow (1) various degrees of recovery of
7 impacted systems, (2) acceleration of recovery of impacted systems, and (3) protection against
8 further damage, recognizing the variability in ecosystem sensitivities to nitrogen and sulfur
9 deposition. At this point in the risk and exposure assessment, we have begun to address many of
10 the policy-relevant questions guiding this review. Our initial case study analyses have begun to
11 show the variability associated with these responses and the degree to which ecological
12 indicators are suited to address a particular type of effect (i.e., acidification or nutrient
13 enrichment in terrestrial or aquatic ecosystems).

14 There are many factors that determine whether or not an ecological effect occurs in
15 response to ambient concentrations of NO_x and SO_x. These “variable factors” may refer to either
16 ecological factors or atmospheric factors, both of which influence deposition or exposure and the
17 associated ecological effects (i.e., acidification or nutrient enrichment). The degree to which any
18 given level of atmospheric deposition of nitrogen and sulfur impacts ecosystems depends in large
19 part on a suite of variable ecological factors at the point of deposition, including geology, surface
20 water flow, soil depth, and weathering rates. The levels of these ecological factors in a given
21 location determine the ecological conditions governing sensitivity to atmospheric deposition of
22 nitrogen and sulfur. The linkages between a given ecological indicator and deposition, as
23 influenced by the variable ecological factors, are discussed in Chapter 8.

24 In the next draft of the risk and exposure assessment, we will look across the results of
25 the case studies to identify ecological indicators that provide the most comprehensive coverage
26 across sensitive ecosystems. These indicators will form the basis for assessing links between
27 adverse effects and potential levels of an ecological indicator to protect against those effects. At
28 this point, the Aquatic Acidification Case Study analysis is the furthest along and will be used to
29 illustrate how an ecological indicator of acidification (i.e., ANC) and the levels of ecological

1 effects associated with changes in that indicator can be used to characterize adversity to public
2 welfare.

3 **7.3 LINKAGES BETWEEN ECOLOGICAL INDICATORS AND ADVERSE** 4 **EFFECTS**

5 Within the limitations of currently available data and models, our case study analyses
6 indicate the degree, if any, to which adverse ecological effects are already occurring in specific
7 sensitive ecosystems at current levels of ambient nitrogen and sulfur deposition. Note that
8 ecological recovery may not immediately result from decreases in atmospheric deposition, but
9 may lag behind changes in atmospheric deposition. Since current ecological conditions reflect
10 past deposition and responses to it, it is possible that an ecological indicator such as ANC may
11 improve over time (years) under current deposition levels. However, the recovery rate may be
12 affected by current levels of deposition, with lower levels of deposition leading to a more rapid
13 recovery compared to current levels.


14 **7.3.1 ANC: The Critical Chemical Parameter**

15 Ambient air concentrations are linked to adverse ecological effects via the process of
16 atmospheric deposition. For our purposes, it is necessary to identify a suitably generalized
17 ecological indicator for these effects because linking individual ecological endpoints directly to
18 atmospheric deposition is difficult. The relationship between ecological effects associated with
19 nitrogen and sulfur deposition and a commonly measured and well understood ecological
20 indicator, ANC, is described below. In addition, we conceptually describe linking ecological
21 effects with potential measures of adversity to public welfare by characterizing impacts on
22 ecosystem services. This information can then help to inform judgments about the extent to
23 which different levels of a particular ecological indicator might affect public welfare.

24 The susceptibility of a catchment to terrestrial or aquatic acidification by nitrogen and
25 sulfur deposition is controlled by the ability of the watershed to neutralize the addition of
26 nonmarine acid anion concentrations of SO_4^{2-} or NO_3^- to the catchment. The neutralizing ability
27 of a catchment is a function of the acid-base balance; the balance between the sum of base
28 cations (i.e., $\text{Ca}^* + \text{Mg}^* + \text{K}^* + \text{Na}^*$) inputs minus the sum of acid anions (i.e., SO_4^{2-} or NO_3^-)
29 inputs. In general, catchments with positive acid-base balance have base cations inputs that

1 exceed acid anions inputs which “buffer” the watershed from becoming acidic. The ability of a
2 catchment to buffer acid anion deposition is known as the “buffering capacity,” which is
3 controlled by a host of environmental factors within the watershed. Such factors include base
4 cation leaching rate, base cation weathering rates, base saturation, uptake by vegetation, rate of
5 surface water flow, soil depth, bedrock, etc. The buffering capacity of a catchment is controlled
6 by the flux of base cations, which act to balance acid anion inputs. The effect of acid anion
7 deposition is to lower this acid-base balance, causing a decline in the ecological system’s
8 composition, structure, and the biological fitness of its organisms over time.

9 The ANC of surface waters is widely used as a chemical indicator of acidic conditions
10 because it has been found in many studies to be the best single indicator of the biological
11 response and health of aquatic communities in acid-sensitive systems (Lien et al., 1992; Sullivan
12 et al., 2006). Logistic regression of species presence/absence data against ANC provides a
13 quantitative dose-response function, which indicates the probability of occurrence of an
14 organism for a given value of ANC. For example, the number of fish species present in a
15 waterbody has been shown to be positively correlated with the ANC level in the water, with
16 higher values supporting a greater richness and diversity of fish species (**Figure 7.3-1**). The
17 diversity and distribution of phyto-zooplankton communities are also positively correlated with
18 ANC.



1
2 **Figure 7.3-1.** Number of fish species per lake or stream versus acidity status, expressed
3 as ANC in lakes in the Adirondacks of New York (Sullivan et al., 2006). Five classes: <0,
4 0–20, 20–50, 50–100 µeq/L, and >100 µeq/L describe the biological effects at a range of
5 ANC concentrations. See Table 7.3-1 for more details.

6 For freshwater systems, ANC levels can be grouped into five major classes: <0, 0–20,
7 20–50, 50–100 µeq/L, and >100 µeq/L with each range representing a probability of ecological
8 damage to the community (see Attachment 3). ANC values above 100 µeq/L are generally not
9 harmful (Figure 7.3-1) to biota. Below 100 µeq/L, it has been shown that fish fitness and
10 community diversity begin to decline, but the overall health of the community remains high as
11 long as ANC concentrations do not go below 50 µeq/L. ANC concentrations below 50 µeq/L
12 result in negative effects on sensitive biota. From 50 to 20 µeq/L, it has been shown that fish
13 diversity and the overall fitness (i.e., health and reproduction) of most aquatic organisms in the
14 waterbody are reduced. Below 20 µeq/L, all biota exhibit some level of negative effects,
15 particularly because surface waters at this level are susceptible to episodic acidification and their
16 associated harmful effects. Fish and plankton diversity and the structure of the communities
17 continue to decline sharply to levels where acidophilic species begin to outnumber all other

1 species. Below an ANC of 0 $\mu\text{eq/L}$, complete loss of fish populations and extremely low
 2 diversity of planktonic communities occur. At these low levels, only acidophilic species are
 3 present, but even their population and community structure are sharply reduced. The five
 4 categories of ANC and expected ecological effects are described in **Table 7.3-1** and are
 5 supported by a large body of research completed throughout the eastern United States (Sullivan
 6 et al., 2006). The connection between the ecological effects described in Table 7.3-1 and
 7 ecosystem services that can be indicative of adverse impacts to public welfare will be explored in
 8 the second draft risk and exposure assessment. However, examples of these ecosystem services
 9 include recreational and subsistence fishing, recreational lake use, natural habitat provision, and
 10 tourism. Potential values of these ecosystem services related to alternative levels of ANC will be
 11 assessed in the second draft risk and exposure assessment.

Table 7.3-1. Aquatic Status Categories

| Category Label | ANC Levels | Expected Ecological Effects |
|------------------|----------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Acute Concern | <0 micro equivalent per Liter ($\mu\text{eq/L}$) | Complete loss of fish populations is expected. Planktonic communities have extremely low diversity and are dominated by acidophilic forms. The numbers of individuals in plankton species that are present are greatly reduced. |
| Severe Concern | 0–20 $\mu\text{eq/L}$ | Highly sensitive to episodic acidification. During episodes of high acidic deposition, brook trout populations may experience lethal effects. Diversity and distribution of zooplankton communities decline sharply. |
| Elevated Concern | 20–50 $\mu\text{eq/L}$ | Fish species richness is greatly reduced (i.e., more than half of expected species are missing). On average, brook trout populations experience sub-lethal effects, including loss of health and reproduction (fitness). Diversity and distribution of zooplankton communities decline. |
| Moderate Concern | 50–100 $\mu\text{eq/L}$ | Fish species richness begins to decline (i.e., sensitive species are lost from lakes). Brook trout populations are sensitive and variable, with possible sub-lethal effects. Diversity and distribution of zooplankton communities begin to decline as species that are sensitive to acidic deposition are affected. |
| Low Concern | >100 $\mu\text{eq/L}$ | Fish species richness may be unaffected. Reproducing brook trout populations are expected where habitat is suitable. Zooplankton communities are unaffected and exhibit expected diversity and distribution. |

12 Whether the biological community of a waterbody is at risk from current deposition loads
 13 depends on the ability of the watershed to neutralize the incoming acid anion deposition and

1 maintain an acid-base balance or ANC level that protects the biological community. This linkage
2 between deposition and biological processes is derived from a series of empirical relationships,
3 that relate the sustainable buffering capacity (preindustrial) of a catchment to the maximum flux
4 of acidic deposition which will maintain a particular level of ANC concentration in the surface
5 water. ANC is operationally defined as the sum of base cations minus the sum of acid anions,
6 with the proportion of ions deriving from neutral sea-spray removed:

$$7 \qquad \qquad \qquad \text{ANC} = [\text{BC}]_t^* - [\text{AA}]_t^* \qquad (1)$$

8 where $[\text{BC}]_t^*$ is the current, measured sum of nonmarine base cation equivalent concentrations
9 (i.e., $\text{Ca}^* + \text{Mg}^* + \text{K}^* + \text{Na}^*$) and $[\text{AA}]_t^*$ is the sum of nonmarine acid anion equivalent
10 concentrations (i.e., $\text{SO}_4^* + \text{NO}_3^*$). It is assumed that all chloride is derived from marine
11 sources. The * denotes the nonmarine component, whereby the marine contribution of each ion is
12 subtracted as a proportion of measured chloride concentration from the known ratio of these ions
13 in seawater. The equation defining ANC forms the basis of the linkage between deposition and
14 surface water ANC concentrations. This linkage is explored in greater detail in Chapter 8.

1 **7.4 REFERENCES**

- 2 Lien, L., G.G. Raddum, and A. Fjellheim. 1992. *Critical Loads of Acidity to Freshwater: Fish*
3 *and Invertebrates*. The Environmental Tolerance Levels Programme. Rep. No. 23/1992.
4 Norwegian Ministry of Environment, Oslo, Norway.
- 5 Sullivan, T.J., C.T. Driscoll, B.J. Cosby, I.J. Fernandez, A.T. Herlihy, J. Zhai, R. Stemberger,
6 K.U. Snyder, J.W. Sutherland, S.A. Nierzwicki-Bauer, C.W. Boylen, T.C. McDonnell,
7 and N.A. Nowicki. 2006. *Assessment of the Extent to Which Intensively-Studied Lakes*
8 *are Representative of the Adirondack Mountain Region*. Final report. Albany, NY: New
9 York State Energy Research and Development Authority (NYSERDA). Available at
10 <http://nysl.nysed.gov/uhtbin/cgisirsi/Qc wd6NzFby/NYSL/138650099/8/4298474>
11 (accessed November 1, 2007).
- 12 U.S. EPA (Environmental Protection Agency). 2008. *Integrated Science Assessment for Oxides*
13 *of Nitrogen and Sulfur – Environmental Criteria (Second External Review Draft)*.
14 Prepared by U.S. Environmental Protection Agency, National Center for Environmental
15 Assessment-RTP Division, Office of Research and Development, Research Triangle
16 Park, NC. EPA/600/R-08-082. August.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

[This page intentionally left blank.]

8. CONSIDERATIONS IN THE STRUCTURE OF NO_x/SO_x SECONDARY STANDARD

The previous chapters have provided an understanding of the risks associated with current deposition of nitrogen and sulfur associated with ambient atmospheric concentrations of NO_x and SO_x. Given that information, the next step is to provide a framework for developing standards for NO_x and SO_x that provide the appropriate level of protection against those risks. The purposes of this chapter are 1) to propose one possible framework for a NO_x and SO_x secondary standard(s), based on the linkages that exist between ambient atmospheric concentrations of NO_x and SO_x, atmospheric deposition of N and S, ecological effects and the related ecological indicators and 2) to illustrate how such a standard might be expressed for aquatic acidity using data from lakes in the Adirondacks case study area. The framework is designed to both meet statutory and legal requirements, and to reflect the complex nature of atmospheric and ecological systems.

In this chapter, we discuss constructing a standard based on achieving a uniform level of ecosystem protection. We explore a framework by which ambient atmospheric concentrations of NO_x and SO_x can be translated into a measure of ecosystem effects, using the transformation functions described below. To facilitate a more concrete understanding of this framework, we illustrate how the atmospheric concentration standards can be calculated using the Adirondacks case study example.

The remainder of this chapter is organized as follows. In Section 8.1, we discuss a possible structure for a secondary NAAQS for NO_x and SO_x, building on the conceptual model introduced in Chapter 1. In section 8.2, we present a discussion of the statutory and legal considerations that affect the design of an ecologically relevant ambient air quality standard for NO_x and SO_x. Section 8.3 discusses the framework for structuring a standard which links atmospheric concentrations of NO_x and SO_x to a specific ecological indicator of effects (i.e., ANC) through the deposition process, taking atmospheric and ecological factors into account. It also builds upon the current conditions risk and exposure assessment presented in Chapters 3 through 6 as well as the linkages presented in Chapter 7 to discuss the nature of a meaningful standard in light of the wide range of ecological effects resulting from deposition of NO_x and SO_x. In so doing, Section 8.3 expands on the standards framework through which the linkages

1 between ambient atmospheric concentrations of NO_x and SO_x, deposition of N and S, and
2 ecological indicators can be described. Section 8.4, which is being developed for inclusion in the
3 next draft, will describe an illustrative example of how ambient atmospheric concentrations for
4 NO_x and SO_x can be compared to a desired level of ecological protection, as expressed using the
5 ecological indicator ANC. This example will also show how observed (or modeled) ambient
6 atmospheric concentrations of NO_x and SO_x can be transformed into ANC values for
7 comparison against the level of a standard defined in terms of a value of the ecological indicator.
8 This example will also provide step-by-step calculations to show how different combinations of
9 atmospheric concentrations of NO_x and SO_x can achieve the same level of ANC, based on the
10 deposition transformation functions and ecological effect functions that are described in Section
11 8.3. In each case, the example calculations will show clearly how the standards framework can
12 incorporate 1) the influence of atmospheric factors that affect deposition, 2) the influence of
13 ecological factors that affect ANC, and 3) the influence of reduced forms of nitrogen (ammonia
14 and ammonium).

15

16 **8.1 POSSIBLE STRUCTURE OF A SECONDARY NAAQS**

17 Throughout this assessment, we have recognized there are important linkages between
18 ambient nitrogen and sulfur and their depositional effects on the environment. Our approach to
19 this joint, multi-pollutant review of two secondary standards has been to evaluate the science
20 driving welfare effects and consider what is needed to develop a secondary standard or suite of
21 standards that appropriately protects public welfare. The structure of the current secondary
22 standards for NO_x and SO_x is based on the ambient concentration of a single indicator that is
23 specified as a nitrogen- or sulfur-containing compound (i.e., NO₂ and SO₂, respectively) where
24 the level of the indicator does not necessarily correlate well with the ecological effects of
25 concern (USEPA, 2008). As discussed in Chapter 7, we recognize that the scientific basis that
26 forms our understanding of the relationship between the criteria air pollutants, NO_x and SO_x and
27 their associated public welfare effects supports the use of a more complex structure for the
28 secondary standard(s). The inherently complex and variable linkages between ambient
29 concentrations of NO_x and SO_x, their deposited forms of nitrogen and sulfur, and the ecological

1 responses that are associated with public welfare effects call for a more complex standard design
2 that reflects these linkages.

3 A critical difference between the current risk and exposure assessment and those from
4 previous reviews is that in the current assessment, exposures and ecological effects are related to
5 current levels of nitrogen and sulfur deposition in selected case study locations, rather than just
6 the ambient concentrations of NO_x and SO_x in those locations. Based on the recognized linkages
7 between ambient concentrations of NO_x and SO_x and deposition of nitrogen and sulfur (see
8 Chapter 3), as well as the linkages between deposition and ecosystem effects, a specified level of
9 ecosystem effect will require limits on ambient concentrations of NO_x and SO_x, all other factors
10 being held constant. A standard that is expressed in terms of atmospheric concentrations of NO_x
11 and SO_x will require a process to translate ambient concentrations through the deposition process
12 into ecological effects, to determine if the specified level of ecosystem effect is achieved. This
13 atmospheric concentration-based standard will explicitly identify the role of modifying factors in
14 affecting the levels of NO_x and SO_x concentrations that would be necessary to achieve the
15 standard.

16 A possible structure for such standards is shown below in Figure 8.1-1, which builds
17 upon the conceptual model presented in Chapter 1. Figure 8.1-1 represents the pathway from
18 atmospheric concentrations of NO_x and SO_x to an ecological effect (aquatic acidification) as
19 represented by an ecological indicator (ANC), through the processes of deposition, with each of
20 the steps along the pathway expanded to reflect the example of aquatic acidification using the
21 ecological indicator ANC. Suites of variable factors reflecting both ecological and atmospheric
22 complexity modify the pathways. Critical to developing an ecologically appropriate secondary
23 standard for NO_x and SO_x is understanding that the different levels of the ecological indicator
24 can be associated with different levels of protection against adverse public welfare effects. The
25 ambient standard is then specified as the level(s) of ambient NO_x and SO_x such that the
26 calculated ecological indicator meets the level of protection of public welfare set by the
27 Administrator. Thus, the standard is expressed in terms of the relationship between ambient NO_x
28 and SO_x and the ecological indicator, linked through the deposition of N and S.




Figure 8.1-1. Possible structure of a secondary NAAQS for NO_x and SO_x based on the ecological indicator ANC for the effect of acidification.

The variable atmospheric factors which influence atmospheric conditions affecting the transformation, transport and deposition of ambient concentrations of NO_x and SO_x include relative humidity, precipitation, wind conditions, and chemical reactivity. Deposition is also affected by seasonally dependent landscape factors such as leaf surface area. Since deposition is the driver for many of the associated ecological effects due to NO_x and SO_x , such understanding of the factors that link atmospheric concentrations with deposition is critical. In addition, the relative contribution of NO_x to total deposition of nitrogen is affected by the ambient concentrations of ammonia and ammonium. The linkages between atmospheric concentrations and deposition as influenced by the variable atmospheric and landscape factors are discussed in Section 8.3.1.

1 The degree to which any given level of deposition of nitrogen and sulfur containing species
2 impacts ecosystems depends in large part on a suite of variable ecological factors at the point of
3 deposition, including soil base saturation, soil thickness, runoff rate, land use conditions, nutrient
4 cycling, and non-atmospheric loadings of sulfur and nitrogen. The particular levels of these
5 factors that occur at any location form the ecological baseline against which ecological effects
6 from deposition are evaluated. The linkages between deposition and ecological effect as
7 influenced by the variable ecological factors are discussed in Section 8.3.2.

8 The relationships illustrated in Figure 8.1-1 in concept can be operated in either direction,
9 from known or assumed ambient atmospheric concentrations to the resulting ecological effects,
10 or from desired ecological conditions in an area of interest to ambient concentrations that would
11 achieve them. In either approach, there may not be a unique combination of ambient NO_x and
12 SO_x that achieves a specified level of ecological protection, because of the multiple
13 combinations of NO_x and SO_x concentrations that can yield the same level of ecological
14 indicator. This is particularly true for the case of aquatic acidity, but may not hold true for other
15 ecological endpoints. Whether precipitation occurs or not also greatly affects the relationship
16 between ambient atmospheric concentrations of NO_x and SO_x and deposition of N and S, so
17 knowledge or an assumption of precipitation is needed to complete the quantitative trail from
18 ambient concentrations to ecological effects, or the reverse. Finally, concentrations and
19 deposition of reduced nitrogen species influence ecological conditions, but these species are not
20 currently encompassed by the listed criteria pollutant NO_x. While reduced species of nitrogen do
21 contribute to the overall loadings of N, as will be demonstrated in a future version of Section 8.4,
22 it may be possible to focus solely on NO_x contributions to loadings while recognizing that there
23 are impacts from reduced forms of nitrogen that must be taken into account.

24 **8.2 STATUTORY AND LEGAL CONSIDERATIONS IN THE** 25 **STRUCTURE OF THE STANDARD**

26 Inherent in the approach described above are various legal considerations as to the
27 requirements of the CAA and EPA's ability to implement various types of standards under the
28 Act. This section will begin to discuss the legal framework for considering appropriate air
29 quality indicator(s), averaging time(s), form(s), and level(s) of standard(s) to protect against
30 known or anticipated adverse impacts on public welfare. As noted above, the structure of such a

1 standard would likely be complex, reflecting the variable factors defining the atmospheric and
2 environmental conditions that affect deposition of NO_x and SO_x, and those variable ecological
3 factors that affect ecosystem responses.

4 In setting NAAQS to protect public health and welfare, EPA has historically established
5 standards which require the comparison of locally monitored ambient concentrations of an air
6 pollutant indicator against a specified numerical level of atmospheric concentration of a
7 pollutant, using a specified averaging time and statistical form. For example, the current
8 secondary standard for NO_x uses ambient concentrations of NO₂ as the indicator, and attainment
9 is determined by comparing the annual arithmetic mean of measured ambient concentrations, for
10 a calendar year, against the level of 0.053 ppm. As more is learned about the effects of pollutants
11 such as NO_x and SO_x on the environment, such a relatively simple framework may not be the
12 most appropriate way to protect sensitive ecosystems from effects associated with ambient
13 concentrations of the air pollutants. EPA is considering in this review of the secondary standard
14 for NO_x and SO_x a conceptual framework for evaluating the environmental effect of ambient
15 concentrations of NO_x and SO_x that takes into account variable factors, such as atmospheric and
16 ecosystem conditions that modify the amounts and effects of deposited N and S on ecosystems.
17 EPA is also considering whether the secondary standard can and should be structured to better
18 take this framework into account, as a way to appropriately protect the public welfare from the
19 effects associated with the presence of these pollutants in the ambient air.

20 As discussed previously, this could involve a standard with an ecological indicator which
21 can be linked to ambient concentrations of SO_x and NO_x. The standard could include a nationally
22 uniform level for a particular ecological indicator, for example a specified level of ANC that
23 would be determined by the Administrator to represent the appropriate level of protection. This
24 framework would also take into account other variables, both atmospheric and ecological that
25 may modify the ecological effects. Compliance would be determined by measuring NO_x and SO_x
26 concentrations, using these measurements and the various transformation functions to calculate
27 an ANC level, and comparing the calculated level of ANC to the level of ANC set by the
28 Administrator, including averaging time and form. Since the effect of a certain ambient
29 concentration of SO_x and NO_x on ANC could vary depending on the other variable factors,
30 whether a certain ambient concentration of SO_x and NO_x would attain the standard would depend
31 in part on these other variables, which could vary geographically and temporally.

1 One important issue in evaluating such an approach is EPA’s authority to structure a
2 NAAQS in such a way. EPA’s review of a NAAQS starts with its review of the air quality
3 criteria for the pollutant. Under Section 109(d)(1) of the CAA, EPA must undertake a thorough
4 review of the air quality criteria for the pollutant at issue as part of reviewing a secondary
5 NAAQS to determine whether a current standard is requisite to protect the public welfare. Under
6 Section 108, air quality criteria are to “reflect the latest scientific knowledge useful in indicating
7 the kind and extent of all identifiable effects” associated with the presence of the pollutant in the
8 ambient air. It is clear from the language of the CAA that where the state of the science provides
9 a basis for considering such effects as relevant, the review of air quality criteria should
10 encompass a broad analysis of “any” anticipated adverse effects, as well as the ways in which
11 variable conditions such as atmospheric conditions or ecological conditions may influence the
12 effect of a pollutant and the ways in which other air pollutants may interact with the criteria
13 pollutant to produce adverse effects. Specifically, Section 108(a)(2) of the CAA provides that:

14 The criteria for an air pollutant to the extent practicable shall include information on the
15 following:

- 16 1. Those variable factors (including atmospheric conditions) which of themselves or in
17 combination with other factors may alter the effects on public health or welfare of
18 such air pollutants;
 - 19 2. The types of air pollutants which, when present in the atmosphere, may interact with
20 such pollutants to produce an adverse effect on public health or welfare; and
 - 21 3. Any known or anticipated adverse effects on welfare.
- 22

23 Taking into account this extensive review of the air quality criteria for an air pollutant,
24 the Administrator is required to review and to revise, as appropriate, the secondary standard. The
25 secondary standard is to “specify a level of air quality which, in the judgment of the
26 Administrator, *based on such criteria*, is requisite to protect public welfare from any known or
27 anticipated adverse effects associated with the presence of such air pollutant in the ambient air.”
28 CAA § 109(b) (emphasis added). “Welfare,” in turn, is defined to include a broad array of
29 effects, including effects on soil, water, crops, vegetation, and manmade materials, “whether
30 caused by transformation, conversion, or combination with other air pollutants.” CAA § 302(h).
31 As with the description of the air quality criteria in Section 108, the CAA uses expansive
32 language in describing the range of effects that EPA should take into account in setting a
33 standard that is requisite to protect public welfare.

1 Thus, Section 108 and the definition of welfare require that EPA analyze and evaluate
2 scientific information on the interaction between ambient concentrations of a pollutant and other
3 variable factors, atmospheric and ecological, that affect the air pollutants impact on the
4 environment. Section 109(b) makes it clear that EPA must base the standard on this scientific
5 information but does not specify how EPA is to incorporate this information into the structure of
6 the standard beyond the requirement that any secondary NAAQS be a level of air quality that is
7 requisite to protect public welfare. Accordingly, while Sections 108 and 109(b) make it clear that
8 EPA must take this kind of information into account when setting the NAAQS, specific
9 directions are not provided on how to accomplish this.

10 It is also important to keep in mind the meaning of the term “requisite,” as interpreted by
11 EPA and the Courts. In considering the meaning of the term “requisite” in the context of the
12 primary standards, the Supreme Court has agreed with EPA that such a standard is one that is
13 “sufficient, but not more than necessary” to protect public health. *Whitman v. American*
14 *Trucking*, 531 U.S. 457, 473 (2001). In addition, while Section 109(b) requires that EPA
15 establish “national” standards, it does not define this requirement in any greater detail than
16 described above.

17 While EPA has most often made decision based on the results of direct exposure to an air
18 pollutant in the ambient air in assessing effects on public health and welfare in establishing
19 NAAQS, such as the health effects on humans when breathing in an air pollutant or the effects
20 on vegetation through the uptake of air pollutants through leaves, EPA has also considered,
21 where appropriate, the effects of exposure to air pollutants through more indirect mechanisms.
22 For example, in 1978, EPA established a NAAQS for lead that included non-respirable particles
23 based on the conclusion that these particles contribute to blood lead levels once they are
24 deposited on the ground and then become a source of human lead exposure through the ingestion
25 of lead-contaminated food or, particularly in children, the placing of hands and other
26 contaminated objects in the mouth. *Lead Industries v. EPA*, 647 F.2d 1130 (DC Cir. 1980). In
27 addition, EPA has previously considered acidification and nutrient enrichment concerns
28 associated with ambient concentrations of NO₂ and other nitrogen compounds, but determined
29 that the scientific and technical evidence that was available at the time did not provide an
30 adequate basis to set a separate secondary standard to address such effects. 61 FR 52852, 52855
31 (October 8, 1996). The deposition of ambient NO_x and SO_x to terrestrial and aquatic

1 environments can affect ecosystems through direct and indirect mechanisms, as discussed in
2 previous chapters. Given Congress’ instruction to set a standard that “is requisite to protect the
3 public welfare from “*any* known or anticipated adverse effects *associated with* the presence of
4 such air pollutant in the ambient air,” 42 U.S.C. § 109 (b)(2) (emphasis added), this review
5 appropriately attempts to take into consideration widely acknowledged effects, such as
6 acidification and nutrient enrichment, which are associated with the presence of ambient SO_x and
7 NO_x, as described in previous chapters.

8 The relationships between the impacts of ambient air concentrations on ecological effects
9 such as acidification and nutrient enrichment, however, are extremely complex. To take these
10 effects into account, including the variability in effects resulting from the same ambient
11 concentrations of SO_x and NO_x, EPA is developing a biologically relevant framework in this risk
12 assessment that relates ambient concentrations of an air pollutant to an ecological indicator,
13 taking into account relevant atmospheric and ecological variables. In that context it is useful to
14 explore whether a standard structured to take these elements into account is an appropriate way
15 to provide protection that is sufficient but not more than necessary, as compared to a standard
16 that treats ambient concentrations the same irrespective of these other biologically relevant
17 variables. The CAA, which requires EPA to establish “national” standards, does not specifically
18 address whether EPA can establish standards which could allow for varying levels of SO_x and
19 NO_x in different areas. However, such an approach may be necessary if EPA is to establish a
20 secondary standard that provides sufficient but not more than necessary to protect the public
21 welfare. Given the lack of specificity in Section 109(b), EPA believes it is appropriate to explore
22 the approach described in this chapter, with an aim of setting a national standard that provides
23 sufficient but not more than necessary protection throughout the United States by taking into
24 account the recognized variation of biologically relevant factors.

25 **8.3 LINKAGES FOR STRUCTURING ECOLOGICALLY RELEVANT** 26 **STANDARDS**

27 Consistent with the framework depicted in Figure 8.1-1, the following sections provide a
28 discussion of both the linkages between ambient air concentrations and resulting deposition
29 metrics, and between the deposition metric and the ecological indicator of concern. These
30 linkages are discussed conceptually in Section 8.3.1. Following that discussion, Section 8.3.2

1 describes the deposition transformation function that relates atmospheric concentrations of NO_x
2 and SO_x to nitrogen and sulfur deposition metrics, while Section 8.3.3 describes the ecological
3 effect function that transforms the deposition metric into the ecological indicator. Modifying
4 factors that alter the relationship between ambient atmospheric concentrations of NO_x and SO_x
5 and depositional loads of nitrogen and sulfur, and those that modify the relationship between
6 deposition loads and the ecological indicator are also discussed more fully.

7 **8.3.1 A Conceptual Model of the Pathway from Ambient Atmospheric** 8 **Concentrations of NO_x and SO_x to Ecological Effects**

9 For this first draft risk and exposure assessment, we have focused on developing an air
10 quality indicator for protection against the adverse effects of aquatic acidification. Figure 7.3-1
11 demonstrates the various levels of protection of fish populations associated with varying ANC
12 levels. Differing levels of an ecological indicator such as ANC will be associated with differing
13 levels of protection of ecosystem functions (e.g. fish populations, biological diversity, fish
14 population health). Different ecological indicators may be necessary for protection against the
15 adverse effects of terrestrial acidification and aquatic and terrestrial nutrient enrichment.
16 Indicators for other welfare effects categories of concern will be explored in future drafts of this
17 risk and exposure assessment.

18 We have created a system of linked functions that translate an atmospheric indicator
19 (concentrations of NO_x and SO_x) into an ecological indicator that expresses the potential for
20 deposition of nitrogen and sulfur to acidify an ecosystem (Figure 8.1-1). The first part of this
21 system is the deposition transformation function, which transforms ambient atmospheric
22 concentrations (the atmospheric indicators) into a deposition metric. In this particular example,
23 the deposition metric referred to in Figure 8.1-1. is the sum of atmospherically deposited
24 oxidized nitrogen and sulfur, weighted by their molar ratio. The second part of this system of
25 functions is an ecological effect function which converts the deposition metric to an ecological
26 indicator (ANC) by adjusting for levels of other ecological factors that contribute to ANC.
27 When the amounts of NO_x and SO_x in the ambient atmosphere can be used to derive a
28 deposition metric (via the deposition transformation function) that can then be used to derive a
29 level of ANC (through the ecological effect function) which falls within the range defined as
30 acceptable by the standard, then those levels of NO_x and SO_x will be considered to meet that

1 standard of protection. The atmospheric levels of NO_x and SO_x that satisfy a particular level of
2 ecosystem protection are those levels that result in an amount of deposition that is less than the
3 amount of deposition that a given ecosystem can accept without degradation of ANC. This
4 latter amount is referred to as the maximum deposition load and is the amount that solves a
5 mass-balance equation for that particular level of ANC. Mass-balance relationships developed in
6 the scientific literature provide the basis for established equations that link levels of ANC to
7 levels of nitrogen and sulfur deposition (given levels of other, non-deposition related parameters
8 such as weathering rates, base cation uptake rates). Quantitative examples of these relationships
9 will be provided in Section 8.4 of this chapter in the next draft of this document.

10 The value of the deposition metric can be calculated in terms of measured (or modeled)
11 ambient NO_x and SO_x concentrations by using established relationships between ambient
12 concentrations of NO_x and SO_x and nitrogen and sulfur deposition. As will be described in
13 Section 8.4, these relationships are derived from the physical processes that govern transport and
14 deposition; they are species dependent; they vary by particle mass, meteorology, compensation
15 point, terrain and land use type.

16 It is important to note for this type of air quality indicator that the same level of
17 protection from acidification can be obtained with different combinations of ambient NO_x and
18 SO_x concentrations and their associated levels of nitrogen and sulfur deposition. As a result, the
19 ambient air quality standard for NO_x and SO_x may be expressed jointly, and the air quality
20 standard may be met by various combinations of NO_x and SO_x.

21 Determining the level of the air quality indicator that is associated with a particular level
22 of protection against adverse effects of acidification may include consideration of factors that
23 could potentially alter the depositional relationship of nitrogen or sulfur to ANC. In addition, the
24 specific concentrations of ambient NO_x and SO_x that will result in a particular deposition metric
25 can vary based on factors that affect the relationship between ambient concentrations and
26 deposition. These two considerations imply that the levels of ambient NO_x and SO_x that would
27 be consistent with a particular level of ANC may differ according to the specific levels of
28 modifying factors in different locations and whether the ecosystem is nitrogen and/or sulfur
29 limited or saturated. While the standard is set to achieve the same level of protection (ANC)
30 nationwide, the atmospheric concentrations of NO_x and SO_x and the related amounts of
31 deposition that can occur and still meet the standard may differ. For example, because of the

1 differences in maximum depositional loads between watersheds with high proportions of
 2 grasslands and those with high proportions of evergreen forest ecosystems, for the same level of
 3 ANC, there may be differences in the ambient concentrations of NO_x and SO_x that would meet
 4 that ANC level. The response variable associated with these examples will vary across effects
 5 categories, and might be based on soil base saturation or aluminum concentrations in soil water
 6 or biodiversity impacts, rather than ANC. Specific modifying factors for the ambient
 7 concentration to deposition relationship are discussed in Section 8.3.2 of this chapter. Specific
 8 modifying factors for the deposition to acidification relationship will be discussed in Section
 9 8.3.3. The overall transformation function of the standard will be national in scope as will be the
 10 ecological indicator level, although specific modifying factors within the transformation function
 11 and maximum depositional load may vary by location. Therefore, while the Administrator may
 12 choose the level of protection that is desired by specifying a particular limiting value of ANC
 13 (ANC_{limit}), there are a series of transformations that link that level of protection to ambient
 14 concentrations of NO_x and SO_x. As a conceptual example, to express the previous discussion in
 15 mathematical terms, consider the following:

$$16 \quad I = \text{ANC}_{\text{limit}} = g(\cdot) - \text{DL}(\text{S}) - \text{DL}(\text{N}) \quad (1)$$

17 where I is an indicator, g(·) is a series of non-atmospheric, non-depositional environmental
 18 factors (e.g., runoff, weathering rates, geology, etc.), ANC_{limit} is a critical limit for ANC
 19 necessary to provide a particular level of protections for biota in an ecosystem, DL(S) is the
 20 implied maximum depositional load of S associated with that ANC limit, and DL(N) is the
 21 implied maximum depositional load of N associated with that ANC limit. Note that DL(S) and
 22 DL(N) are jointly determined by the equation g(·) - ANC_{limit} = DL(S) + DL(N), and that there is no
 23 unique solution for either one unless the other is fixed. Further, we can express the deposition of
 24 NO_x and SO_x through the following relationships:

$$25 \quad S_D = f_S(\text{NO}_x, \text{SO}_x | Z) \text{ and } N_D = f_N(\text{NO}_x, \text{SO}_x | Z) \quad (2)$$

26 where S_D is sulfur deposition, N_D is nitrogen deposition, Z is a series of parameters affecting
 27 deposition of ambient SO_x and NO_x (including NH₃), f_S is a function relating S_D to ambient
 28 concentrations of NO_x and SO_x, given Z, and f_N is a function relating N_D to ambient
 29 concentrations of NO_x and SO_x, given Z.
 30
 31

1 It is clear that the indicator of NO_x and SO_x effects (I) is a function of deposition.
2 However, by substituting in the ambient to deposition conversion functions in (2), the indicator I
3 can be made into an ambient indicator:

$$4 \quad I = g(\cdot) - S_D(f_S(\text{NO}_x, \text{SO}_x | Z)) - N_D(f_N(\text{NO}_x, \text{SO}_x | Z)). \quad (3)$$

5 Not all values of S_D and N_D cause adverse welfare effects. The point where the levels of
6 deposition associated with the ambient NO_x and SO_x lead to an ANC that does not achieve the
7 specified ANC limit are when S_D and N_D are equal to or greater than the DL(S) and DL(N) for a
8 particular location.

$$9 \quad I = g(\cdot) - DL(f_S(\text{NO}_x, \text{SO}_x | Z)) - DL(f_N(\text{NO}_x, \text{SO}_x | Z)). \quad (4)$$

10 Clearly, the establishment of the functional forms for g(·), f_S, and f_N, as well as for the
11 values of the parameter series Z, will be critical. Note that variable factors in g(·) and Z can be in
12 many forms, some continuous, e.g. precipitation, and some discrete, e.g. type of bedrock.
13 However, continuous variables can be converted to discrete categories, which may be easier to
14 implement in the form of a standard. For example, precipitation could be expressed as inches per
15 year, or precipitation could be categorized as high, medium, and low, possibly based on
16 percentiles of the observed distributions of precipitation values.

17 **8.3.2 Linkages between Ambient Concentrations and Deposition**

18 Defining the linkages between ambient concentrations of NO_x and SO_x and deposition of
19 N and S is a crucial piece of the framework as described in Section 8.1. Atmospheric deposition
20 of nitrogen from ambient concentrations of NO_x (N_{NO_x}) and atmospheric deposition of sulfur
21 from ambient concentrations of SO_x (S_{SO_x}) to a surface is a complex function of numerous
22 meteorological parameters and surface properties. A transformation function that incorporates
23 the nationwide variability of the important meteorological and surface parameters is expected to
24 be a useful approach to describing the deposition process. The derivation of this transformation
25 function, which converts ambient concentrations of NO_x and SO_x to their associated depositions,
26 is described below. The derivation takes algorithms from the U.S. EPA-developed Community
27 Multi-scale Air Quality (CMAQ) modeling system (U.S. EPA, 1999) into consideration, and
28 CMAQ output is used to calculate key parameters in the resulting function. The use of CMAQ

1 data is intended to be illustrative of the process. Specific models, data sets and methods
 2 appropriate to use in other locations or on different spatial/temporal scales may differ.

3 8.3.2.1 Data and Tools

4 8.3.2.1.1 Physics and Chemistry of Atmospheric Deposition

5 Atmospheric pollutants deposit onto land and water surfaces through at least two major
 6 mechanisms: direct contact with the surface (dry deposition), and transfer into liquid
 7 precipitation (wet deposition). The magnitude of each deposition process is related to the
 8 ambient concentration through the time-, location-, process- and species-specific *deposition*
 9 *velocity* (Seinfeld and Pandis, 1998):

$$10 \quad Dep_i^{Dry} = v_i^{Dry} \cdot C_i^{Amb} \quad (19)$$

$$11 \quad Dep_i^{Wet} = v_i^{Wet} \cdot C_i^{Amb} \quad (20)$$

12 where v_i^{Dry} and v_i^{Wet} are the dry and wet deposition velocities, Dep_i^{Dry} and Dep_i^{Wet} are the dry and
 13 wet deposition fluxes, C_i^{Amb} is the ambient concentration, and the i subscript indicates the
 14 pollutant species under study. The total deposition of each pollutant is

$$15 \quad Dep_i^{Tot} = Dep_i^{Dry} + Dep_i^{Wet} \quad (21)$$

16 Substituting Equations 19 and 20 into 21 yields

$$17 \quad Dep_i^{Tot} = v_i^{Dry} \cdot C_i^{Amb} + v_i^{Wet} \cdot C_i^{Amb} \quad (22)$$

18 The total deposition of sulfur and nitrogen would therefore be:

$$19 \quad Dep_{S+N}^{Tot} = \sum_i (v_i^{Dry} + v_i^{Wet}) \cdot m_i \cdot C_i^{Amb} \quad (23)$$

20 where m_i is the molar ratio of the atom (sulfur or nitrogen) of interest to the i th pollutant.

21 Ambient sulfur- and nitrogen-containing pollutants include gases such as sulfur dioxide (SO_2),
 22 ammonia (NH_3), various nitrogen oxides (NO , NO_2 , N_2O_5), nitric acid (HNO_3), and peroxyacetyl
 23 nitrate (PAN); and particulate species such as sulfate (SO_4^{2-}), nitrate (NO_3^-), and ammonium (NH_4^+)
 24). Oxides of nitrogen and oxides of sulfur are currently regulated by the NAAQS as defined in

1 the CAA. These compounds include the sulfur-containing species above as well as the above
2 oxidized forms of nitrogen (referred to here as NO_x); ammonia and ammonium, which are
3 reduced forms of nitrogen, are not currently included as regulated pollutants.

4 For this section, Equation 5 is termed the *deposition transformation function*, and it
5 provides the functional form of the relationship between instantaneous atmospheric pollutant
6 concentrations and deposition fluxes of sulfur and nitrogen. Equation 5 can also be thought to
7 represent the relationship between cumulative or average values of the variables over time and/or
8 space, but doing so raises issues of exactly how integration/averaging over time and/or space is
9 to be done. To some extent, the degree of temporal and spatial averaging needed will depend on
10 the ability to measure or model concentrations at particular spatial (e.g. at specific water bodies)
11 and temporal (e.g. hourly or daily) scales. However, the need for temporal or spatially
12 aggregated values may also be driven by the form of the ecological indicator, which may be
13 annual or seasonal, and may reflect local ecological attributes. The remainder of Section 8.3.2
14 discusses this and other policy-relevant scientific and mathematical issues and proposes methods
15 for determining parameter values (i.e. v_i^{dry} and v_i^{wet}) for input to Equation 23; results of example
16 calculations will be shown in Section 8.4.

17 **8.3.2.1.2 Air Quality Simulation Models**

18 Deposition velocities necessary for employing the deposition transformation function can
19 be obtained either through models or direct measurements, although the logistics of accurate
20 measurements are currently prohibitive to deploying a measurement network. For purposes of
21 the current discussion and accompanying future example calculations (Section 8.4), deposition
22 velocities will be obtained from CMAQ output, and research will be conducted to assess if this is
23 the optimal source of these parameters for future regulatory analyses.

24 CMAQ determines the deposition fluxes of the suite of modeled air pollutants at each
25 time step and location in the user-specified spatial grid. Deposition velocities can be found in
26 CMAQ output in the following ways:

- 27 1. v^{dry} values of gaseous pollutants are calculated in the CMAQ weather module called
28 the Meteorology-Chemistry Interface Processor (MCIP) through a complex function
29 of meteorological parameters (e.g. temperature, relative humidity) and properties of
30 the geographic surface (e.g. leaf area index, surface wetness)

- 1 2. v^{dry} values for particulate pollutants are calculated in the aerosol module of CMAQ,
2 which, in addition to the parameters needed for the gaseous calculations, also
3 accounts for properties of the aerosol size distribution
- 4 3. Equation 20 is not directly expressed in CMAQ source code: the cloud processing
5 module of CMAQ performs simulations of mass transfer into cloud droplets and
6 aqueous chemistry to incorporate pollutants into rainwater, all of which is
7 conceptually contained in the v^{wet} parameter. Explicit v^{wet} values can and will be
8 calculated in a post-processing step by applying Equation 20 to the wet depositions
9 and ambient concentrations output by the model.

10 Due to lack of direct measurements, no performance evaluations of CMAQ's dry
11 deposition calculations can be found; however, the current state of MCIP is the product of
12 development work that has been based on peer-reviewed literature from the past two decades
13 (U.S. EPA, 1999) and is considered to be EPA's best estimate of dry deposition velocities. Some
14 bias has been found between CMAQ's wet deposition predictions and measured values (Morris
15 et al., 2005); recent analyses suggest that poor simulation of precipitation could be responsible
16 for this (Davis and Swall, 2006), which should be considered if v_i^{wet} values are applied to
17 ambient measurements (this will be discussed in Section 8.4). Although the model is continually
18 undergoing improvement, CMAQ is EPA's state-of-the-science computational framework for
19 calculating deposition velocities, and was therefore the first choice for exploring inputs to
20 Equation 23. Gaseous v^{dry} values are directly output by MCIP, whereas aerosol v^{dry} values and all
21 species' v^{wet} values currently need to be calculated from standard CMAQ output. In all cases,
22 data aggregation issues need to be carefully considered: while MCIP and CMAQ give hourly
23 output values, the particular averaging time relevant for ecological processes for NO_x and/or SO_x
24 is likely to be the same as the time resolution of the ecological indicator of choice (which is
25 likely to be seasonal or annual). As such, values of v^{dry} and v^{wet} for each species should either be
26 representative of the desired averaging period, or aggregation of resulting depositions after
27 applying hourly deposition velocities to concentrations will need to be performed. It is proposed
28 that the spatial scale of the deposition velocities used here reflect that of future analyses, e.g. be
29 relevant at scales that may range from individual water bodies up to regional or national scales,
30 as discussed in Chapters 4 and 5. Specific methods for performing calculations are discussed
31 below.

8.3.2.2 Analytical Techniques

8.3.2.2.1 Hourly Deposition Velocities

As stated above, hourly v^{dry} values of gaseous species are complex functions of meteorology and land surface properties that are directly available in the outputs of MCIP. Deposition velocities of aerosol species were calculated by applying Equation 19 to CMAQ grids of aerosol C_i^{Amb} and Dep_i^{Dry} values to calculate hourly grids of v_i^{Dry} values for the example in a future Section 8.4. Plans are underway to obtain the size-specific aerosol deposition velocities from future CMAQ model runs. This will provide v_i^{Dry} for fine as well as coarse mode aerosols.

Unlike dry deposition velocity, the wet deposition velocity is independent of land surface properties and can be effectively expressed in a simple form (Seinfeld and Pandis, 1998):

$$v_i^{Wet} = w_i^r \cdot p^0 \quad (24)$$

$$w_i^r = \frac{C_i^{Aq}}{C_i^{Amb}} \quad (25)$$

where w^r is the washout ratio, C_i^{Aq} is the concentration of species i in the rainwater, and p^0 is the precipitation intensity.

If Equation 25 is substituted into 24, and Equation 24 is substituted into Equation 20, the ambient concentration cancels out and wet deposition can be expressed as a function of only rainwater concentration and precipitation (both of which are routinely measured by the NADP network). However, deposition can be expressed in terms of ambient concentrations since the atmospheric indicators for the framework discussed above are ambient air concentrations of NO_x and SO_x. Calculating the transformation function will therefore use values of w^r derived from CMAQ output for input into Equation 24 (which is then input to Equation 5). Solving for v_i^{wet} in Equations 2 and 6, equating them and rearranging these equations yields:

$$w_i^r = \frac{Dep_i^{wet}}{p^0 \cdot C_i^{Amb}} \quad (26)$$

which expresses the washout ratio in terms of CMAQ outputs. As mentioned above, simulated precipitation is a source of error in CMAQ's wet deposition simulations. The washout ratio offers a method of dealing with this error: CMAQ output can be used to calculate w_i^r values and

1 precipitation measurements (e.g. from NADP) can be used to obtain p^0 values for input to
2 Equation 24 to obtain measurement-corrected wet deposition velocities.

3 **8.3.2.2.2 Aggregation Issues**

4 The deposition velocities and washout ratio are dynamic quantities that depend on
5 numerous conditions that are highly variable with both space and time. However, an ecologically
6 relevant standard is likely to be based on an averaging time greater than hourly, which might
7 require temporal aggregation of deposition velocities. Use of deposition velocities that reflect a
8 spatial scale consistent with this review might also require some level of spatial aggregation of
9 deposition velocities. The v^{dry} , v^{wet} , and w_i^r parameters are all expressible as ratios of other
10 quantities, which allow for multiple possible methods of calculating aggregate values, including
11 the following:

- 12 1. Average ratio: Averaging hourly values of the deposition velocities and the washout
13 ratio is equivalent to find the average ratios of their component quantities.
- 14 2. Weighted Average Ratio: Same as above but weight the hour-specific ratios by either
15 concentration or deposition during the averaging process.
- 16 3. Ratio of averages: Instead of working with hourly deposition velocities and w_i^r
17 values, concentrations and depositions can be aggregated (concentrations are
18 averaged, depositions and precipitation are summed) over space and/or time and the
19 ratios of the resulting aggregate values can be computed. This method removes the
20 outlier effect of unrealistically large ratios resulting from small denominators (this
21 seems most likely to be an issue in calculations of w_i^r when precipitation is very small
22 but non-zero).
- 23 4. Conservatism: Single hourly values from the pool of available data can be chosen that
24 provide a more conservative calculation of depositions, such as predicting the largest
25 possible depositions from a given concentration (i.e., choose the annual and spatial
26 maximum values of v_i^{dry} and/or w_i^r for each species and region).

27 Temporal aggregation is known to produce substantial biases in dry depositions when
28 correlation exists between v^{dry} and C_i^{Amb} (Clarke et al., 1997), and it is presumed that this bias
29 needs consideration when performing spatial aggregation as well. Some exploration of the
30 conservative method of finding v^{dry} values has been performed, and it was found that this
31 produced unrealistically high annually representative values. For purposes of the example
32 calculations that will be presented in Section 8.4, the Ratio of Averages method will be
33 employed to obtain all annual deposition velocities, but the best method of obtaining such values
34 remains an open research question.


1 **8.3.2.3 Results and Findings**

2 Potential applications of the transformation function to the standard setting and
3 implementation processes will be presented in Section 8.4. In this section, the variability of the
4 deposition velocities resulting from CMAQ output is explored to 1) highlight the complexity of
5 the process of converting between ambient concentrations and depositions and 2) illustrate some
6 possible visualization techniques

7 Fields of dry deposition velocities of SO_2 output by MCIP are shown in **Figure 8.3-2**.
8 The figure shows maps of mean and standard deviation of hourly dry deposition velocities over
9 24 hours during four seasonally distinct days across the eastern U.S. output by MCIP. The plots
10 show that v^{dry} values vary by at least two orders of magnitude over the spatial domain, and that
11 there appears to be some spatial correlation between the magnitude of v^{dry} values (mean plots)
12 and the hourly variability (standard deviation plots). These maps highlight the difficulty of
13 choosing a single nationally representative v^{dry} value.

14 Therefore, v^{dry} values could also be studied at smaller spatial scales. **Figures 8.3-3** and
15 **8.3-4** show distributions of v^{dry} values from a Case Study Area (Pamlico Sound, NC) of the same
16 four days used for Figure 8.3-2. v^{dry} values in Figure 8.3-3 show an increasing trend in the
17 January 1 plot, a decreasing trend on April 1, and relatively stable trends on July 1 and October
18 1, but with much greater spatial variability in the July 1 plot. Figure 8.3-4 shows maps of the
19 diurnal distributions of the same data, and reveals that a substantial portion of the spatial
20 variability might be explained by differences in the dominant land use of the grid cells (see
21 **Appendix A** for the full list of land use categories in the model), especially when distinguishing
22 between water and land surfaces. This suggests that v^{dry} values specific to at least the land use
23 type, if not the grid cell, should be considered when using the transformation function.

1




2

3

4

5

Figure 8.3-2: Mean and standard deviations of $SO_2 v^{dry}$ values across the CMAQ eastern United States modeling domain for the first day of four seasonally distinct months. Note that the high value of the legend represents the maximum of the scale, not of the data.




1

2

3

4

Figure 8.3-3: Hourly distributions of $SO_2 v^{dry}$ values for one HUC (#3020105: Pamlico Sound, NC) over the first day of four seasonally distinct months.



1

2

3

4


5

Figure 8.3-4: Distributions of $SO_2 v^{dry}$ in each gridcell of one HUC (#3020105: Pamlico Sound, NC) over the first day of four seasonally distinct months. Colors indicate the dominant land use category (DLUSE) of the gridcell; see **Appendix A** for a description of each categorical value.

1
2 **Figure 8.3-5** shows domain-wide SO_2 washout ratios values for the same four seasonally
3 distinct days as above, calculated using the Average Ratio and the Ratio of Averages methods.
4 These plots illustrate that the potential for unrealistically extreme values is far greater using the
5 Average Ratio method. Exploration of the numbers in these calculations showed that, despite
6 excluding times when p^0 was 0, very low ambient concentration values could also cause extreme
7 values such as those shown in Figure 8.3-5. These results thus suggest that the Ratio of Averages
8 might be a better choice for calculations of w_i^r rather than the Average ratio method to help
9 mitigate the use of unrealistically large outlier values.

10 **8.3.2.4 Uncertainties**

11 The most prominent uncertainty in the process of converting between concentrations and
12 depositions is the known lack of accuracy of the deposition velocities that are input to Equation
13 23. The CASTNET and NADP networks measure concurrent weekly values of wet deposition
14 and ambient concentrations of select species; these data might be used to estimate the bias
15 present in calculations of v_i^{Wet} . Such data explorations are also expected to aid in understanding
16 the stability of w_i^r values, which might be low due to the dependence of cloud chemistry
17 dynamics on meteorology of the upper atmosphere. Estimates of deposition velocities might be
18 improved over CMAQ predictions by packaging the relevant portions of the MCIP and CMAQ
19 source code into a stand-alone tool that generates values using local meteorological
20 *measurements* (as opposed to the simulated values that are used during model runs). Further
21 discussions of uncertainty will be included in future drafts of this document.



1
2
3
4

Figure 8.3-5: Daily average washout ratios of SO_2 across the CMAQ eastern United States modeling domain for the first day of four seasonally distinct months (calculated using two different aggregation methods).

8.3.3 Linkages between Deposition and Ecological Indicators

The equation defining ANC forms the basis of the linkage between deposition and surface water acidic condition and the modeling approach used. In the discussion below, the functions are framed in terms of a maximum depositional load given a specified ANC limit, for ease of discussion. However the maximum depositional load is not the deposition metric employed in Fig 8.1. The deposition metric is the deposition associated with a certain measured or assumed level of atmospheric concentration of NO_x and SO_x, and under Fig 8.1 the deposition metric is used to calculate an ANC level associated with that atmospheric level. The next draft will also frame the functions in terms of calculating an ANC level from a specific deposition metric, whether it is above or below the depositional load.

Given an ANC concentration [ANC_{level}], the corresponding depositional load (DL) is simply the input flux of acid anions from atmospheric deposition, which gives the [ANC_{level}] when subtracted from the sustainable acidity or charge balance, defined as the uptake (N_{upt}), immobilization (N_{imm}), and denitrification (N_{den}) of NO_x deposition, the in-lake retention of NO_x deposition (N_{re}) and SO_x deposition (S_{ret}), and the acid anion leaching from the catchment as a function of the pre-industrial flux of base cations [BC]₀^{*} from weathering, as described in Equation 5:

$$DL(N_{NO_x}) + DL(S_{SO_x}) = \{fN_{upt} + (1 - r)(N_{imm} + N_{den}) + r(N_{ret} + S_{ret})\} + ([BC]_0^* - [ANC_{level}])Q \quad (5)$$

where *f* and *r* are dimensionless parameters that define the fraction of forest cover in the catchment and the lake/catchment ratio, respectively. The right hand side of equation (5) is equivalent to the *g*(·) function discussed in Section 8.3.1. To convert into fluxes, concentrations ([*x*] = *X*/*Q*) are simply multiplied by runoff (*Q*) (in m/yr) from the site. The charge balance equates the deposition inputs of acid anions with the sum of processes which control their long term storage, removal and leaching exports. Several major assumptions are made: 1) steady-state conditions, 2) the effect of nutrient cycling between plants and soil is ignored 3) there are no significant N inputs from sources other than atmospheric deposition, 4) ammonium leaching is negligible because any inputs are either taken up by biota, adsorbed onto soils or nitrate compounds, and 5) long-term sinks of sulfate in the catchment soils are negligible.

Using equation 5 to derive the terms for internal catchment processes, the charge balance provides those combinations of N_{NOx} and S_{SOx} deposition that defines the DL function given by:

$$DL_{max}(S_{SOx}) = (([BC]_0^* - [ANC_{level}])Q) / (1 - p_s) \quad (6)$$

$$DL_{min}(S_{SOx}) = 0 \quad (7)$$

$$DL_{max}(N_{NOx}) = fN_{upt} + (1-r)(N_{imm} + N_{den}) + (([BC]_0^* - [ANC_{level}])Q) / (1 - p_n) \quad (8)$$

$$DL_{min}(N_{NOx}) = fN_{upt} + (1-r)(N_{imm} + N_{den}) \quad (9)$$

where p_s and p_n define the fraction of in-lake retention of N_{ret} and S_{ret} , which is similarly proportional to the net input of N_{NOx} and S_{SOx} to the lake. It is not possible to define a single value of maximal loading of total acidity, since the acid anions sulfate and nitrate behave differently in the way they are transported with hydrogen ions; one unit of deposition of S will not have the same net effect on surface water ANC as an equivalent unit of N deposition. Equations 2 and 4 define the deposition loads for N_{NOx} and S_{SOx} individually. $DL_{max}(S_{SOx})$ defines the loading for sulfur when total N_{NOx} deposition is less than $DL_{min}(N_{NOx})$. $DL_{min}(N_{NOx})$ defines the deposition of N_{NOx} at which terrestrial catchment processes effectively remove N_{NOx} , so that deposition loads lower than $DL_{min}(N_{NOx})$ result in no leaching of nitrate. $DL_{max}(N_{NOx})$ defines the load for total N_{NOx} deposition when S deposition is zero. In reality, neither N_{NOx} and S_{SOx} deposition will ever be zero, so the DL for the deposition of one is fixed by the deposition of the other, according to the line defining in **Figure 8.3-1**. It is our intention to construct a similar line for N deposition originating from NO_x only for the second draft of this risk assessment.




Figure 8.3-1. The deposition load function defined by the model.

The thick lines indicate all possible pairs of DL of N_{NO_x} and S_{SO_x} acidity. Note that in the above formulation individual DL of N_{NO_x} and S_{SO_x} are not specified; each pair of depositions (N_{NO_x} and S_{SO_x}) fulfilling equations 2-5.

Given a $[ANC_{level}]$ for a desired level of protection, the modeling has to provide the calculations of the N_{NO_x} and S_{SO_x} sinks (N_{upt} , N_{imm} , N_{den} , N_{ret} , S_{ret}), landscape characteristics (f and r) and critical leaching of acid anions ($([BC]_0^* - [ANC_{level}])Q$), which are described below.

Given a maximum deposition load, ANC concentration at a site can be provided by the difference between future base cation and acid anion concentrations, which is equal to the predicted, steady-state sulfate (S_{leach}) and nitrate (N_{leach}) leaching fluxes:

$$[ANC] = [BC]_0^* - (S_{leach} + N_{leach})/Q \quad (10)$$

where S_{leach} and N_{leach} are:

$$S_{leach} = (1 - p_s)S_{dep} \quad (11)$$

$$N_{leach} = (1 - p_n)(N_{dep} - fN_{upt} - (1 - r)(N_{imm} + N_{den})) \quad (12)$$

8.3.3.1 Parameters

Net growth uptake of N_{NO_x} in forest (N_{upt})

At steady-state or over long timescales, the only permanent sink for N (from any source) in the terrestrial system occurs when there is removal of biomass from the catchment. Short term, seasonal cycling of N does not affect the mean annual budget over these timescales. The main removal of N from the catchment is through the harvesting of the forest or grazing. There are various ways for estimating this N loss from the catchment, which take into account the N content of various tree species and the time periods of harvesting cycle. For forests without harvesting, like the Adirondack region of New York, N_{upt} is zero.

Immobilization (N_{imm}) and denitrification (N_{den}) of N_{NO_x} in catchment soils

For long-term immobilization of N in forest soils, a constant value of 2 kg N / ha/ yr is used. This value represents the lower end of the range suggested from studies (Downing et al 1993). Thus, the amount of N removed for a given N_{NO_x} depositional load is simply proportional to its total N_{tot} load (N_{NO_x}/N_{tot})

It is assumed that the denitrification is related to the net input of N_{NO_x} loading into the soil system after accounting for the removal of N_{NO_x} by immobilization and uptake as follows:

$$N_{den} = fde(N_{nox} - N_{imm} - N_{upt}) \quad (13)$$

where fde is the denitrification fraction (value between 0 and 1), ascribed to the catchment soil. Denitrification is an anaerobic process and therefore a function of soil moisture content, with higher rates in peat soils. A typical upland soils denitrify N at a rate of 10% of N input while peat soils work at a rate of 70 to 80 %. Therefore, the average denitrification fraction for a catchment is approximated by the following linear relationship:

$$fde = 0.1 + 0.7f_{peat} \quad (14)$$

where f_{peat} is the proportion of peat soils within the catchment.

In-lake retention of N_{NO_x} and S_{SO_x}

The in-lake retention of acid anions is assumed to be a linear function of net input of acidity. It is sequentially the final sink of acidity encountered by inputs that move through the catchment.

1 The in-lake retention fractions for N_{NOx} (p_n) and S_{SOx} (p_s) are calculated from a kinetic equation
2 accounting for water retention time (Kelly et al 1987; Posch et al 1997).

$$3 \quad p_n = (s_n / s_n + Q/r) \text{ and } p_s = (s_s / s_s + Q/r) \quad (15)$$

4
5
6 where Q is the runoff and s_n and s_s are the net mass transfer coefficient for N_{NOx} and S_{SOx} . Due to
7 a lack of data, the net mass transfer coefficient for sulfate (s_s) was taken from a retention model
8 calibration to mass balance data of 11 lakes located in North America and northern Europe
9 (Baker and Brezonik 1988). The mean value of s_s for these lakes, 0.5 m/yr, was used for the
10 maximum depositional load calculations. The value for s_n , 5 m/yr, was obtained from a similar
11 study of 12, mostly Canadian, lakes (Dillon and Molot 1990).

12 **Pre-Industrial Base Cation Concentration**

13 The pre-industrial concentration of base cations $[BC]_0^*$ effectively set the long term
14 capacity of the catchment to neutralize acidic deposition, because it represents the only source of
15 base cation input that is sustainable over the long-term. Input of cations from weathering is
16 assumed to be a relatively constant process driven largely by the reaction of CO_2 with primary
17 minerals in the soils and bedrock. Base cations are removed by leaching from the soil solution
18 through surface water runoff. At a steady-state, the leaching rate of base cation occurs at lesser or
19 greater rates than the weathering supply. However, base cation leaching is not at steady-state
20 today because anthropogenic acid deposition actually increases the leaching of base cations
21 through ion-exchange within catchment soils. Soils contain a store of adsorbed base cations, as
22 measured as base saturation, which are derived from weathering, but have accumulated in the
23 soil over millennia, until eventually a steady-state is achieved, whereby the supply of base
24 cations from weathering was in approximate equilibrium with the removal of base cations by
25 rainwater, itself in equilibrium with the atmosphere. For this reason, $[BC]_0^*$ cannot be derived
26 from measured data in runoff, but derived from a empirical relationships (i.e., pre-industrial base
27 cation concentration).

28 The pre-industrial base cation concentration is the sum of weathering ($[BC^*_w]$) supply
29 plus base cation deposition ($[BC^*_{dep}]$), if it is assumed that base cation deposition has not

1 significantly changed since pre-industrial times, minus long-term average uptake of base cations
 2 in the biomass (i.e., the annual average removal of base cations due to harvesting):

$$3 \quad [BC]_0^* = [BC_w] + [BC_{dep}^*] - [BC_u]. \quad (16)$$

4 The first step is to quantify the proportion of measured base cation leaching which is
 5 derived from transient ion-exchange processes (BC_{ex}) and is proportional to the load of acid
 6 anions. This proposition is represented by the term ‘F’, calculated according to the methodology
 7 of Brakke and colleagues (1990):

$$8 \quad F = \sin(\pi/2 \cdot [BC^*]_t/[S]) \quad (17)$$

9 where $[BC^*]_t$ is the measured non-marine base cation concentration and S is a constant which
 10 varies regionally according to geology, but from empirical studies is taken as 400 $\mu\text{eq/L}$
 11 (Harriman and Christie, 1995). This constant determines the measured non-marine base cation
 12 concentration which represents a catchment likely to be unaffected by acid deposition; $[BC^*]_t =$
 13 S, $F = 1$ and base cation leaching is increased by exactly the value of the acid anion load,
 14 resulting in no change in the ANC of runoff. For values of $[BC^*]_t$ greater than S, F is 1.

15 This F value is then used to calculate the pre-industrial base cation concentration
 16 according to the following equation:

$$17 \quad [BC^*]_0^* = [BC^*]_t - F([AA]_t^* - [AA]_0^*) \quad (18)$$

18 where $[AA]_0^*$ is the pre-acidification concentration of non-marine acid anions from weathering
 19 and natural atmospheric sources and the measured leaching rate of non-marine base cations
 20 $[BC^*]_t$ represent the sum of weathering, non-marine deposition and leaching sources. We
 21 assumed the steady-state concentration of nitrate ($[AA]_0$) was zero ($[AA]_0^* = 0$). Since $[BC^*]_0^*$ is
 22 now known, the maximum depositional load can be defined by Equation 9.

23 **Catchment Parameters (r, f, and Q)**

24 Lake and catchment areas can be estimated from topographical maps (1:50,000) and from
 25 the EMAP lake survey. The annual runoff can be obtained by interpolation from a national
 26 runoff map (Krug et al., 1990) for water years 1951-80. The fraction of forest in the catchment
 27 can be estimated from 2001 National Land Cover Data (NLCD).

1 **8.4 EXAMPLE APPLICATIONS OF SECONDARY NAAQS**
2 **STRUCTURE**

3 This section is intended to provide an illustrative example applying actual data sets for
4 the Adirondacks to the above derivations and linkage calculations. It is our intent that this
5 section will be developed in the second draft of this document.

6 **8.5 REFERENCES**

- 7 Baker, L. A., and P. L. Brezonik. 1988. Dynamic model of in-lake alkalinity generation. *Water*
8 *Resources Research* 24: 65–74.
- 9 Brakke, D.F., A. Henriksen, and S.A. Norton. 1990. A variable F-factor to explain changes in
10 base cation concentrations as a function of strong acid deposition. *Verh. Internat. Verein.*
11 *Limnol.* 24:146–149.
- 12 Clarke, J., Edgerton, E., Martin, B., 1997. Dry deposition calculations for the Clean Air Status
13 and Trends Network. *Atmospheric Environment* 31 (21), 3667–3678.
- 14 Davis, J. M., Swall, J. L., 2006. An examination of the CMAQ simulations of the wet deposition
15 of ammonium from a Bayesian perspective. *Atmospheric Environment* 40, 4562–4573.
- 16 Dillon, P. J., and L. A. Molot. 1990. The role of ammonium and nitrate in the acidification of
17 lakes and forested catchments. *Biogeochemistry* 11:23–43.
- 18 Downing, R. J., J.-P. Hettelingh, and P. A. M. de Smet (eds.). 1993. Calculation and mapping of
19 critical loads in Europe, status report 1993. Coordination Center for Effects, National
20 Institute of Public Health and Environmental Protection (RIVM), Bilthoven, The
21 Netherlands.
- 22 Harriman R, Christie AEG. 1995 Estimating critical loads for biota: the steady-state water
23 chemistry Henriksen.model. *Critical Loads of Acid Deposition for United Kingdom*
24 *Freshwaters:7_8.*

- 1 Kelly, C. A., J. W. M. Rudd, R. H. Hesslin, D. W. Schindler, P. J. Dillon, C. T. Driscoll, S. A.
2 Gherini, and R. H. Heskey. 1987. Prediction of biological acid neutralization in acid
3 sensitive lakes. *Biogeochemistry* 3:129–140.
- 4 Krug, W. R., W. A. Gebert, D. J. Graczyk, D. L. Stevens, B. P. Rochelle, and M. R. Church,
5 1990. Map of Mean Annual Runoff for the Northeastern, Southeastern, and Mid-Atlantic
6 United States, Water Years 1951-1980. U.S. Geological Survey Water- Resources
7 Investigations Report 88-4904, Madison, Wisconsin.
- 8 Morris, R. E., McNally, D. E., Tesche, T. W., Tonnesen, G., Boylan, J. W., Brewer, P., 2005.
9 Preliminary evaluation of the community multiscale air quality model for 2002 over the
10 Southeastern United States. *Journal of the Air and Waste Management Association* 55,
11 1694–1708.
- 12 Posch, M, J.K.MA, R. Forsius, A. Henriksen, A. Wilander 1997 Exceedance of Critical Loads
13 for Lakes in Finland, Norway, and Sweden: Reduction Requirements for Acidifying
14 Nitrogen and Sulfur Deposition. *Environmental Management* Vol. 21, No. 2, pp. 291–
15 304.
- 16 Seinfeld, J., Pandis, S., 1998. *Atmospheric Chemistry and Physics*. John Wiley and Sons, Inc.,
17 New York.
- 18 U.S. EPA. 2008. Integrated Science Assessment for *Oxides of Nitrogen and Sulfur –*
19 *Environmental Criteria (Second External Review Draft)*. Prepared by U.S. Environmental
20 Protection Agency, National Center for Environmental Assessment-RTP Division, Office
21 of Research and Development, Research Triangle Park, NC. EPA/600/R-08-082. August.
- 22 U.S. EPA, 1999. Science Algorithms of the EPA Models-3 Community Multiscale Air Quality
23 (CMAQ) Modeling System. Tech. Rep. EPA/600/R-99/030, U.S. Environmental
24 Protection Agency, Washington DC.
- 25
26
27
28

1 **APPENDIX A**

2 Land Use Categories in MCIP (from USGS).

- 3 1. Urban Land
- 4 2. Dryland Cropland and Pasture
- 5 3. Irrigated Cropland and Pasture
- 6 4. Mixed Dryland and Irrigated Cropland
- 7 5. Cropland/Grassland Mosaic
- 8 6. Cropland/Woodland Mosaic
- 9 7. Grassland
- 10 8. Shrubland
- 11 9. Mixed Shrubland/Grassland
- 12 10. Savannah
- 13 11. Deciduous Broadleaf Forest
- 14 12. Deciduous Needleleaf Forest
- 15 13. Evergreen Broadleaf Forest
- 16 14. Evergreen Needleleaf Forest
- 17 15. Mixed Forest
- 18 16. Water
- 19 17. Herbaceous Wetland
- 20 18. Wooded Wetland
- 21 19. Barren or Sparsely Vegetated
- 22 20. Herbaceous Tundra
- 23 21. Wooded Tundra
- 24 22. Mixed Tundra
- 25 23. Bare Ground Tundra
- 26 24. Snow or Ice
- 27

United States
Environmental Protection
Agency

Office of Air Quality Planning and Standards
Health and Environmental Impacts Division
Research Triangle Park, NC

Publication No. EPA-452/P-08-005b
August 2008
