

2008-2009

Lime and Nutrient Recommendations

Contents

Basis of Nutrient Recommendations.....	2
Secondary Nutrients and Micronutrients.....	4
Soil Buffer Test	5
Tobacco	7
Corn	9
Soybean	10
Small Grains (Barley, Oats, Rye, Wheat, and Triticale)	11
Corn Silage and Small Grain Hay/Silage.....	12
Grain Sorghum	12
Canola	13
Hay and Pastures—New Seedings.....	13
Hay and Pastures—Established Stands of Legumes and Grass-Legume Mixtures.....	14
Hay and Pastures—Renovation of Grass with Clovers or Annual Lespedeza	15
Hay and Pastures—Established Stands of Cool-Season Grasses.....	16
Hay and Pastures—Warm-Season Forages.....	17
Hay and Pastures—Horse Pastures.....	18
Conservation and Wildlife Land	19
Lawns and General Turf	21
Tree Fruits, Blackberries, Raspberries, Blueberries, and Grapes	22
Strawberries	23

Basis of Nutrient Recommendations

Recommended nutrient additions, based on a soil test, are only made when a crop yield or economic response has been measured for that crop under Kentucky soil-climatic conditions. Many field studies have been conducted by the Kentucky Agricultural Experiment Station under Kentucky farm conditions to determine the extent of any primary, secondary, or micronutrient needs. Yield and soil test data from these studies serve as guidelines for establishing recommendations contained in this publication. Recommendations in this publication strive to supply the plant nutrients needed to achieve maximum economic return assuming good management practices.

Sometimes, nutrients are supplied at rates greater than those recommended by soil test, specifically when using manure and other waste products as a nutrient source. In these situations, one should never exceed the nitrogen recommendation in a given year. Some soils in Kentucky have naturally very high P soil tests. If soil test phosphorus (P) is greater than 400 lb P per acre, then environmental constraints may limit additional P applications. Special consideration should also be given to environmentally sensitive areas, such as fields near streams or springs, or in karst areas (which contain caves, sinkholes, or depressions). The Kentucky Agriculture Water Quality Act requires nutrient management planning for all operations 10 acres or larger regardless of the source of the plant nutrients (fertilizer, biosolids, or manure). For soils testing above 400 lb P per acre, please consult the USDA Natural Resources Conservation Service Practice Code 590 for specific guidance in utilizing organic sources of nutrients and for appropriate manure (biosolid) application guidelines. In all cases, applications of P (from any source) are not permitted if soil test levels exceed 1066 lb P per acre.

Nutrient recommendations in this publication are based on soil test values obtained using testing methods in the laboratories operated as part of the Kentucky Agricultural Experiment Station. This laboratory uses the Mehlich III solution to extract phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg),

and zinc (Zn). Water pH is determined in a 1:1 (v/v) ratio of de-ionized water:soil, and buffer pH is determined with the Sikora buffer (Soil Science Society of America Journal 70, 2006). These methods are described in Bulletin 190 of the Southern Cooperative Series (November 1984), *Procedures Used by the State Soil Testing Laboratories in the Southern Region of the United States*. The nutrient recommendations in this publication should not be used for soil test results obtained by other testing methods.

The recommendations assume average climatic and management conditions for Kentucky. Rates should be adjusted upward or downward to reflect any deviations from these assumptions.

The rates recommended are for production of a crop to be grown each year and will increase soil test values slowly for P and K. Using these recommended rates would likely take four years or longer of annual nutrient application at the recommended rates to result in appreciably higher soil test levels of P and K.

CEC and Percent Base Saturation

Values for cation exchange capacity (CEC) and percent base saturation (% BS) are reported for all soil samples analyzed routinely in the University of Kentucky Soil Testing Laboratories. The CEC is an estimate that is calculated according to accepted practices, with a modification to better reflect specific Kentucky conditions.

Reported values for CEC include an estimation of acidity (expressed as hydrogen, H), as reflected in the buffer pH measurement and extractable calcium (Ca), magnesium (Mg), and potassium (K) by the Mehlich III extracting procedure. The total CEC is the sum of milliequivalents per 100 grams of soil (meq/100 g) of the following cations: H, Ca, Mg, and K. The value for % BS is the sum of the meq/100 g of the basic cations (Ca, Mg, and K) expressed as a percentage of the total CEC. The dominant cation in most Kentucky soils with a pH above 5.8 is calcium (Ca) and hydrogen (H) when soil pH is below 5.8. Rarely does the amount of magnesium (Mg) or potassium (K) greatly affect CEC estimation or % BS.

While CEC is not used directly in making nutrient recommendations for agronomic crops, this information does affect lime recommendations. The buffer pH is directly related to the acidity or H fraction of the CEC. The values for % BS are a good indicator of the relative presence of the basic cations in the soil. Soils with 70% or greater BS are unlikely to limit agronomic crop growth due to acidity. Agronomic crop yields on soils with a % BS below 50 can be affected by excessive acidity. Other Extension publications or your county Cooperative Extension agent should be consulted for the soil pH associated with optimal growth and yield of specific horticultural or agronomic crops.

Soil Drainage Classes and Nitrogen Fertilization

Nitrogen fertilizer recommendations are related to soil drainage differences. Soils differ in the amount of water that infiltrates the soil surface and in the subsequent rate of percolation through the root zone. Important soil series are listed in Table 1. Their drainage class grouping there does not always coincide with the NRCS/USDA drainage classes (indicated by an asterisk).

Table 1. Soil drainage classes.			
Well-Drained Soils			
Allegheny Ashton Baxter Caneyville Crider Cuba Elk	Frederick Hagerstown Huntington Jefferson Loradale Maury McAfee	Memphis Nolin Pembroke Pope Shelbyville Shelocta Trimble	Uniontown Vicksburg Wellston Wheeling Whitley
Moderately Well-Drained Soils			
Beasley* Bedford Captina Collins Commerce	Cotaco Eden* Grenada Heitt* Loring	Lowell* Mercer Morehead Mountview Nicholson	Otwell Sadler Tilsit Zanesville
Somewhat Poorly and Poorly Drained Soils			
Belknap Bonnie Calloway	Falaya Henry Johnsburg	McGary Melvin Newark	Stendal Tyler Weinbach
* NRCS/USDA drainage classes for these soils may be different from above.			

Soils with Naturally High Contents of P and K

Some soils naturally contain higher levels of P and K, thus having the ability to supply higher amounts for crop production. Soils developed from phosphatic limestones will likely maintain high soil test P levels without fertilization as will some soils containing high native K levels.

Table 2. Converting elemental and oxide values.

To Convert		Multiply By:
From:	To:	
P ₂ O ₅	P	0.44
P	P ₂ O ₅	2.29
K ₂ O	K	0.83
K	K ₂ O	1.20

Elemental and Oxide Values for P and K

Soil test values for phosphorus and potassium are reported as pounds of elemental P or K per acre (lb/A). Nutrient recommendations are made on the oxide basis: pounds of phosphate (P₂O₅) or potash (K₂O) per acre. Use the factors in Table 2 when converting from elemental to oxide and vice versa.

Plant Analysis

A plant analysis may be used to verify a suspected nutrient problem or to evaluate the nutrient status of a crop. Plant analysis is not a substitute for a soil test but should be used along with a soil test. Your county Extension agent has information on plant analysis services available for various crops, or see University of Kentucky Extension publication AGR-92, *Sampling Plant Tissue for Nutrient Analysis*.

Collecting a Representative Sample

In order to get reliable recommendations, it is important that the submitted soil sample accurately represent the field or area from which it was taken. Analytical results provided on the soil test report form are for the sample submitted, and the listed recommendations are based on those results. All recommendations are made on the assumption that a properly representative soil sample was properly taken. If soil sampling procedures are questionable, accurate nutrient and lime recommendations for the sampled field or area cannot be assured. Because results vary somewhat between fall and spring, it is better to consistently sample any given field in fall or spring. See University of Kentucky Extension publication AGR-189, *Managing Seasonal Fluctuations of Soil Tests*, for details. When sampling untilled fields in the fall, an equal number of cores should be collected from both between and close to the rows. Following a prolonged drought, soil water pH can be as much as 0.5 pH unit lower if the soil was not rewet prior to sample collection. To calculate a lime recommendation under these conditions, assume that the current water pH is about 0.5 units lower than normal and that the buffer pH has not been affected. Using the tables in this publication, apply lime as recommended with the estimated water pH (now raised by 0.5 pH units) and the sample buffer pH.

Sampling Depth and Frequency

For tilled areas, take soil cores to a depth of 6 to 7 inches. With pastures, lawns, no-tilled areas, and turf, take soil cores to a depth of 3 to 4 inches. Because of high fertilizer and commodity prices, each production field should be sampled every two years. Annual sampling is preferable for high-value crops

that remove large amounts of soil nutrients, such as alfalfa and double-crop silage. Sampling each year after manure application is also recommended. See University of Kentucky Extension publication AGR-16, *Taking Soil Test Samples*, for details.

Nutrient Recommendations Samples Testing Low in P and K

If soil tests for phosphorus and potassium are low, one-third to one-half of the recommended amounts of P₂O₅ and/or K₂O for corn can be used if it is banded 2 to 4 inches from the row.

Samples Testing High in P and K

When soil test levels for P and K are so high that no nutrient recommendation is made for the current year, there is no assurance that these high levels will be maintained for optimal production in the following years. When soil test levels are in the lower portion of the high range, the area should be sampled again the following year.

Recommendations without Soil Tests

If nutrient recommendations must be made without soil test results, assume low levels of residual N, P, and K.

Recommendations for Multiple Years

If one nutrient recommendation is made for two years of sequential cropping, the recommended rates of phosphate and potash for each crop are added together and applied to the first crop grown. However, this method is not recommended for crops with a high nutrient demand, e.g., alfalfa, corn for silage, tobacco, etc. Double-cropping recommendations for small grains and soybean can be found in the “Small Grains” and “Soybean” sections.

Nutrient Value of Manures and Tobacco Stalks

Animal manure and tobacco stalks add nutrients when applied to soils. These should be considered when deciding on materials to use in fulfilling crop nutrient recommendations. The best method to determine the nutrient content of these materials is through sampling and analysis of the manure/stalks. In cases where it is not possible to take samples in a timely manner, Table 3 can serve as a guide in estimating nutrients contained in the materials listed until samples can be obtained and tested.

When applying organic materials to soil, it is important to remember that some of the nutrients they contain are not as available to the next crop as those nutrients contained in commercial fertilizers. While almost 100% of the potash is available, only about 80% of the phosphate is expected to be available to the next crop. Organic nitrogen availability is variable, with nitrogen from manure especially dependent on livestock species/diet, storage and handling methods, and the timing and method of application.

Animal manures also contain significant amounts of calcium, magnesium, sulfur, zinc, copper, and molybdenum that may be of value to crops. The added organic matter can also be of significant benefit to soils low in organic matter.

University of Kentucky Extension publication AGR-146, *Using Animal Manures as Nutrient Sources*, and a computer

Table 3. Typical total nitrogen, phosphate, and potash content (pounds of nutrient per ton) of some manures and tobacco stalks.

Animal Manures ¹	Water (%)	Lb/Ton		
		N ₂	P ₂ O ₅	K ₂ O
Dairy cattle	80	11	9	12
Swine	80	9	9	8
Beef	80	11	7	10
Broiler litter	20	55	55	45
Broiler layers	40	35	55	30
Broiler pullets	30	40	45	40
Goat	70	22	5	15
Horse	80	12	6	12
Tobacco stalks	20	30	10	70

¹ Animal manures contain chloride, which can reduce the quality of tobacco. Limit rates to 10 tons per acre of cattle or swine manure. Poultry manure should not be applied to land in the year in which tobacco will be grown.
² Plant-available N can range from 20 to 80% of the total N in the year of application. See University of Kentucky Extension publication AGR-146, *Using Animal Manures as Nutrient Sources*, for more details.

spreadsheet (<http://soils.rs.uky.edu/manureprogram.htm>) are available to help determine application rates and fertilizer credits. Additional assistance can be obtained at your county Cooperative Extension Service office.

Tobacco stalks that are stored under cover to prevent leaching of nutrients by rainfall are a significant source of nutrients. See University of Kentucky Extension publication AGR-23, *Tobacco Stalks and Stems Fertility Value and Use*, for more information on the storage and use of tobacco stalks.

Surface Mine Reclamation

See the following University of Kentucky Extension publications:

- AGR-40, *Lime and Fertilizer Recommendations for Reclamation of Surface-Mined Coal Spoils*
- AGR-41, *Sampling Surface Mine Lands before and after Mining*.

Secondary Nutrients and Micronutrients

Magnesium

Magnesium levels in soils range from very high (in loess-derived soils) to low (some sandstone-derived soils). Despite low-testing soils, crop yield responses to magnesium fertilization have been very limited. It is sufficiently important in some areas in determining CEC that a soil test is offered. Magnesium needs of animals can best be met by direct feeding rather than with magnesium fertilization. Table 4 gives soil test levels and associated magnesium rate recommendation.

Table 4. Recommendations for magnesium.

Soil Test Level	Lb Mg/A ¹	Oz Mg/100 Sq Ft ²
0 - 6	50	2
7 - 18	45	2
19 - 30	40	2
31 - 42	35	1
43 - 54	30	1
55 - 60	25	1
Above 60	0	0

¹ These rates may be applied when no lime is needed or where dolomitic lime is not available. When lime is needed, the addition of dolomitic lime is preferred.
² Epsom salts (MgSO₄·7H₂O, 10% Mg) is readily available and may be more convenient for applying Mg to small areas.

Iron, Copper, Sulfur, Boron, and Molybdenum

Responses to added iron, copper, or sulfur have never been found, nor have symptomatic indicators been observed in Kentucky. Yield responses to boron and molybdenum have been observed for certain crops under certain conditions. Boron is recommended for topdressing on alfalfa. Consult the sections on tobacco, soybean, alfalfa, and pasture renovation with legumes for molybdenum recommendations.

Zinc and Manganese

Yield responses to zinc applications on corn and to foliar applications of manganese on soybean have been observed in Kentucky. The responses to manganese on soybean have been on a few soils in Daviess, McLean, and Webster counties. After diagnosis of manganese deficiency, responses to foliar applications have been superior to soil applications of manganese at planting. To date, soil testing has not been advantageous in solving any isolated cases of manganese deficiency in Kentucky soils.

Zinc deficiency in corn is significant in central and south-central Kentucky and in other areas with high soil test P levels. A soil test for zinc is performed routinely on all samples submitted to the University of Kentucky Soil Testing Lab. The zinc test results along with the soil test results for P and soil pH are used to identify soils/fields needing zinc.

Table 5. Crop nutrient removal values at standard harvest moisture.

Crop	Yield Unit	Nutrients Removed		
		N	P ₂ O ₅	K ₂ O
Lb/Yield Unit				
From IP-56:				
Alfalfa hay	ton	50	14	55
Grass/legume hay	ton	35	12	53
Fescue hay	ton	35	18	50
Pasture forage	ton	10	4	16
Corn for grain	bu	0.7	0.4	0.35
Silage corn	ton	7.5	3.5	8
Corn stalks	ton	14	7	29
Wheat grain	bu	1.2	0.5	0.3
Wheat straw	ton	12	4	20
Wheat forage/silage	ton	44	4	20
Sorghum grain	bu	1	0.4	0.3
Sorghum fodder	ton	12	5	17
Soybean grain	bu	3	0.7	1.1
Soybean hay (R3)	ton	100	7	12
Soybean hay (R5)	ton	150	12	27
Burley tobacco	100 lb	7	1.1	7.5
Dark-air tobacco	100 lb	7	0.6	6.0
Dark-fired tobacco	100 lb	7	0.6	6.0
Barley grain	bu	0.9	0.4	0.3
From NRCS Code 590:				
Rye grain	bu	1.16	0.33	0.32
Oats grain	bu	0.6	0.25	0.2
Bermudagrass hay	ton	37.6	8.7	34
Reed canary hay	ton	27	8.2	25
Eastern gamma hay	ton	35	16	31.2
Other warm-season hay	ton	20	6.8	25

Nutrients Removed by Agronomic Crops

Good nutrient management involves effective use of applied nutrients at rates utilized by crops. As a basis of assessing long-term soil fertility trends, crop nutrient removal should be used. Crop nutrient removal is the quantity of nutrients removed from a field in the harvested portion of the crop. This should not be confused with crop nutrient uptakes, which is the total amount of nutrients taken up by the entire crop (roots, stems, leaves, and seed) in a field. For quick reference, Table 5 includes crop nutrient removal values, published in University of Kentucky Extension publication IP-56, *Assessment of the Potential for Livestock and Poultry Manure to Provide the Nutrients Removed by Crops and Forages in Kentucky*, and in "NRCS Nutrient Management Standard Code 590."

Soil Buffer Test

The pH of the soil is a measurement made from a mixture of soil and water and is reported for all samples. The soil buffer test is performed and used to determine lime requirement for samples with a soil pH of 6.4 or below. In the soil buffer test, a buffer solution is mixed with soil, and the pH of the resulting suspension is measured. The result from the buffer test is reported as buffer pH. The buffer pH together with the soil pH can be used to determine the lime requirement to change the soil pH to a greater desired level. The University of Kentucky soil test laboratories started using the Sikora buffer in 2005, replacing the SMP buffer which contained hazardous chemicals (chromium and p-nitrophenol). The Sikora buffer provides the same buffer pH as the SMP buffer, so changes in agronomic interpretation of the data were not required.

All recommendations for agricultural lime are now based on 100% effective lime (100% RNV). Therefore, the lime recommendations in Tables 6, 7, and 8 are not for bulk lime but for 100% effective lime.

To determine how much lime is required to raise soil-water pH, see the table with your target pH in the heading. Tables 6, 7, and 8 give lime rate recommendations for target soil pH values of 6.4, 6.6, and 6.8, respectively. Limestone rate recommendations in these tables are given for water pH and buffer pH combinations in increments of 0.2 pH units. (Computerized recommendations use 0.1 pH unit increments.) To determine limestone rates, read down the left side of the appropriate table to the sample's water pH value, then read across until reaching the column head with the appropriate buffer pH value. If hydrated lime is applied, use the rate shown as 100% effective lime.

Table 6. Rate of 100% effective limestone (tons/acre) needed to raise soil pH to 6.4.

Water pH of Sample	Buffer pH of Sample								If Buffer pH Is Unknown
	5.5	5.7	5.9	6.1	6.3	6.5	6.7	6.9	
4.5	4.67	4.0	4.0	3.33	2.67	2.67	2.0	2.0	2.67
4.7	4.67	4.0	4.0	3.33	2.67	2.67	2.0	2.0	2.67
4.9	4.67	4.0	4.0	3.33	2.67	2.67	2.0	2.0	2.67
5.1	4.67	4.0	3.33	3.33	2.67	2.0	2.0	1.33	2.67
5.3	4.67	4.0	3.33	2.67	2.67	2.0	2.0	1.33	2.33
5.5	4.0	3.33	3.33	2.67	2.67	2.0	1.33	1.33	2.00
5.7	4.0	3.33	2.67	2.67	2.0	2.0	1.33	1.33	1.67
5.9	-	3.33	2.67	2.0	2.0	1.33	1.33	0.67	1.33
6.1	-	-	2.0	2.0	1.33	1.33	0.67	0.67	1.00

Table 7. Rate of 100% effective limestone (tons/acre) needed to raise soil pH to 6.6.

Water pH of Sample	Buffer pH of Sample								If Buffer pH Is Unknown
	5.5	5.7	5.9	6.1	6.3	6.5	6.7	6.9	
4.5	4.67	4.67	4.00	4.00	3.33	2.67	2.67	2.00	4.00
4.7	4.67	4.67	4.00	4.00	3.33	2.67	2.67	2.00	3.67
4.9	4.67	4.67	4.00	4.00	3.33	2.67	2.67	2.00	3.33
5.1	4.67	4.67	4.00	3.33	2.67	2.67	2.67	2.00	3.00
5.3	4.67	4.67	4.00	3.33	2.67	2.67	2.00	1.33	2.67
5.5	4.67	4.00	3.33	3.33	2.67	2.00	2.00	1.33	2.33
5.7	4.00	4.00	3.33	2.67	2.67	2.00	1.33	1.33	2.00
5.9	-	3.33	2.67	2.67	2.00	2.00	1.33	1.33	1.67
6.1	-	-	2.67	2.00	2.00	1.33	1.33	0.67	1.33
6.3	-	-	-	2.00	1.33	1.33	0.67	0.67	1.00

Table 8. Rate of 100% effective limestone (tons/acre) needed to raise soil pH to 6.8.

Water pH of Sample	Buffer pH of Sample								If Buffer pH Is Unknown
	5.5	5.7	5.9	6.1	6.3	6.5	6.7	6.9	
4.5	4.67	4.67	4.67	4.00	3.33	3.33	2.67	2.00	4.00
4.7	4.67	4.67	4.67	4.00	3.33	3.33	2.67	2.00	4.00
4.9	4.67	4.67	4.67	4.00	3.33	3.33	2.67	2.00	4.00
5.1	4.67	4.67	4.67	4.00	3.33	3.33	2.67	2.00	4.00
5.3	4.67	4.67	4.67	4.00	3.33	2.67	2.00	2.00	3.67
5.5	4.67	4.67	4.00	3.33	3.33	2.67	2.00	2.00	3.33
5.7	4.67	4.67	4.00	3.33	3.33	2.67	2.00	1.33	3.00
5.9	-	4.00	3.33	3.33	2.67	2.67	2.00	1.33	2.33
6.1	-	-	3.33	2.67	2.67	2.00	1.33	1.33	2.00
6.3	-	-	-	2.00	2.00	2.00	1.33	0.67	1.67
6.5	-	-	-	-	1.33	1.33	1.33	0.67	1.33

Because agricultural limestone is rarely 100% effective, the bulk lime rate required will need to be determined for each available lime source, based on the RNV. The bulk lime recommendation can be determined from Table 9 using the 100% effective lime recommendation and the RNV for the lime to be purchased. It can also be determined using this formula:

$$\text{Bulk Lime Rate} = \frac{100\% \text{ Effective Lime Rate Recommendation}}{\text{RNV}} \times 100$$

The bulk lime values in Table 9 are rounded to the nearest 0.5 ton. Therefore, the amount calculated by the formula above will not always be the same as the amount in Table 9.

The adjustment of soil pH by limestone is affected by: (1) thoroughness of mixing into the soil; (2) depth of mixing into soil (top 6 inches is assumed); (3) time of reaction (four years are needed

for complete reaction of limestone; however, the reaction time for hydrated lime is much shorter); (4) quality of limestone; and (5) use of acid-forming nitrogen fertilizers that can lower the effective soil pH obtained. When applying lime rates greater than 4 tons per acre (T/A), the lime should be thoroughly mixed in the plow layer by applying one-half the recommended rate before plowing and the other half after plowing followed by discing. Consult Uni-

versity of Kentucky Extension publication ID-163, *Determining the Quality of Aglime: Relative Neutralizing Value (RNV)*, and your county Extension agent about lime quality information for your area. Adjusting lime applications based on relative neutralizing values and comparing economic values of various lime sources in the state can be done with calculators found on the Internet at <http://soils.rs.uky.edu/calculators.php3>.

Table 9. Conversion table to bulk lime (rounded to the nearest 0.5 tons).

RNV % Unknown ¹	Recommended 100% Effective Lime Rate												
	0.67	1.00	1.33	1.67	2.00	2.33	2.67	3.00	3.33	3.67	4.00	4.33	4.67
	Bulk Lime Tons/Acre												
40	2.0	2.5	3.5	4.0	5.0	6.0	6.5	7.5	8.5	9.0	10.0	11.0	12.0
42	1.5	2.5	3.5	4.0	4.5	6.0	6.5	7.0	8.0	9.0	9.5	10.5	11.0
44	1.5	2.5	3.0	3.5	4.5	5.5	6.0	7.0	7.5	8.5	9.0	10.0	10.5
46	1.5	2.0	3.0	3.5	4.0	5.0	5.5	6.5	7.5	8.0	9.0	9.5	10.0
48	1.5	2.0	3.0	3.5	4.0	5.0	5.5	6.5	7.0	7.5	8.5	9.0	10.0
50	1.5	2.0	3.0	3.5	4.0	4.5	5.5	6.0	6.5	7.5	8.0	8.5	9.5
52	1.5	2.0	2.5	3.0	4.0	4.5	5.0	6.0	6.5	7.0	7.5	8.5	9.0
54	1.5	2.0	2.5	3.0	3.5	4.5	5.0	5.5	6.0	7.0	7.5	8.0	9.0
56	1.5	2.0	2.5	3.0	3.5	4.0	5.0	5.5	6.0	6.5	7.0	8.0	8.5
58	1.5	1.5	2.5	3.0	3.5	4.0	4.5	5.0	6.0	6.5	7.0	7.5	8.0
60	1.0	1.5	2.0	2.5	3.0	4.0	4.5	5.0	5.5	6.2	6.5	7.5	8.0
62	1.0	1.5	2.0	2.5	3.0	3.5	4.0	5.0	5.5	6.0	6.5	7.0	7.5
64	1.0	1.5	2.0	2.5	3.0	3.5	4.0	4.5	5.0	6.0	6.5	7.0	7.5
66	1.0	1.5	2.0	2.5	3.0	3.5	4.0	4.5	5.0	5.5	6.0	6.5	7.0
68	1.0	1.5	2.0	2.5	3.0	3.5	4.0	4.5	5.0	5.5	6.0	6.5	7.0
70	1.0	1.5	2.0	2.5	3.0	3.5	4.0	4.5	5.0	5.5	6.0	6.5	7.0
72	1.0	1.5	2.0	2.5	3.0	3.5	3.5	4.0	4.5	5.0	5.5	6.0	6.5
74	1.0	1.5	2.0	2.5	3.0	3.5	3.5	4.0	4.5	5.0	5.5	6.0	6.5
76	1.0	1.5	2.0	2.0	2.5	3.0	3.5	4.0	4.5	5.0	5.5	5.5	6.0
78	1.0	1.5	1.5	2.0	2.5	3.0	3.5	4.0	4.5	4.5	5.0	5.5	6.0
80	1.0	1.5	1.5	2.0	2.5	3.0	3.5	4.0	4.0	4.6	5.0	5.5	6.0
82	1.0	1.5	1.5	2.0	2.5	3.0	3.5	3.5	4.0	4.5	5.0	5.5	5.5
84	1.0	1.0	1.5	2.0	2.5	3.0	3.0	3.5	4.0	4.5	5.0	5.0	5.5
86	1.0	1.0	1.5	2.0	2.5	2.5	3.0	3.5	4.0	4.5	4.5	5.0	5.5
88	1.0	1.0	1.5	2.0	2.5	2.5	3.0	3.5	4.0	4.0	4.5	5.0	5.5
90	1.0	1.0	1.5	2.0	2.0	2.5	3.0	3.5	3.5	4.0	4.5	5.0	5.0
92	1.0	1.0	1.5	2.0	2.0	2.5	3.0	3.5	3.5	4.0	4.5	4.5	5.0
94	1.0	1.0	1.5	2.0	2.0	2.5	3.0	3.0	3.5	4.0	4.5	4.5	5.0
96	1.0	1.0	1.5	1.5	2.0	2.5	3.0	3.0	3.5	3.8	4.0	4.5	5.0
98	1.0	1.0	1.5	1.5	2.0	2.5	2.5	3.0	3.5	3.8	4.0	4.5	5.0
100	1.0	1.0	1.5	1.5	2.0	2.5	2.5	3.0	3.5	3.5	4.0	4.5	4.5

¹ Unknown RNV assumes 67%.

TOBACCO

Lime

Limestone should be applied in the fall and thoroughly mixed with the soil one to two years ahead of the crop. If applied in the spring before transplanting, or if more than 4 T/A are applied, plow one-half down and disc in the other half for soils with a water pH below 6.0.

Rates—If water pH is below 6.4, see page 5, and use the rate appropriate for a target pH of 6.6.

Nitrogen

Rates—Nitrogen fertilization rates (see Table 10) depend primarily on the field cropping history and soil drainage class. See page 2 for soil drainage classes. Rotation to other crops is strongly recommended after two or more years of burley tobacco production in the same field. More frequent rotation may be necessary when growing dark tobacco or burley tobacco varieties with low levels of disease resistance.

Sources—All commonly available N sources can be used satisfactorily on tobacco, particularly on well-drained soils where a good liming program is followed and soil pH is maintained in the range of 6.0 to 6.6. If soil pH is moderately to strongly acid (pH 6.0 or less) and no lime is applied, using a nonacid-forming source of N (sodium nitrate, calcium nitrate, or sodium-potassium nitrate) will lower the risk of manganese toxicity. Use these sources (or ammonium nitrate or potassium nitrate) for sidedressing because nitrate-nitrogen is more mobile in soil than ammonium nitrogen. If tobacco is grown on sandy soils or soils that tend to waterlog regardless of pH, using ammonium sources (urea, ammonium nitrate, ammoniated phosphates, ammonium sulfate, nitrogen solutions) will lower the risk of leaching and denitrification losses.

Time and Method—The entire nitrogen requirement can be broadcast pre-plant on well-drained soils. However, Kentucky often has large amounts of rainfall during April and May, so applying the broadcast nitrogen as near to transplanting as possible will significantly lessen the chances of loss. Apply the nitrogen after plowing and disc into the surface soil.

Because losses of fertilizer nitrogen can occur on sandy soils or soils with poor drainage, it is helpful to split nitrogen applications on these soils, applying one-third of the nitrogen before transplanting and the remaining amount two or three weeks after transplanting. The use of poorly drained or somewhat poorly drained soils for tobacco production is not recommended.

Further efficiencies in nitrogen use, decreased manganese toxicity, and increased early growth can be obtained by banding (sidedressing) most

of the nitrogen after transplanting. These bands should be applied 10 to 12 inches to the side of the row in either one or two bands and at depths of 4 to 5 inches. The nitrogen should be banded at one time 0 to 10 days after transplanting or split into two applications with two-thirds at 0 to 10 days and one-third at four to five weeks after transplanting. If one-third or more of the total nitrogen is applied after transplanting, the rate from Table 10 should be reduced by 15 to 25 lb N per acre.

Animal Manures

Animal manures are known to contain chloride in concentrations high enough to reduce the quality of cured tobacco. Cured tobacco leaf containing more than 1% chloride is considered unacceptable by the tobacco industry. Cattle and swine manure applications should be limited to no more than 10 tons per acre. Poultry manures should not be applied in the year tobacco is grown. Fall applications of poultry litter should not exceed 4 tons per acre on ground where tobacco will be planted the following spring. Fall manure applications should be made only when a living cover crop will be present to take up and recycle some of the available N.

Phosphate and Potash

Rates—Phosphorus and potassium fertilizer additions should be guided by soil testing. Based on soil test results, apply the recommended amounts indicated in Table 11. Research indicates that when soil test potassium is below 225 lb per acre, a broadcast application of potassium fertilizer is more effective than banding.

Sources—Research at the University of Kentucky has shown that applications after January 1 of chloride-containing nutrient sources such as muriate of potash at rates greater than 50 lb of chloride per acre lead to excessive levels of chloride in the cured burley tobacco leaf, increased curing and storage problems, decreased combustibility of the leaf, and, ultimately, greatly reduced quality and usability of the cured leaf. Consequently, sulfate of potash should be the major potassium fertilizer used after January 1. Because animal manures contain chloride, applications of dairy and swine manure should not exceed 10 tons per acre. Poultry manures should not be applied in the year tobacco is grown. Excessive rates of manure or manure used in conjunction with chlorine-containing fertilizers may result in unacceptable chlorine levels in the cured leaf.

Molybdenum

Molybdenum (Mo) is recommended for use on burley tobacco either as a broadcast soil application or dissolved in transplant setter water when the soil pH is below 6.6. Recent research and field trials have shown that setter water applications are equally as effective as broadcast applications in supplying molybdenum to the crop. Molybdenum can be purchased in dry solid or liquid forms. Either source is satisfactory when molybdenum is needed.

Soil Broadcast—Apply at the rate of 1 lb of sodium molybdate (6.4 oz of molybdenum) per acre. Dissolve this amount of dry sodium molybdate (or 2 gallons of 2.5% Mo liquid product) in 20 to 40 gallons of water and spray uniformly over each acre. Apply before transplanting and disc into the soil. Because sodium molybdate is compatible with many herbicides used on tobacco,

Table 10. Nitrogen recommendations (lb/A), burley and dark tobacco.

N Levels	Soil Drainage Class	
	Well-Drained	Moderately Well-Drained
Low ¹	225 - 250	250 - 275
Medium ²	200 - 225	225 - 250
High ³	150 - 175	175 - 200

¹ Following tobacco or row crops.

² First-year tobacco following a grass or grass-legume sod.

³ First-year tobacco following legume sod or legume cover crop.

it can be applied with herbicides normally applied as water-based sprays. Combining the two chemicals can result in application cost savings because only one trip over the field is necessary. It is recommended that not more than 2 lb of sodium molybdate (12.8 oz of molybdenum) per acre be applied during a five-year period.

Setter Water—Use 0.25 to 0.50 lb sodium molybdate (1.6 to 3.2 oz of molybdenum) per acre. If dry sodium molybdate is used, divide the total recommended amount (0.25 to 0.50 lb/A) equally among the number of 52-gallon barrels of water used per acre. For example, if 8 barrels of water per acre are used, add one-eighth (0.2 to 0.4 oz sodium molybdate) of the total recommended amount to each barrel, and fill the barrel with water. Adding the dry material before filling the barrel will aid in dissolving and mixing. If a 2.5% liquid source of molybdenum is used with 8 barrels of setter water per acre, add 0.50 pint to 1 pint (1 to 2 cups) of the liquid product per barrel before filling the barrel with water.

Conventional Plant Beds

Site Selection—A gentle slope facing south or southeast is preferable. The soil should be high in fertility with a soil pH of 6.0 to 6.6. Rotate with sod crops after one or two years, and use a green manure crop between consecutive years.

Fertilizer Rates—Salt damage is often the greatest problem in establishing tobacco seedlings in the plant bed and is commonly caused by excessive use of plant bed fertilizer. Recommendations for plant beds are shown in Table 12.

Incorporate needed fertilizer into the surface 1 to 2 inches of soil before seeding. If additional N is needed later by growing transplants, topdress with 0.5 to 1.0 lb actual N for each 100 sq yd. Nitrogen materials should be evenly broadcast when plants are dry, followed immediately by enough irrigation to remove fertilizer residues from the plant leaves.

Float Plant Beds

Source—Choose a water-soluble fertilizer that has less P₂O₅ than N or K₂O (i.e., 20-10-20, 15-5-15, etc.). Nitrate should be the major source of nitrogen with little or no urea to avoid problems with plant toxicity (see AGR-163).

Rates and Timing—The initial application should be made to bring the N concentration in the water to approximately 100 parts per million (ppm). For a fertilizer that has 20% N (20-10-20), this would be 4.2 lb of fertilizer for 1000 gallons of float water. For 15% N, this would be 5.6 lb of fertilizer per 1000 gallons of float water. For good growth and reduced disease susceptibility, N should be maintained in the range of

75 to 100 ppm. Inexpensive conductivity meters can be used to monitor fertility levels in the float bed (see AGR-174). The initial application can be made at seeding time. However, waiting 4 to 7 days after seeding may reduce the chance of salt injury to young seedlings. When fertilizer is added after seeding, care should be exercised to ensure adequate mixing and even distribution of the fertilizer in the float water.

Category	Burley and Dark		Burley		Dark	
	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed	Test Result: K	K ₂ O Needed
Very high	>80	0	>450	0	>450	0
High	73 - 79	30	424 - 449	30	398 - 450	30
	71 - 72	40	417 - 423	40	383 - 397	40
	68 - 70	50	409 - 416	50	368 - 382	50
	66 - 67	60	402 - 408	60	353 - 367	60
	64 - 65	70	394 - 401	70	338 - 352	70
	62 - 63	80	387 - 393	80	323 - 337	80
	58 - 61	90	379 - 386	90	308 - 322	90
			372 - 378	100	296 - 307	100
			364 - 371	110		
			357 - 363	120		
			349 - 356	130		
			342 - 348	140		
			334 - 341	150		
		327 - 333	160			
		319 - 326	170			
		312 - 318	180			
		304 - 311	190			
Medium	54 - 57	100	296 - 303	200	286 - 295	110
	50 - 53	110	286 - 295	210	276 - 285	120
	46 - 49	120	276 - 285	220	266 - 275	130
	41 - 45	130	266 - 275	230	256 - 265	140
	37 - 40	140	256 - 265	240	246 - 255	150
	33 - 36	150	246 - 255	250	236 - 245	160
	29 - 32	160	236 - 245	260	226 - 235	170
			226 - 235	270	216 - 225	180
216 - 225			280	206 - 215	190	
206 - 215			290			
Low	25 - 28	170	195 - 205	300	195 - 205	200
	22 - 24	180	184 - 194	310	184 - 194	210
	18 - 21	190	173 - 183	320	173 - 183	220
	14 - 17	200	162 - 172	330	162 - 172	230
	11 - 13	210	151 - 161	340	151 - 161	240
	7 - 10	220	140 - 150	350	140 - 150	250
			129 - 139	360	129 - 139	260
			118 - 128	370	118 - 128	270
			107 - 117	380	107 - 117	280
			96 - 106	390	96 - 106	290
Very low	<7	230	<96	400	<96	300

pH Level	Lb/Bed Aglime Needed	N Level	Lb/Bed N Needed	P Level	Lb/Bed P ₂ O ₅ Needed	K Level	Lb/Bed K ₂ O Needed
6.0 +	0		5	>80	0	>300	0
<6.0	100 - 150			<80	5 - 10	<300	5
Note: Fertilization without soil test: 5 lb N, 5-10 lb P ₂ O ₅ , 5 lb K ₂ O per bed.							
¹ A bed measures 900 square feet.							

CORN

Lime

If water pH is below 6.2, see page 5, and use the rate appropriate to attain a target pH of 6.4.

Nitrogen

Also see the “Fertility Management” chapter in University of Kentucky Extension publication ID-139, *A Comprehensive Guide to Corn Management*.

Winter Legume Cover Crops—A winter legume cover crop can provide a substantial amount of nitrogen for corn with either no-tillage or conventional tillage.

Research conducted by the University of Kentucky on no-tillage corn indicates that some legume cover crops can provide yield advantages beyond that provided by fertilizer nitrogen. Hairy vetch performed better than crimson clover or big flower vetch. The dense mulch resulting from a killed legume cover crop conserves soil water, aids in weed control, and helps to control soil erosion.

Three important factors should be considered when using a legume cover crop:

1. The amount of nitrogen provided will depend on the amount of growth the legume makes before it is chemically killed or plowed under. However, corn planting should not be delayed later than mid-May, particularly on well-drained soils.
2. A cover crop, legume or non-legume, can deplete soil water during a dry spring, resulting in decreased germination and seedling growth of corn.
3. Some vetch seeds are hard and can remain in the soil for one or more years before germinating. The result can be volunteer vetch in small grains grown in rotation with corn unless the vetch is killed with herbicides in the early spring.

Placement—Small amounts of N plus K₂O can be applied in the row, but if more than 15 lb/A of N plus K₂O is banded, it should be banded at least 2 inches below the soil surface and 2 inches to the side of the seedrow center. No more than 100 lb/A of N plus K₂O should be banded near the row. Fertilizer for banding near or in the row should not contain urea.

Adjustments to Nitrogen Recommendations

Irrigation—The nitrogen rate on irrigated corn should be increased to 175 to 200 lb N/A due to increased risk of depletion of available N from crop uptake, leaching, and denitrification.

Sidedressed N—On moderately well-drained to poorly drained soil, rates of nitrogen can be decreased by 35 lb/A if as much as two-thirds of the N is applied 4 to 6 weeks after planting.

Table 13. Recommended application of nitrogen (lb N/A), corn.

	Tillage ¹	Soil Drainage Class ²		
		Well-Drained	Moderately Well-Drained	Poorly Drained
Corn, sorghum, soybean, small grain, fallow	Intensive	100 - 140	140 - 175	175 - 200
	Conservation	125 - 165	165 - 200	
Grass, grass-legume sod (4 years or less), winter annual legume cover	Intensive	75 - 115	115 - 150	150 - 175
	Conservation	100 - 140	140 - 175	
Grass, grass-legume sod (5 years or more)	Intensive	50 - 90	90 - 125	125 - 150
	Conservation	75 - 115	115 - 150	

¹ Intensive tillage has less than 30% residue cover, and conservation tillage has more than 30% residue cover on the soil at planting.
² Soil drainage class examples are given on page 2.

Table 14. Phosphate and potash recommendations (lb/A), corn.

Category	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed
Very high			>420	0
High	>60	0	355 - 420	0
			336 - 354	0
			318 - 335	0
			301 - 317	0
Medium	46 - 60	30	282 - 300	30
			264 - 281	30
			242 - 263	30
			226 - 241	40
			209 - 225	50
			191 - 208	60
Low	23 - 27	80	173 - 190	70
			155 - 172	80
			136 - 154	90
			118 - 135	100
			100 - 117	110
			6 - 8	120
Very low	1 - 5	200	<100	120

Conservation Tillage—On moderately well-drained to poorly drained soils, the risk of denitrification loss is great from N applied at or near planting on conservation-till corn. Alternative practices may include:

1. sidedressed N application as noted above;
2. use of maximum rate in appropriate recommended range at planting; or
3. use of a nitrification inhibitor at planting with N sources that include urea, N-solutions, or anhydrous ammonia with minimum rate of N within the appropriate recommended range (see AGR-185 for additional information).

Surface-Applied Urea—Volatilization losses of N from urea-based products can be significant when they are surface-applied after May 1. Alternative practices include:

1. irrigation or incorporation within two days after application;
2. use of a urease inhibitor;
3. use of maximum N fertilization rate in appropriate recommended range (see AGR-185 for additional information).

Tiled Soils—Poorly drained soils that have been tile drained should be considered as moderately well-drained.

Zinc

Where zinc deficiency of corn has previously occurred or is suspected, a zinc soil test is helpful in determining if zinc should be applied. The following table gives the soil test zinc levels at various soil pH and soil test P levels below which a response to zinc fertilization is likely to occur. However, many other factors including weather conditions and cool soil temperatures affect soil zinc availability to corn, making it difficult to predict a response to added zinc for a specific growing season. Zinc can be a broadcast or banded treatment. Broadcast zinc fertilizer should raise the Zn soil test to acceptable levels for several years.

Table 15. Zinc recommendations¹ (lb/A), corn.

Test Result: P	Soil Water pH																	Zn Needed	
	6.0	6.1	6.2	6.3	6.4	6.5	6.6	6.7	6.8	6.9	7.0	7.1	7.2	7.3	7.4	7.5	7.6	Broadcast	Banded
25	0.5	0.7	0.9	1.0	1.2	1.4	1.6	1.7	1.9	2.1	2.3	2.4	2.6	2.8	3.0	3.1	3.3	10-15	2-3
50	1.1	1.2	1.4	1.6	1.8	1.9	2.1	2.3	2.5	2.6	2.8	3.0	3.2	3.3	3.5	3.7	3.9		
75	1.4	1.5	1.7	1.9	2.1	2.2	2.4	2.6	2.8	3.0	3.1	3.3	3.5	3.6	3.8	4.0	4.2		
100	1.6	1.8	1.9	2.1	2.3	2.5	2.6	2.8	3.0	3.2	3.4	3.5	3.7	3.9	4.1	4.2	4.4		
150	1.9	2.1	2.3	2.4	2.6	2.8	3.0	3.1	3.3	3.5	3.7	3.8	4.0	4.2	4.4	4.5	4.7		
200	2.1	2.3	2.5	2.7	2.8	3.0	3.2	3.4	3.5	3.7	3.9	4.1	4.2	4.4	4.6	4.8	4.9		
250	2.3	2.5	2.7	2.8	3.0	3.2	3.4	3.5	3.7	3.9	4.1	4.2	4.4	4.6	4.8	4.9	5.1		
300	2.4	2.6	2.8	3.0	3.2	3.3	3.5	3.7	3.9	4.0	4.2	4.4	4.6	4.7	4.9	5.1	5.3		
350	2.6	2.7	2.9	3.1	3.3	3.4	3.6	3.8	4.0	4.2	4.3	4.5	4.7	4.9	5.0	5.2	5.4		
400	2.7	2.9	3.0	3.2	3.4	3.6	3.7	3.9	4.1	4.3	4.4	4.6	4.8	5.0	5.1	5.3	5.5		
450	2.8	2.9	3.1	3.3	3.5	3.6	3.8	4.0	4.2	4.3	4.5	4.7	4.9	5.1	5.2	5.4	5.6		
500	2.8	3.0	3.2	3.4	3.6	3.7	3.9	4.1	4.3	4.4	4.6	4.8	5.0	5.1	5.3	5.5	5.7		

¹ Zinc and phosphorus levels shown are from soil extraction by the Mehlich III procedure. To determine if zinc is needed, find the appropriate soil test P level in the left column and read across the table to the appropriate soil pH level. If soil test zinc is less than that shown for the appropriate soil test P level and pH, apply fertilizer zinc as recommended in the table.

SOYBEAN

Lime

If water pH is below 6.2, see page 5, and use the rate appropriate to attain a target pH of 6.4.

Nitrogen

No nitrogen is recommended for well-nodulated soybean.

Double Cropping of Small Grains and Soybean—The phosphate recommendation should be taken from small grains, and the potash recommendation should be taken from soybean. This recommendation can be applied in the fall before seeding the small grain.

Inoculation

Soybean should be inoculated when planted in fields where soybean has not been grown in the past three to five years or where previously grown soybean had few nodules. If inoculation is necessary, the inoculant should be applied to the seed or in the row at planting. Delays in planting inoculated seed often result in poor nodulation. Numbers of live rhizobia in inoculum decrease rapidly under dry conditions, with exposure

to sunlight and high temperatures, or when dry-packaged with sodium molybdate and fungicides.

Molybdenum

If soils are limed to maintain pH values at 6.2 or above, aluminum and manganese toxicities and molybdenum deficiency usually do not occur in soybean. However, in soils with pH values below 6.2 at seeding time, molybdenum application to soybean is recommended. Apply 1 to 2 oz of sodium molybdate (0.4 to 0.8 oz of elemental molybdenum) per acre as a seed treatment. This is a satisfactory method of applying a small amount of molybdenum where no seed inoculant is needed. Where soybean seed is to be inoculated, including sodium molybdate with the inoculum may seriously lower the numbers of live rhizobia if seed is not planted immediately. If both inoculum and molybdenum are needed, apply inoculum to the seed, and broadcast the molybdenum on the soil. For each acre, dissolve 1 lb sodium molybdate (6.4 oz molybdenum) in 20 to 40 gallons of water and spray uniformly ahead of final seedbed preparation. Not more than 2 lb sodium molybdate (13 oz molybdenum) per acre should be used during any five-year period. Use of molybdenum should not be substituted for a good liming program.

Manganese

Foliar applications of manganese to manganese-deficient soybean have been superior to soil applications made at planting. Foliar manganese spray is recommended in two forms: (1) as chelated manganese at rates recommended by the manufacturer on the label and (2) as manganese sulfate at a rate of 1.0 to 1.5 lb of actual manganese per acre in 20 to 25 gallons of water when the soybean is 10 inches high or higher. The symptom of Mn deficiency is interveinal chlorosis of the emerging (youngest) leaves. Tissue analysis can also be used to evaluate the Mn status of the plant. The sufficiency range is 15 to 200 parts per million (ppm) Mn in the uppermost mature trifoliolate leaves (petiole discarded). See *Soil Science News and Views*, volume 10, number 8, "Manganese Fertility of Soybean" for additional information.

Table 16. Phosphate and potash recommendations (lb/A), soybean.

Category	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed
High	>60	0	>300	0
Medium	40 - 60	30	242 - 300	30
	34 - 39	40	226 - 241	40
	28 - 33	50	209 - 225 191 - 208	50 60
Low	22 - 27	60	173 - 190	70
	16 - 21	70	155 - 172	80
	11 - 15	80	136 - 154	90
	9 - 10	90	118 - 135	100
	7 - 8 6	100 110	100 - 117	110
Very low	1 - 5	120	82 - 99	120
			64 - 81	130
			46 - 63	140
			<46	150

SMALL GRAINS

(BARLEY, OATS, RYE, WHEAT, AND TRITICALE)

Lime

If water pH is below 6.2, see page 5, and use the rate appropriate to attain a target pH of 6.4.

Nitrogen

Fall Application—Only enough N to provide for good ground cover and to stimulate tillering is necessary. Seedlings following tobacco, soybean, or well-fertilized corn will likely have enough carryover N for fall growth. For optimal fertilizer N efficiency, the total fall application should not exceed 40 lb N/A for seedlings in fields with insufficient N carryover. Fall-applied N will be of little benefit where little fall growth is expected.

Spring Application—Application from late February to early April is the most effective. Where excessive rainfall occurs in late winter or early spring, causing excessive wetness on less than well-drained soils, split applications of spring-applied N may be justified.

Sources—Experimental results have shown little difference among nitrogen materials commonly used to supply supplemental N to small grains.

Small Grains for Grazing—Total forage production from small grains can be increased by splitting nitrogen applications between fall and spring. For fall grazing, apply 50 to 60 lb N/A at seeding. A late winter or early spring topdressing of 30 to 50 lb N/A will stimulate early growth for additional grazing.

Intensively Managed Wheat—When managed for high yields (70 to 100 bu/A), wheat should receive higher rates of N in the spring. If spring N is split into two applications (early to mid-February and mid- to late March), yields will be 3 to 5 bu/A higher than if all N is applied in a single application in mid- to late March. The February application should be made at green-up, and the March application should be made at Feekes growth stage 5 or 6 (just prior to or at jointing). Green-up may not occur until March in central and northern Kentucky.

Table 17. Spring nitrogen rates (lb N/A), small grains.

Seedbed	Lb N/A
Tilled	60 - 90
No-till	90 - 120

Table 18. Spring nitrogen rates (lb N/A), intensive wheat.

Applications	Feb	Mar	Total
Single	0	95	95
Split	30 - 60	75 - 45	105

Table 19. Phosphate and potash recommendations (lb/A), small grains.

Category	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed
High	>60	0	>300	0
Medium	48 - 60	30	213 - 300	30
	45 - 47	40	187 - 212	40
	41 - 44	50		
	38 - 40	60		
	34 - 37	70		
	31 - 33	80		
Low	24 - 30	90	159 - 186	50
	17 - 23	100	132 - 158	60
	10 - 16	110	104 - 131	70
Very low	<10	120	<104	80

Double Cropping of Small Grains and Soybean—The phosphate recommendation should be taken from small grains, and the potash recommendation should be taken from soybean. This recommendation can be applied in the fall before seeding the small grain.

CORN SILAGE AND SMALL GRAIN HAY/SILAGE

Because the entire plant is removed, silage/hay production results in much different nutrient removal than when growing the same crops for grain. Additionally, because of land area limitations, silage is usually produced for several consecutive years without rotation. The high nutrient removal and high recommended N application rates mandate that soils under continuous silage production be sampled every year to assure appropriate pH and nutrient levels. Much of the required nutrients can be supplied via appropriate use of animal manures from either on or off-farm confined animal enterprises. For additional information see University of Kentucky Extension publications AGR-17, *Double Crop Silage Production*, and AGR-165, *The Agronomics of Manure Use for Crop Production*.

If water pH is below 6.2, see page 5, and use the rate appropriate to attain a target pH of 6.4.

Nitrogen and Zinc

Fall-applied nitrogen (N) for the small grain is usually not needed, but most producers apply some if they use 18-46-0 or 11-52-0 for fall phosphorus applications. An additional top-dressing of N should be made to the small grain in late winter or early spring just before growth begins. Total N application to

the small grain should not exceed 110 lb N/A. There is no zinc (Zn) recommendation for small grains grown in Kentucky.

Nitrogen and Zn application for corn silage is no different from that for corn grown for grain. See Table 13 (page 9) and Table 15 (page 10) for specific recommendations.

Table 20. Phosphate and potash recommendations (lb/A) for corn and small grain silage (hay) production.

Category	Test Result: P	P ₂ O ₅ Needed			Test Result: K	K ₂ O Needed ²	
		Corn	Small Grain	Dbl-Crop Silage ¹		Corn	Small Grain
Very high					>420	0	0
High	>60	0	0	0	355 - 420	30	40
					336 - 354	40	40
					318 - 335	50	40
					301 - 317	60	40
Medium	46 - 60 41 - 45 37 - 40 33 - 36 28 - 32	30	30	60	282 - 300	70	70
		40	40	80	264 - 281	80	70
		50	50	100	242 - 263	90	70
		60	60	120	226 - 241	100	70
		70	70	140	209 - 225	110	70
					191 - 208	120	70
Low	23 - 27 19 - 22 14 - 18 9 - 13 6 - 8	80	80	160	173 - 190	130	80
		90	90	180	155 - 172	140	80
		100	100	200	136 - 154	150	90
		110	110	220	118 - 135	160	110
		120	120	240	100 - 117	170	120
Very low	1 - 5	200	120	320	<100	180	120

¹ For double-crop silage, all of the P can be applied in the fall before the small grain (as long as total fall N does NOT exceed 30 lb N/A).

² Potassium should be applied according to the recommendation before each of the crops are planted to minimize luxury consumption of K (see page 16 for additional information).

GRAIN SORGHUM

Lime

If water pH is below 6.2, see page 5, and use the rate appropriate to attain a target pH of 6.4.

Placement—Banded fertilizer should be placed 2 inches below the soil surface and 2 inches to the side of the row with a maximum of 40 lb/A N plus K₂O. Additional fertilizer should be broadcast.

Table 21. Recommended application of nitrogen (lb N/A), grain sorghum.

Previous Crop	Lb N/A Needed ¹
Corn, sorghum, soybean, small grain, fallow, set-aside	100 - 125 ²
Grass, grass-legume sod (4 years or less)	75 - 100
Grass, grass-legume sod (5 years or more)	50 - 75

¹ Recommended rates are for moderately well-drained soils which comprise the basis of current field data. See page 2 for soil drainage class.

² **Note:** Rates of nitrogen fertilization can be decreased 25 lb/A if two-thirds or more of the nitrogen is applied 4 to 6 weeks after planting.

Table 22. Phosphate and potash recommendations (lb/A), grain sorghum.

Category	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed
High	>60	0	>300	0
Medium	46 - 60	30	242 - 300	30
	41 - 45	40	226 - 241	40
	37 - 40	50	209 - 225	50
	33 - 36	60	191 - 208	60
	28 - 32	70		
Low	23 - 27	80	173 - 190	70
	19 - 22	90	155 - 172	80
	14 - 18	100	136 - 154	90
	9 - 13	110	118 - 135	100
	6 - 8	120	100 - 117	110
Very low	<6	200	<100	120

CANOLA

Lime

If water pH is below 6.2, see page 5, and use the rate appropriate to attain a target pH of 6.4.

Nitrogen

Fall Application—Only enough N to provide for good ground cover and good root growth is necessary to aid in winter survival. Seedings following tobacco, soybean, or well-fertilized corn will likely have enough carryover N for fall growth. For optimum fertilizer N efficiency, the total fall application should not exceed 30 lb N/A for seeding in fields with insufficient N carryover.

Spring Rates

Previous Crop	Lb N/A
Corn, small grains, soybean, fallow, set-aside	120
Grass-legume or legume sod	90

Spring Applications—Applications between late February and late March are the most effective time. This coincides with spring green-up. All N should be applied before stem elongation.

Table 23. Phosphate and potash recommendations (lb/A), canola.

Category	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed
High	>60	0	>300	0
Medium	48 - 60	30	213 - 300 187 - 212	30
	45 - 47	40		40
	41 - 44	50		
	38 - 40	60		
	34 - 37	70		
	31 - 33	80		
Low	24 - 30	90	159 - 186	50
	17 - 23	100	132 - 158	60
	10 - 16	110	104 - 131	70
Very low	<10	120	<104	80

Double Cropping of Canola and Soybean—The phosphate recommendations should be taken from canola, and the potash recommendation should be taken from soybean. This recommendation can be applied in the fall before seeding the canola.

HAY AND PASTURES

NEW SEEDINGS

Lime

See page 5, and use the rate appropriate to attain a target pH of 6.4. For long-term production of alfalfa and alfalfa grass, it is important to raise pH and maintain it in a range between 6.5 and 7.0. (See page 5 for the rate of lime needed to attain a target pH of 6.8 when the water pH is below 6.6.)

Nitrogen

Apply 0 to 30 lb N/A at seeding for legumes or grass-legume mixtures and 0 to 50 lb N/A for grass-only seedings. If the field has a history of high N application, omit N at seeding.

Surface Mine Reclamation

See page 4 for more details.

Molybdenum

If soils are limed to maintain pH values at 6.2 or above, aluminum and manganese toxicities and molybdenum deficiency usually do not occur in forage legumes. However, for soils having pH values below 6.2 at seeding time, molybdenum application to forage legumes is recommended. Apply at the rate of 1 lb of sodium molybdate (6.4 oz of molybdenum) per acre. Dissolve this amount of molybdate in 20 to 40 gallons of water, and spray uniformly over each acre. Apply before planting, and disc into the soil unless using no-till establishment. It is recommended that not more than 2 lb of sodium molybdate (12.8 oz of molybdenum) per acre be used during a five-year period.

Inoculation

Appropriate good-quality inoculant should be applied to legume seed or in the row at planting. Delays in planting inoculated seed may result in poor root nodulation. Numbers of live rhizobia decrease rapidly under dry conditions, with exposure to sunlight and high temperatures, or when dry-packaged with sodium molybdate and fungicides.

Table 24. Phosphate and potash recommendations (lb/A), hay and pasture, new seedings.

Category	Alfalfa, Alfalfa-Grass, Clover, Clover-Grass		Alfalfa, Alfalfa-Grass		Clover, Clover-Grass		Cool-Season Grasses			
	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed	Test Result: K	K ₂ O Needed	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed
High	>60	0	>450	0	>300	0	>60	0	>300	0
			394 - 450	60						
			363 - 393	90						
			338 - 362	100						
			313 - 337	110						
297 - 312	120									
Medium	51 - 60 48 - 50 46 - 47 43 - 45 40 - 42 37 - 39 35 - 36 32 - 34 28 - 31	30 40 50 60 70 80 90 100 110	291 - 296	130	266 - 300	30	48 - 60	30	213 - 300	30
			285 - 290	140	256 - 265	40	45 - 47	40	187 - 212	40
			279 - 284	150	246 - 255	50	41 - 44	50		
			272 - 278	160	236 - 245	60	38 - 40	60		
			266 - 271	170	226 - 235	70	34 - 37	70		
			260 - 265	180	216 - 225	80	31 - 33	80		
			254 - 259	190	206 - 215	90				
			247 - 253	200	191 - 205	100				
			241 - 246	210						
			235 - 240	220						
			229 - 234	230						
			222 - 228	240						
			216 - 221	250						
			210 - 215	260						
			204 - 209	270						
Low	23 - 27 19 - 22 14 - 18 9 - 13	120 130 140 150	194 - 203	280	173 - 190	110	24 - 30	90	159 - 186	50
			180 - 193	290	155 - 172	120	17 - 23	100	132 - 158	60
			166 - 179	300	136 - 154	130	10 - 16	110	104 - 131	70
			152 - 165	310	118 - 135	140				
			139 - 151	320	100 - 117	150				
			125 - 138	330						
111 - 124	340									
97 - 110	350									
Very low	<9	160	<97	360	<100	160	<10	120	<104	80

HAY AND PASTURES

ESTABLISHED STANDS OF LEGUMES AND GRASS-LEGUME MIXTURES

Topdressing Legumes and Grass-Legume Mixtures

Top production from hay and pasture fields can best be obtained if soil test levels are in the range of 50 to 60 P and 270 to 300 K and are maintained at those levels. If initial soil test levels are below these ranges, they can be raised more quickly by heavy nutrient applications at seeding and then maintained by annual topdressings as outlined in the following tables. Another alternative is to raise soil test levels gradually over a longer period of years by foregoing a heavy nutrient application at seeding but increasing topdressing rates each year over the rates shown. An initial soil test, followed by periodic soil tests, will indicate changes in field fertility levels that are taking place.

Lime

See page 5, and use the table to determine the rate appropriate to attain a target pH of 6.4. For alfalfa or alfalfa-grass, use the table on page 5 to find the rate needed for a target pH level of 6.8 when the water pH is below 6.6.

Nitrogen

Topdress applications of N are not recommended for legumes or legume-grass mixtures containing more than 25% legumes. If there is less than 25% legume in a legume-grass stand, topdress with N at rates shown for established grass stands.

Boron

For alfalfa production, apply 1.5 to 2.0 lb per acre of elemental boron (B) per acre every two years either as a borated fertilizer or as fertilizer borate. If boron-containing materials or wastes from a coal-fired power plant have been applied in successive years to established stands of alfalfa, the field should be tested for boron. If soil test B exceeds 2.0 lb per acre, additional B should not be applied.

Surface Mine Reclamation

See page 4 for more details.

Higher Yields

For alfalfa yields above 5 tons per acre and red clover yields above 3 tons per acre, fields should be soil sampled every year to monitor P and K levels.

Luxury Consumption of Potassium

Luxury consumption is a phenomenon that all alfalfa producers should be aware of and should try to avoid. Luxury consumption occurs when the alfalfa plant takes up more K than is needed for maximum yield. The additional K is removed with hay harvest and is not available for future cuttings. To minimize luxury consumption, K fertilizer should not be applied in the spring prior to the first cutting for existing stands. For new stands of alfalfa, K fertilizer should be thoroughly incorporated prior to planting.

Table 25. Phosphate and potash recommendations (lb/A), hay and pasture, annual topdressing.

Category	Alfalfa or Alfalfa-Grass				Clover or Clover-Grass			
	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed
High	>60	0	>450	0	>60	0	>300	0
			394 - 450	60				
			363 - 393	90				
			338 - 362	100				
			313 - 337	110				
			297 - 312	120				
Medium	46 - 60 41 - 45 37 - 40 33 - 36 28 - 32	30 40 50 60 70	291 - 296	130	41 - 60 36 - 40 31 - 35	30 40 50	271 - 300	30
			285 - 290	140			263 - 270	40
			279 - 284	150			255 - 262	50
			272 - 278	160			246 - 254	60
			266 - 271	170			238 - 245	70
			260 - 265	180			230 - 237	80
			254 - 259	190			221 - 229	90
			247 - 253	200			213 - 220	100
			241 - 246	210			205 - 212	110
			235 - 240	220				
			229 - 234	230				
			222 - 228	240				
			216 - 221	250				
			210 - 215	260				
204 - 209	270							
Low	23 - 27 19 - 22 14 - 18 9 - 13	80 90 100 110	194 - 203	280	27 - 30 23 - 26 19 - 22 16 - 18 12 - 15 8 - 11	60 70 80 90 100 110	191 - 204	120
			180 - 193	290			173 - 190	130
			166 - 179	300			155 - 172	140
			152 - 165	310			136 - 154	150
			139 - 151	320			118 - 135	160
			125 - 138	330			100 - 117	170
			111 - 124	340				
			97 - 110	350				
Very low	<9	120	<97	360	<8	120	<100	180

HAY AND PASTURES

RENOVATION OF GRASS WITH CLOVERS OR ANNUAL LESPEDEZA

Lime

If water pH is below 6.2, see the table on page 5 for the rate appropriate to attain a target pH of 6.4.

Nitrogen

One factor that is critical to legume establishment in established grass sods is grass competition with young legume seedlings. Use of N at renovation will stimulate grass growth and increase the likelihood of legume stand failure. However, if there is a need for increased grass production during the fall preceding spring legume renovation, a small amount of N (up to 50 lb N/A) can be topdressed between Aug. 1 and Aug. 15. Be sure any increased grass growth is grazed off before legume renovation.

Annual Topdressing

See recommendations for topdressing legume-grass mixtures.

Table 26. Phosphate and potash recommendations (lb/A), clover or annual lespedeza (renovation into established grass).

Category	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed
High	>60	0	>300	0
Medium	48 - 60	30	257 - 300	30
	45 - 47	40	244 - 256	40
	41 - 44	50	232 - 243	50
	38 - 40	60	219 - 231	60
	34 - 37	70	207 - 218	70
	31 - 33	80	187 - 206	80
Low	24 - 30	90	159 - 186	90
	17 - 23	100	132 - 158	100
	10 - 16	110	104 - 131	110
Very low	<10	120	<104	120

Inoculation

Appropriate good-quality inoculant should be applied to the legume seed or in the row at planting. Delays in planting inoculated seed often result in poor inoculation. Numbers of live rhizobia decrease rapidly under dry conditions, with exposure to sunlight and high temperatures, or when dry-packaged with sodium molybdate and fungicides.

Molybdenum

Molybdenum deficiency usually does not occur in forage legumes when soil pH is 6.2 or higher. However, with soil pH values below 6.2 at seeding, molybdenum application to forage legumes is recommended. Apply at a rate of 1 lb of sodium molybdate (6.4 oz actual Mo) per acre. Dissolve this amount of sodium molybdate in 20 to 40 gallons of water, and spray uniformly over each acre. Apply before planting, and disc into the soil unless performing no-till renovation. Not more than 2 lb of sodium molybdate (12.8 oz of Mo) per acre should be used during a five-year period.

HAY AND PASTURES

ESTABLISHED STANDS OF COOL-SEASON GRASSES

Topdressing

Cool-season grasses such as tall fescue, bluegrass, timothy, orchardgrass, and bromegrass grow best from early spring into early summer and again in the fall. If moisture and soil test levels of P and K are adequate (P >30, K >200), the use of nitrogen greatly stimulates their growth during these peak production periods. The use of nitrogen should depend on what is expected from the grass. Nitrogen fertilization can help increase total production and protein content and shift the seasonality of production. However, unless the increased grass and forage production is utilized, there will be no return to the added N expense.

Nitrogen Management

Increase Total Production—Research data for a period of years indicate that dry matter can be increased from about a ton per year with no nitrogen up to about 4 tons per year with up to 200 lb N per acre.

Shifting the Seasonality of Production—Timely nitrogen applications will increase the productivity of these grasses during particular seasons. Topdressing in late winter or early spring, just before growth begins, will increase growth so that grazing can begin about two weeks ahead of pastures receiving no nitrogen. This effect can be used to lower overwinter feed costs. Topdressing with N in late spring, following a grazedown or hay clipping, will increase growth that will carry over into the normally low summer production period. Topdressing with N following a grazedown or clipping in late summer will stimulate fall growth, which, if properly utilized, will extend grazing several weeks later into the fall and winter. This is a practical method to lower winter feed cost by keeping animals on pasture longer.

Table 27. When to topdress nitrogen.

Date	Lb N/A per Application ¹
Feb. 15 – Mar. 15	up to 100
May 1-15	up to 50
Aug. 1-15	up to 80

¹ Total amount of N to topdress should depend on how much additional production is needed. If a total of more than 100 lb of N per acre per year is to be used, it should be applied in split applications. Suggested dates and rates for topdressing with N are shown above.

Sources—Research in Kentucky has shown that during late winter and early spring there is little difference among the N sources commonly used for topdressing cool-season grasses. After early May, there is an increased risk that topdressed urea will not be as effective as other N sources. Average efficiency values for topdressed urea after early May ranged from 51 to 78% of that observed with ammonium nitrate, depending largely on the length of time between urea application and the next rainfall. When a urease inhibitor is used in conjunction with urea, the efficiency of urea becomes comparable to that of ammonium nitrate. For urea applied after early May without an inhibitor to be an economical substitute for ammonium nitrate, the cost per pound of nitrogen from urea probably should be 15 to 20% less than that of ammonium nitrate. Research indicates the efficiency of liquid nitrogen (urea-ammonium nitrate solutions) applied after early May is between that of urea and ammonium nitrate. A urease inhibitor can also be used in conjunction with liquid nitrogen.

Grass tetany (magnesium deficiency-hypomagnesaemia) problems with cattle are sometimes encountered on straight grass pastures, particularly with nursing cows where grass

Table 28. Phosphate and potash recommendations (lb/A), cool-season grasses, annual topdressing.

Category	Test Result: P	P ₂ O ₅ Needed	Test Result: K	Pasture: ¹ K ₂ O Needed	Hay: K ₂ O Needed
Very high			>420	0	0
High	>60	0	321 - 420	0	30
			301 - 320	0	40
Medium	46 - 60	30	267 - 300	30	50
			240 - 266	30	60
			213 - 239	30	70
			187 - 212	40	80
			28 - 32	70	
Low	23 - 27	80	159 - 186	50	90
			132 - 158	60	100
			104 - 131	70	110
			9 - 13	110	
			<9	120	
Very low	<9	120	<104	80	120

¹ If pasture is clipped and harvested for hay in the spring, K applications should be according to the hay recommendations.

pasture is the only source of feed. Tetany in such animals results from an imbalance of magnesium in their blood. Supplemental feeding of magnesium to nursing cows on such fields is recommended as a means of lowering tetany risk. Applying fertilizer containing magnesium is not recommended to offset potential grass tetany problems. There is little guarantee that the plant will take up the additional applied magnesium when soil test magnesium levels are adequate. See University of Kentucky Extension publication ASC-16, *Grass Tetany in Beef Cattle*, for detailed recommendations.

Grass Seed Production—For pure stands of tall fescue and bluegrass from which seed will be harvested, an additional topdressing of 60 to 70 lb N/A around Dec. 1, after grazedown, will increase seed yields the following year.

Lime

See page 5, and use the rate appropriate to attain a target pH of 6.4.

Surface Mine Reclamation

See page 4 for more details.

HAY AND PASTURES

WARM-SEASON FORAGES

Sudangrass, Millets, Sorghum-Sudangrass Hybrids

This category includes annual warm-season grasses. Consult University of Kentucky Extension publication AGR-88, *Producing Summer Annual Grasses for Emergency or Supplemental Forage*, for management details.

The soil pH should be maintained between 5.8 and 6.4. If establishing a stand and the water pH is less than 6.2, add lime to attain a target pH of 6.4 (see page 5). If the stand is already established, apply lime to reach a target pH of 6.4 when the water pH is less than 5.8. Apply 60 to 100 lb N/A at seeding plus 40 to 60 lb N/A topdressed after first and second grazedowns or hay clippings. Apply phosphate and potash according to soil test results (Table 29).

Bermudagrass

Bermudagrass is a perennial warm-season grass. Consult University of Kentucky Extension publication AGR-48, *Bermudagrass: A Summer Forage in Kentucky*, for management details.

The pH should be maintained between 5.8 and 6.4. If a stand is being established and the water pH is less than 6.2, then add lime to attain a target pH of 6.4 (see page 5). If the stand is already established, apply lime to reach a target pH of 6.4 when water pH is less than 5.8. For establishing new stands, apply 30 to 60 lb N/A one month after seeding or sprigging. As ground cover is attained, an additional 30 to 50 lb applied around Aug. 15 can be beneficial in stimulating first-year growth. The total amount of N used should be based on the amount of forage needed and whether it is grazed or cut for hay. Nitrogen should be applied in split applications for best results. Apply 60 to 100 lb N/A in the spring when green-up begins. A nitrogen application near 100 lb N/A is needed for top hay production. Apply an additional 60 to 100 lb N/A after each hay cutting or 40 to 60 lb N/A after each grazedown. The last N application should be made by mid-August. Apply phosphate and potash according to soil test results. Table 30 summarizes nitrogen, phosphate, and potash rate recommendations.

Table 29. Phosphate and potash recommendations (lb/A), annual warm-season crops.

Category	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed
High	>60	0	>300	0
Medium	48 - 60	30	271 - 300	30
	45 - 47	40	263 - 270	40
	41 - 44	50	255 - 262	50
	38 - 40	60	246 - 254	60
	34 - 37	70	238 - 245	70
	31 - 33	80	230 - 237	80
			221 - 229	90
		213 - 220	100	
		205 - 212	110	
Low	24 - 30	90	191 - 204	120
	17 - 23	100	173 - 190	130
	10 - 16	110	155 - 172	140
			136 - 154	150
			118 - 135	160
		100 - 117	170	
Very low	<10	120	<100	180

Table 30. Annual nitrogen, phosphate, potash applications (lb/A), bermudagrasses.

Soil Test Level	Annual Application					
	Pasture			Hay		
	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O
High >60 P 300 K	120 - 240	0	0	120 - 400	0	0
Medium 60 - 30P 300 - 200K	120 - 240	30 - 65	30 - 90	120 - 400	30 - 95	30 - 180
Low <30P 200K ¹	120 - 240	65 - 120	90 - 180	120 - 400	95 - 120	180 - 360

¹ The maximum P₂O₅ or K₂O rate shown should be used if P is less than 10 or K is less than 100.

Native Warm-Season Grasses

Native warm-season perennial grasses can be used for forage. These plant species include bluestems, switchgrass, indiagrass, side oats grama, and eastern gamagrass. See University of Kentucky Extension publication AGR-145, *Native Warm-Season Perennial Grasses for Forage in Kentucky*, for detailed information on management.

Do not apply nitrogen at the time of seeding to avoid growth of other plants that can compete with the young seedlings. An application of 40 to 60 lb N/A can be applied in July of the seeding year to aid establishment. Apply 40 to 60 lb N/A

for established stands after the grass begins to green up in the spring. If N is applied too early, it will promote the growth of cool-season plant species that will compete with the warm-season grass. An additional 40 to 60 lb N/A can be applied after harvest in June or July for increased yields. Apply phosphate and potash according to Table 31.

Table 31. Annual phosphate and potash applications (lb/A), establishment or forage use of native warm-season grasses.

Soil Test Level	P ₂ O ₅	K ₂ O
High >60 P 300 K	0	0
Medium 60 - 30 P 300 - 200 K	30 - 40	30 - 50
Low¹ <30 P 200 K	40 - 80	50 - 100

¹ The maximum rates should be used if P is less than 6 lb/A or K is less than 90 lb/A.

HAY AND PASTURES

HORSE PASTURES

General Fertility Requirements

As with all pastures, soil samples should be taken from horse pastures every three years. Then pastures should be limed and fertilized with phosphorus (P) and potassium (K) to soil test recommendations to maintain vigorous growth of desired grasses and legumes and provide competition against undesirable weeds (Table 34). Pasture fertilization is not a sound approach to alleviate macro- and micronutrient deficiencies of livestock. These problems may be better addressed by more direct actions, such as feeding of mineral supplements.

Topdressing Nitrogen on Cool-Season Grass Pastures

The need to topdress nitrogen (N) on horse pastures in Kentucky is often less than that for cattle pastures because the goal with mature horses is weight/condition maintenance, whereas the goal on most beef cattle pastures is maximum growing animal weight gain. The guidelines shown below base N topdressing rates on stocking rate (horses per acre) and the percentage of clover (pasture containing more than 25% clover will supply most N needs of the pasture grass) in the stand (Tables 32 and 33).

Cool-season grasses, such as Kentucky bluegrass, orchardgrass, timothy, and tall fescue, grow most vigorously from early spring into early summer and then again in the fall into early winter. When soil moisture, pH (>6.0), and soil test levels of P and K are adequate (P >30, K >200), fertilizer N will stimulate cool-season grass growth during these peak production periods. The use of N should depend on what is expected from the pasture. Nitrogen fertilization can help increase total production and protein content, extend spring grass growth into the early summer, and extend fall pasture production into early winter. However, unless the increased pasture yield is utilized, there is no return to the added N expense, and there are several disadvantages. Stimulating grass growth in the spring can be helpful to maintaining a high stocking rate, but frequent clipping will be required to maintain pasture quality in a lightly stocked horse pasture. Late spring N application may have the unintended result of promoting the growth of

summer weeds such as crabgrass, yellow foxtail, nimblewill, and ragweed. In addition, there is an extra caution when broodmares are on pastures that contain tall fescue. In these pastures, spring N has been known to increase levels of ergot alkaloids, leading to an increased likelihood of fescue toxicity symptoms (prolonged gestation, difficult birth, and lack of milk production). Fortunately, this is only an issue with broodmares since tall fescue toxicity symptoms are not known to occur in other classes of horses.

Kentucky N fertilizer recommendations for cool-season grass horse pastures are shown in Tables 32 and 33. At low stocking rates, on soils with good productivity (>2 acres per mature horse), N should only be topdressed during the fall. Fall applications stimulate tillering of individual grass plants, producing a denser grass stand, which will suppress weed growth. Fall applications also lengthen the period of active photosynthesis, which promotes root growth and winter survival. If the primary goal is a more dense grass stand, then one late fall application between late October and early November is sufficient. If fall pasture growth is important, then also apply N between mid-August and mid-September. For suggested dates and N application rates for horse pastures managed at low stocking rate, see Table 32.

At higher stocking rates, on soils with good productivity (<2 acres per mature horse), fertilizer N may be topdressed on cool-season grasses throughout much of the growing season (Table 33). A late winter application will stimulate an early spring growth flush. An N application in May will help extend the pasture into the early summer, and a late summer application in August will stimulate cool-season pasture grass production in the fall and early winter. The major limitation to summer N fertilization is its stimulation of the growth of summer weeds. Excessive N applications may merely result in wasted forage. For suggested dates and N application rates for high stocking rate horse pastures, see Table 33.

The above-stated stocking rates are estimates for Kentucky soils of average productivity. The highest productivity soils in Kentucky will support one mature horse on 2 acres or less, while those with the lowest productivity (often with significant slopes) require 10 or more acres per horse. It is important to determine the recom-

Table 32. Topdressing nitrogen (lb/A) on cool-season horse pastures when maintained at low stocking rates (more than 2 acres/horse).

Date	N per Application ¹
Aug. 15 – Sept. 15	30 - 40
Oct. 15 – Nov. 15	30 - 60

¹ Total amount of N to topdress depends on desired result. No N is recommended if clover makes up more than 25% of the pasture. If primary goal is increased tillering for a denser grass stand winter, then one late fall application is sufficient. If fall pasture growth is important then also apply N in late August-early September. Suggested dates and rates for topdressing with N are shown above.

Table 33. Topdressing nitrogen (lb/A) on cool-season horse pastures when maintained at high stocking rates (less than 2 acres/horse).

Date	N per Application ¹
Feb. 15 – Mar. 15	up to 40 - 80
May 1-15	up to 30 - 40
Aug. 15-30	up to 40 - 80

¹ Total amount of N to topdress should depend on how much additional production is needed. Late spring N applications may have the unintended effect of stimulating unwanted summer weeds. Little or no N is recommended if clover makes up more than 25% of the pasture. Suggested dates and rates for topdressing with N are shown above.

Table 34. Phosphate and potash recommendations (lb/A) for cool-season grass horse pastures when applying annual fertilizer applications.

Category	Test Result: P	P ₂ O ₅ Needed	Test Result: K	K ₂ O Needed
Very high			>420	0
High	>60	0	321 - 420 301 - 320	0 0
Medium	46 - 60 41 - 45 37 - 40 33 - 36 28 - 32	30 40 50 60 70	267 - 300 240 - 266 213 - 239 187 - 212	30 30 30 40
Low	23 - 27 19 - 22 14 - 18 9 - 13	80 90 100 110	159 - 186 132 - 158 104 - 131	50 60 70
Very low	<9	120	<104	80

mended stocking rate for your soil type. See your county Extension agent or USDA-NRCS specialist for the average stocking rates associated with the soil types found in your area. Or go to the Web Soil Survey Web site for this information (www.websoilsurvey.nrcs.usda.gov). Often, stocking rate information in soil surveys is given in animal unit months (AUM). To convert animal unit months to recommended acres per horse, use the following formula: $(12 \div \text{AUM}) \times (\text{wt of horse} \div 1000) = \text{acres required per horse}$.

Sources of Nitrogen—Research in Kentucky has shown that during late fall, late winter, and early spring, there is little difference among the N sources commonly used for topdressing cool-season grasses. After May 1, there is an increased risk that topdressed urea will not be as effective as other N sources. Average efficiency values for urea N applied after early May ranged from 51 to 78% of that observed with ammonium nitrate, depend-

ing largely on the length of time between urea application and the next rainfall. When a urease inhibitor is used in conjunction with urea, the efficiency of urea becomes comparable to that of ammonium nitrate. For urea applied after early May and without an inhibitor, to be an economical substitute for ammonium nitrate, the cost per pound of nitrogen probably should be 15 to 20% less than that for ammonium nitrate-nitrogen. Research indicates that efficiency of liquid urea-ammonium nitrate (UAN) applied after early May is between that of urea and ammonium nitrate. A urease inhibitor can also be used in conjunction with UAN. To avoid burning symptoms on grass forages, remember to apply any granular N source when the leaf surface is dry. Also, avoid leaving piles of granular N in the field, either from misapplication or improper equipment operation because ingestion of significant quantities of N fertilizer can be toxic to horses.

CONSERVATION AND WILDLIFE LAND

Land can be managed to improve soil and water quality, enhance growth of native plant species, or provide habitat and food for wildlife. These types of land management are commonly referred to as conservation programs. The government can offer cost-sharing money to promote the management of land using specific conservation methods.

Riparian Buffer and Filter Strips

Riparian buffers and filter strips are land uses designed to improve water quality. Riparian buffers include land kept in an undisturbed condition for some distance away from the edge of streams. Filter strips are land containing grasses that filter out sediment in runoff from agricultural fields before entering surface water. Riparian buffers can include grasses, bushes, or trees. Filter strips can include warm- and cool-season grasses useful for trapping particles suspended in runoff water.

The pH should be maintained between 5.8 and 6.4. If water pH is below 5.8, apply lime to attain a target pH of 6.4 (see page 5). To ensure thick and vigorous growth, apply 40 to 60 lb N/A at seeding for cool-season grass or in July of the establishment year for warm-season grass. Apply phosphate and potash at seeding according to Table 35. Fertilizer or other sources of nutrients should not be applied on established stands. The management of this land can involve mowing, with no plant removal, so the nutrients in the grass are recycled back into the soil.

Table 35. Phosphate and potash recommendations (lb/A) for establishing riparian buffers and filter strips.

Soil Test Level	P ₂ O ₅	K ₂ O
High >60 P >300 K	0	0
Medium 60 - 30 P 300 - 200 K	30 - 40	30 - 50
Low¹ <30 P <200 K	40 - 80	50 - 100

¹ The maximum rate should be used if soil test P is less than 6 lb/A or soil test K is less than 90 lb/A.

Wildlife Food Plots

Land can be planted with a varied mixture of plant species to provide food for wildlife. An example of the various plant species are shown in Table 36. Land management depends on the composition of the plant species. Details on managing wildlife food plots can be found in University of Tennessee Extension publication PB 1743, *Growing and Managing Successful Wildlife Food Plots in the Mid-South*.

The soil pH should be maintained between 5.8 and 6.4. If the water pH is below 5.8, apply lime to attain a target pH of 6.4 (see page 5). Most of the seeding mixtures recommended for wildlife food plots contain one or more legume species that will add nitrogen to the soil. Therefore, only 0 to 30 lb N/A is required at seeding to establish the stand, and no N is applied to established stands. Be sure to inoculate the legume with species-specific inoculant prior to planting. If a legume is not in the seeding mixture, apply 40 to 60 lb N/A in the spring for cool-season grasses or after green-up occurs in warm-season grasses. Do not apply N at time of seeding warm-season grasses. Apply phosphate and potash according to Table 37.

Native Grassland Restoration

Native warm-season grasses can be planted in conservation land to restore native grassland conditions in Kentucky. Native warm-season grass species include bluestem, switch-

grass, indiangrass, side oats grama, and eastern gamagrass. Warm-season grasses offer good habitat for wildlife with their bunch-type growth characteristic providing open spaces for travel and feeding. Deer also use warm-season grasses as bedding sites. Warm-season grasses are not as good a wildlife food source because the grass is not as palatable as other plant species. For information on establishing native warm-season grasses, see University of Tennessee Extension publication PB 1747, *A Landowner's Guide to Warm-Season Native Grasses in the Mid-South*. Information on establishing native warm-season grasses in Kentucky can be found on the Internet at http://www.uky.edu/Ag/Forestry/TBarnes/Web%20pages/how_to_NWSG.htm.

Fertilizer should not be applied at seeding (usually May to early June). Fertilizer, particularly nitrogen, will encourage growth of undesirable plant species that will compete with the young seedlings. Apply 15 to 30 lb N/A in the summer of the seeding year when the plants are 4 to 6 inches high and there is adequate soil moisture to help the stand become established. Native warm-season grasses are adapted to nutrient-deficient soils with low pH. Therefore, lime and fertilizer are needed only if the soil is extremely low in pH or deficient in nutrients. Apply 100% effective lime at a rate of 2 tons/A if water pH is below 5. Apply 50 lb P₂O₅/A if soil test P is below 10 lb/A. Apply 60 lb K₂O/A if soil test K is below 100 lb/A.

Table 36. Sample plant species for wildlife food plots.	
Cool-Season Legumes	
Alsike clover (perennial)	Subterranean clover (annual)
Arrowleaf clover (annual)	White-Dutch clover (perennial)
Ball clover (annual)	Sweetclover, yellow or white (biennial)
Crimson clover (annual)	Alfalfa (perennial)
Ladino white clover (perennial)	Austrian winter pea (annual)
Red clover (biennial)	Bird's-foot trefoil (perennial)
Warm-Season Legumes	
American jointvetch (annual)	Lablab (annual)
Alyceclover (annual)	Soybean (annual)
Catjang pea (annual)	Common (Kobe) lespedeza (annual)
Cowpea (annual)	Partridge pea (annual)
Cool-Season Grasses	
Oats (annual)	Ryegrass (annual or perennial)
Rye (annual)	Wheat (annual)
Warm-Season Grasses	
Corn (annual)	Japanese millet (annual)
Grain sorghum (milo; annual)	Pearl millet (annual)
Browntop millet (annual)	White proso millet (annual)
German (foxtail) millet (annual)	
Other Plantings	
Buckwheat (annual; warm-season)	Rape, typhoon (annual, cool-season)
Chicory, puna (perennial)	Sunflower (annual, warm-season)
Chufa (annual; warm-season)	Turnip, forage-type (annual; cool-season)
Rape, dwarf essex (perennial; cool-season)	

Table 37. Annual phosphate and potash recommendations for wildlife food plots (lb/A).		
Soil Test Level	P₂O₅	K₂O
High >60 P >300 K	0	0
Medium 60 - 30 P 300 - 200 K	30 - 40	30 - 50
Low¹ <30 P <200 K	40 - 80	50 - 100
¹ The maximum rate should be used if soil test P is less than 6 lb/A or soil test K is less than 90 lb/A.		

LAWNS AND GENERAL TURF

Establishing New Turf

Mix lime, nitrogen, P₂O₅, and K₂O into the top 4 to 6 inches of soil before seeding. Use 1.5 lb N/1000 sq ft and lime, P₂O₅, and K₂O rates recommended in the following tables.

Lime—Use the rate of limestone needed to attain a target pH of 6.4 as indicated in Table 38.

Maintenance of Turf

Lime—Based on the soil test pH, apply limestone at rates indicated under the “Establishing New Turf” section. Apply no more than 70 to 100 lb/1000 sq ft at any one time. Additional lime can be supplied as repeated applications at three- to six-month intervals.

Phosphate and Potash—According to soil test results, apply phosphate or potash at rates indicated in Table 39.

Nitrogen—Apply 1 to 1.5 lb actual N/1000 sq ft per application or 40 to 60 lb N/A. The frequency of nitrogen applications depends on the level of overall maintenance. The following low- and medium-maintenance levels are best for general lawns that get little or no summer irrigation. The high and very high levels usually require some irrigation and a high mowing frequency (see Table 40).

Table 39. Phosphate and potash recommendations for lawns and turf.

Lb/A Soil Test Level	Lb/1000 Sq Ft	
	P ₂ O ₅	K ₂ O
High >60 P 300 K	0 - 1	0 - 1
Medium 60 - 30 P 300 - 200 K	1 - 2	1 - 2
Low <30 P 200 K	3 - 5	3 - 5

Table 38. Rate (lb/1000 sq ft)¹ of bulk agricultural limestone needed to raise soil pH to 6.4, lawns and turf.

Water pH of Sample	Buffer pH of Sample								If Buffer pH Is Unknown
	5.5	5.7	5.9	6.1	6.3	6.5	6.7	6.9	
4.5	320	300	280	250	220	180	150	130	180
4.7	320	300	280	240	200	170	140	120	170
4.9	310	290	260	230	190	150	130	110	160
5.1	310	290	260	220	180	130	100	80	150
5.3	300	280	240	210	160	120	90	70	130
5.5	290	270	230	190	140	100	70	60	120
5.7	280	260	220	170	120	90	60	50	100
5.9	—	240	200	150	100	80	50	40	80
6.1	—	—	180	120	80	60	40	40	60
6.3	—	—	—	90	60	40	40	30	40

¹ Based on 67% RNV ag limestone.

See University of Kentucky Extension publication AGR-53, *Lawn Fertilization in Kentucky*, for more specific details on lawn fertilization.

Table 40. Best months to make nitrogen applications for turf maintenance.

N Applications per Year		Cool-Season Grasses ¹	Warm-Season Grasses ²
Medium	2	Sept. – Oct., Nov. – Dec.	May, July
High	3	Sept. – Oct., Oct. – Nov., Nov. – Dec.	April, June, August
Very high	4	Sept. – Oct., Oct. – Nov., Nov. – Dec., late May – early June (½ rate)	April, May, June, August

¹ Kentucky bluegrass and tall fescue. Red fescue and all cool-season grasses grown in shady lawns should be fertilized only once a year.

² Bermudagrass and zoysiagrass. Zoysiagrass needs only a minimal amount of N after lawn is fully established.

TREE FRUITS, BLACKBERRIES, RASPBERRIES, BLUEBERRIES, AND GRAPES

New Plantings

Lime—Limestone should be applied three to six months before planting and worked into the soil. Tree fruits, blackberries, and raspberries are most productive when soil pH is between 6.4 and 6.6. For plantings of American and French-American hybrid grapes, adjust the soil pH to 6.5 at establishment, and maintain between 5.5 and 6.0 during production. European or *Vitis vinifera* grapes perform best when soil pH is between 6.0 and 7.0. See page 5, and use the table appropriate to the target pH; apply the amount of recommended lime rate.

For blueberries, adjust soil pH to a value between 4.5 and 5.2. See University of Kentucky Extension publication HO-60, *Growing Highbush Blueberries in Kentucky*, for further information on adjusting soil pH to fall in this range.

Nitrogen—Nitrogen fertilization rates depend on the field cropping history and soil types. Apply no more than 100 lb N/A (3.7 oz nitrogen per 100 sq ft). Nitrogen is most effective when it is applied at planting time.

The above recommendations are for establishment only. During subsequent seasons, fertilizer application should be based

Table 41. Phosphate and potash for tree fruits, blackberries, raspberries, blueberries, and grapes.

Lb/A Soil Test Level	P ₂ O ₅		K ₂ O	
	Lb/A	Oz/100 Sq Ft	Lb/A	Oz/100 Sq Ft
High >70 P >300 K	0	0	0	0
Medium 70-35 P 300-200 K	0 - 80	0 - 3	0 - 80	0 - 3
Low <35 P <200 K	80 - 120	3 - 5	80 - 120	3 - 5

Table 42. Magnesium recommendations (lb/A) for tree fruits, blackberries, raspberries, blueberries, and grapes.

Lb/A Soil Test Level	Lb/A Mg
Low below 60	80
Medium 61-120	20 - 80
High above 120	0

on the plant growth rate and condition. Tissue analysis is the most accurate method for determining plant nutrient status in commercial plantings. Contact your county Extension agent to obtain information and a source for plant analysis kits.

Magnesium—Fruit crops require more magnesium than most agronomic crops. The soil should be adjusted if soil test Mg is less than 120 lb of Mg/acre prior to planting. Use Table 42 for adjusting Mg levels in the soil according to the soil test Mg value.

STRAWBERRIES

Establishment

Lime—Limestone should be applied three to six months before planting and worked into the top 4 inches of soil. Strawberries are most productive when the soil pH falls between 6.0 and 6.5. See page 5, and apply the rate of lime to attain a target pH of 6.4. If established plantings need lime, an application of agricultural limestone during the dormant season is best.

Nitrogen—Nitrogen fertilizer should not exceed 60 lb N/A (2.2 oz nitrogen per 100 sq ft) broadcast before planting. Sidedressing with 30 lb N/A in two bands, one on each side of the row with each band placed 2 to 4 inches deep and 8 inches from the plants, is just as effective as broadcast N applications and reduces competition from weeds.

For an early fall application, apply 30 to 40 lb N/A between Aug. 15 and Sept. 10 to promote fruit bud development in the next season.

Spring nitrogen applications are generally avoided during fruiting years because these applications lead to greater vegetative growth, lower fruit yield, delayed ripening, and increased fruit rot.

Magnesium—See Table 42 for recommendations.

Renovation

Use plant analysis during mid-season harvest and fertilize accordingly, or obtain a soil test following harvest. Apply 30 lb N/A (1.1 oz N/100 sq ft), follow lime rate recommendations to attain a target pH of 6.4, and apply phosphate and potash as recommendations in the “Establishment” section. Fertiliza-

Table 43. Phosphate and potash recommendations for strawberries.

Lb/A Soil Test Level	P ₂ O ₅		K ₂ O	
	Lb/A	Oz/100 Sq Ft	Lb/A	Oz/100 Sq Ft
High >70 P >300 K	0	0	0	0
Medium 70 - 35 P 300 - 200 K	0 - 80	0 - 3	0 - 40	0 - 1.5
Low <35 P <200 K	80 - 150	3 - 6	40 - 80	1.5 - 3

tion should be done before any cultivation during renovation. Contact your county Extension office to obtain plant analysis kits.

Types of Application

Broadcast—Apply fertilizer over the tops of the plants when leaves are dry. Avoid possible foliage burn by brushing nitrogen granules off the plant leaves. A canvas attached to the back of the fertilizer applicator works well in brushing fertilizer from the plants.

Sidedress—Banded fertilizer should be placed 2 inches below the soil surface and 6 to 8 inches from the plants in established stands.

