

COLORADO POTATO BEETLE MANAGEMENT

Ric Bessin, Extension Entomologist

The common black and yellow-striped "potato bug", a very familiar insect to home gardeners, is the most serious insect pest of potatoes. Both the striped beetle and the black-spotted, red larva feed on potato leaves. Their damage can greatly reduce yield and even kill plants. In addition to potato, Colorado potato beetle can be a serious pest of tomato, eggplant, and pepper.

The Colorado potato beetle is notorious for its ability to rapidly develop resistance to insecticides that are used repeatedly for control. This has been a serious problem on the east coast for some time, and is becoming more of a problem in Kentucky. With a limited number of insecticides available, some homeowners feel they have exhausted their control options when it becomes resistant to one or more insecticides.



Biology

Colorado potato beetles overwinter in the soil as adults. They become active in the spring as temperatures rise and begin to feed on weeds and volunteer or early planted potatoes, even entering the soil to attack emerging foliage. Female beetles lay orange-yellow eggs in batches of about two dozen or so on the underside of the leaves. Each female can lay 500 or more eggs over a four to five week period. Eggs hatch in four to nine days and the larvae begin to feed on potato foliage. The larvae are humpbacked with two rows of black spots on each side. They usually feed in groups and damage can be severe. The larval stage lasts two to three weeks.

Full grown larvae burrow in the ground to pupate. In five to 10 days, the adult beetle emerges. This insect can go from egg to adult in as little as 21 days. The newly emerged adult female feeds for a few days before egg

laying begins. There are two full and occasionally a partial third generation each year. If foliar sprays are used, an effort should be made to treat just after most eggs have hatched but before serious plant damage occurs.



Resistance Management

Insecticides in the same chemical class usually have the same mode of action, the same method of killing the insect. Resistance develops more rapidly to an insecticide when that insecticide is used repeatedly as the only control measure. Repeated use of one class kills susceptible beetles, leaving those that are resistant. Overuse of one insecticide may favor the development of resistance to other insecticides in the same chemical class. Consequently, to delay or prevent resistance it is important to avoid repeated usage of one particular insecticide by rotating the insecticides used.

Rotation needs to be among different classes of insecticides (see table below). For example, rotation among Warrior and Asana would not be as effective as Platinum, Asana rotation. Because Warrior and Asana are in the same class of chemicals and have the same mode of action, little is gained with this type of rotation. Note that the insecticides marked with an asterisk in the table are available to homeowners

Table 1. Availability of Insecticides to Control Colorado Potato Beetle on Different Crops

INSECTICIDE CLASS	PRODUCT NAME	POTATOES	EGGPLANT	TOMATOES
Organophosphate	Imidan*	OK		
	Di-Syston	OK		OK
Carbamate	Sevin*	OK	OK	OK
	Furadan	OK		
Pyrethroid	Ambush or Pounce	OK	OK	
	Asana XL	OK	OK	OK
	Baythroid	OK		OK
	Capture		OK	
	Mustang			OK
	New Spectracide*	OK		
	Warrior			OK
Chlorinated Hydrocarbon	Methoxychlor*	OK		OK
	Thiodan*	OK	OK	OK
Insect Growth Regulator	Azatin*	OK	OK	OK
Chloronicotinyl	Actara	OK		
	Admire 2F	OK	OK	OK
	Assail			OK
	Platinum	OK	OK	OK
	Provado 1.6F	OK	OK	OK
Spinosad	SpinTor		OK	OK
Abamectin	Agri-Mek	OK		OK

Bacillus thuringiensis var *tenebrionis* (Bt) is effective against small larvae (less than 1/4") and should be applied at egg hatch or when larvae are first seen. A premature treatment may lose much of its effectiveness before the eggs hatch. Larger larvae are more difficult to control with Bt. Azatin, an extract of the neem seed, prevents the larvae from developing normally.

Frequently, control failures with Colorado potato beetle are due to other factors besides just insecticide resistance alone. Timing of sprays is critical for control. Overwintering beetles are attracted to fields over a period of several weeks; spraying an insecticide too early may only control a portion of those beetles. However, waiting until larvae are nearly full grown also increases the chances of control failure. Small larvae are much easier to control with an insecticide than large ones. Using the correct amount of insecticide as well as obtaining complete coverage of the plants is important.

Insecticides should only be used when needed. Potato

plants can withstand considerable defoliation without yield loss. Plants can lose up to 30% of their foliage without yield loss. Generally, insecticides do not need to be applied unless there is more than an average of one beetle or larva per plant. Additionally, some beneficial insects such as birds, predatory stink bugs, and parasitic flies will help to reduce Colorado potato beetle numbers somewhat.

Other non-chemical control measures such as hand picking of adult beetles and immature stages is encouraged as this will aid to delay the development of resistance. Hand picking can be particularly effective in reducing the numbers of overwintering beetles coming to the young plants in the spring. Resistance by Colorado potato beetles should be managed on a field to field basis. While they may be resistant to one insecticide in a particular location, those in other areas within the same county may not have developed resistance to that insecticide.

Revised: 11/03