

## REDBERRY JUNIPER

*Juniperus pinchotii* Sudw.

Plant Symbol = JUPI

Contributed by: USDA NRCS National Plant Data Center



from Native Trees of Texas

### Alternate Names

Pinchot's juniper

### Uses

*Ethnobotanic:* The Comanche used Pinchot's juniper as a treatment for headaches, vertigo, and ghost sickness. For these ailments the Comanche would sprinkle dried leaves over hot coals and inhale the smoke. The Comanche also used an extract of the dried and pulverized roots of Pinchot's juniper for menstrual problems.

### Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's

current status (e.g. threatened or endangered species, state noxious status, and wetland indicator values).

### Description

*General:* Cypress Family (Cupressaceae). Pinchot's juniper grows to be a shrub or small tree, reaching a maximum height of 6 meters. Multiple stems coming from the base of the tree form a dense clump. The bark is thin and ashy-gray colored with longitudinal fissures. The branches are rigid, with slender ascending tips. Pinchot's juniper has white sapwood and reddish-brown heartwood. The leaves on mature fruiting branches are triangular-ovate and pressed together in groups of two or three (1.5-2.5 mm long). The leaf margins are serrated with teeth that point forward. The fruits are variable in size, ranging in color from reddish to copper-brown. Each fruit has either one or two seeds. The seeds are 5 mm long, have a broad oval shape, and are chestnut brown in color.

*Distribution:* For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site.

*Habitat:* Found on open flats, dry hills, in arroyos, and in canyons.

### Adaptation

Low to moderate intensity fires will kill seedlings and saplings. Mature trees, will resprout after top-kill by fire if soil is protecting the basal bud zone. Once top-killed, mature trees require 3-50 years to attain pre-fire height.

### Establishment

Pinchot's juniper requires two years of above average precipitation for seedling establishment. The optimal soil temperature for germination is 64 degrees Fahrenheit. Reproduction also occurs from resprouting of already established plants. Following injury or top removal, Pinchot's juniper will resprout from the base of the stem.

### Management

Fire has been successfully used to prevent the encroachment of Pinchot's juniper on rangeland. For further information regarding the use of fire to manage the tree consult your local land management agency.

### **Pests and Potential Problems**

Grown in its native habitat and using local seed stock, Pinchot's juniper should not be prone to debilitating pests.

### **Cultivars, Improved and Selected Materials (and area of origin)**

Contact your local Natural Resources Conservation Service (formerly Soil Conservation Service) office for more information. Look in the phone book under "United States Government." The Natural Resources Conservation Service will be listed under the subheading "Department of Agriculture."

### **References**

Correl, D.S. & M.C. Johnston 1970. *Manual of the vascular plants of Texas*. Texas Research Foundation, Renner, Texas. 1881 pp.

Great Plains Flora Association 1986. *Flora of the Great Plains*. University Press of Kansas, Lawrence, Kansas. 1392 pp.

Jones, D.E. 1968. *Comanche plant medicine*. Papers in Anthropology 9:1-13.

McPherson, G.R. & H.A. Wright 1997. *Factors affecting reproductive maturity of redberry juniper (Juniperus pinchotii)*. Forest Ecology and Management 21:191-196.

Moerman, D.E. 1998. *Native American ethnobotany*. Timber Press, Portland, Oregon. 927 pp.

Moerman, D.E. 1999. *Native American ethnobotany database: Foods, drugs, dyes and fibers of native North American peoples*. The University of Michigan-Dearborn, Michigan.  
<http://www.umd.umich.edu/cgi-bin/herb>.

Texas A&M University 2002. *Native trees of Texas*. Department of Horticulture, College Station, Texas. <<http://aggie-horticulture.tamu.edu/ornamentals/natives/indexscientific.htm>>. Accessed: 30May2002.

United States Department of Agriculture, Forest Service 2001. *Fire effects information system*. Rocky Mountain Research Station, Fire Science Laboratory, Boise, ID.  
<http://www.fs.fed.us/database/feis/plants/tree/junpin/index.html>

### **Prepared By:**

*Matthew D. Hurteau*  
Formerly USDA, NRCS, National Plant Data Center, c/o Environmental Horticulture Department, University of California, Davis, California

### **Species Coordinator:**

*M. Kat Anderson*  
USDA, NRCS, National Plant Data Center, c/o Plant Sciences Department, University of California, Davis, California

Edited: 30May2002 jsp; 21may03 ahv; 060801 jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

*The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).*

*To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.*

*Read about [Civil Rights at the Natural Resources Conservation Service](#).*