

West National Technical Center 511 NW Broadway, Room 248 Portland, Oregon 97209-3489

June 18, 1990

ECONOMICS TECHNICAL NOTE NO. W-7 200-VI

SUBJECT: ECN - CALCULATING MANURE SPREADER CAPACITY

Purpose. To transmit the above named technical note.

Effective Date. When received.

Filing Instructions. File in Technical Note binder - Economics Section.

ROBERT L. CALDWELL

Head, ESS

Enclosure

DIST:

S (West)

N (ESS)

T (ESS)

United States Department of Agriculture

Soil Conservation Service

West National Technical Center

Portland, Oregon

June 1990

Technical Note

Calculating Manure Spreader Capacity

Economics Series No. W7

United States Department of Agriculture

Soil Conservation Service

WNTC Portland Oregon

ECONOMICS NO. W7

Calculating Manure Spreader Capacity

Though disposing of manure with the least possible cost may be a major concern on some specialized livestock farms, the objective on most farms should be to obtain the maximum crop nutrient benefit from the manure. The producer should also minimize pollution from manure. Crop budgets that include manure as a crop nutrient use some type of manure spreading equipment. A handy way to estimate spreader capacity in different units is given in the following table.

Prepared by:

Shirley J. Elliott Agricultural Economist NENTC October 24, 1989

Page 2 Technical Note Economics No. W7

| MANURE SPREADER CAPA | CITY |
|--|---------------------------|
| Spreader size | Tons of Manure |
| in gallons (8 lb/gal): 1,000 2,000 4,000 | 4 8 16 |
| in bushels (75 lb/bu): 75 100 125 150 | 2.8 3.75 4.7 5.6 |
| in cubic feet (609 lb/cu. ft): 100 200 300 | 3 6 9 |

The dollar value of manure will depend on the handling system and the nutrient content when applied to crops. It can vary considerably from the original amount in the manure, depending on storage, time and method of application. This data is available in the Animal Waste Management Field Manual, currently under revision, and should be used when developing crop budgets. Budgets for individual farmers should be based on a current nutrient analysis of their stored waste.