

United States Department of Agriculture

Forest Service

Rocky Mountain Research Station

General Technical Report RMRS-GTR-125

February 2004



Fourwing Saltbush (Atriplex canescens) Seed Transfer Zones

Stewart C. Sanderson and E. Durant McArthur



Sanderson, Stewart C., and McArthur, E. Durant. 2004. Fourwing saltbush (Atriplex canescens) seed transfer zones. Gen. Tech. Rep. RMRS-GTR-125. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 10 p.

Abstract

Atriplex canescens (Pursh.) Nutt. is the most widespread species of perennial *Atriplex* in North America. Throughout its distributional range, *A. canescens* shows considerable between-population variation. Some of this variation may be due to phenotypic plasticity but most of it appears to be genetic. Mutations, polyploidy, introgressive hybridization, and segregation from interspecific hybrids all appear to have contributed to its extensive heritable variation. Polyploidy is unusually common with numerous chromosome races (2x, 4x, 6x, 8x, 10x, 12x, 14x, 20x).

Fourwing saltbush is widely used for reclamation plantings. Proper identification is important to the utilization of fourwing saltbush in such plantings. While many of the races have been formally named as varieties, others have not. Even though differentiated by ploidy, chemical constituents, geographic distribution, and statistical distribution of morphological characters, races may lack sufficient diagnostic characters to allow facile identification, at least in the herbarium. Rather than combining unnamed races under those that do have a taxonomic name, it seems better at present not to use the formal infraspecific categories in treating the fourwing saltbushes, but to consider them all as races. Seed transfer should be within the geographical distribution limits of each race. The most common race, by far, is Occidentalis. We recommend four overlapping seed transfer zones for race Occidentalis in the United States: (1) Northern Intermountain, (2) Western Great Plains, (3) Colorado Plateau/ Great Basin/Columbia Basin, and (4) Southwestern. Source seed populations from near the planting sites generally do well; and populations generally perform better when moved south and/or to lower elevations than when moved north and/or up in elevation.

The Authors

Stewart C. Sanderson is a research geneticist and **E. Durant McArthur** is a research geneticist and project leader of the Shrubland Biology and Restoration Research Work Unit, Shrub Sciences Laboratory, Provo, Utah. Dr. Sanderson has been at the Shrub Sciences Laboratory since 1981; he received B.S. and M.S. degrees from Brigham Young University, a Ph.D. from the University of Texas at Austin, and postdoctoral training at the University of Durham, England. Dr. McArthur has been with Forest Service Research since 1972 (at the Shrub Sciences Laboratory since 1983); his B.S., M.S., and Ph.D. degrees are from the University of Utah. He spent a postdoctoral sojourn at the University of Leeds, England.

You may order additional copies of this publication by sending your mailing information in label form through one of the following media. Please specify the publication title and series number.

Fort Collins Service Center

Rocky Mountain Research Station Natural Resources Research Center 2150 Centre Avenue, Building A Fort Collins, CO 80526

Contents

Introduction	. 1
Materials and Methods	. 5
Application of Results	. 5
Planting Recommendations	. 6
Selected Germplasms	. 7
References	. 8

Acknowledgments

This work was stimulated and partially funded by the USDI Bureau of Land Management's Great Basin Native Plant Selection and Increase Project and by the USDA Forest Service's National Fire Plan. It draws from work initiated many years ago by our colleagues A. Perry Plummer of the Intermountain Forest and Range Experiment Station (now Rocky Mountain Research Station) and Howard C. Stutz of Brigham Young University. We appreciate review comments from Mike Pellant, Dan Ogle, Loren St. John, Nancy Shaw, and Howard Stutz on earlier versions of the manuscript.

Introduction

Fourwing saltbush, Atriplex canescens (Pursh.) Nutt., is an important shrub for revegetation plantings (Plummer and others 1966; Blauer and others 1976; Plummer 1984). Its uses include mine land rehabilitation (Aldon 1981), shrub-herb interplantings for animal forage (Pendery and Provenza 1987; Rumbaugh and others 1982), fire rehabilitation plantings (Ott and others 2003), rangeland improvement (Monsen and Shaw 1995), and stock for nutrient-poor soils (Glenn and others 1996, 1998). It is widespread in North America, being found from southern San Luis Potosi, Mexico, to southern Alberta and Saskatchewan, Canada, and from the Pacific Coast of California and Baja California to Texas, Oklahoma, Nebraska, Kansas, and the Dakotas (figures 1-5). It is a quality forage plant. Not only does it have fine nutritive quality but it is palatable to wild and domestic animals and tolerates browsing (Cibils and others 1998; Maywald and others 1998; Welch and Monsen 1981, 1984). It shows abundant variation in response to seasons and to rainfall, as well as genetic differentiation. This has resulted in increasing confusion, even by reputable authors (see for example Taylor and Wilkens 1993 who do not recognize the extent of subspecific variation). Fourwing saltbush is widely used for reclamation and restoration plantings. Over 100,000 pounds of seed is sold annually (McArthur and Young 1999). Last year, one seed company alone sold 70,000 pounds (McArthur and Sanderson, personal communication). With this much seed in commerce, land management agencies and seed vendors need guidance in planting. This paper is intended to meet that need.

The species has been planted successfully not only in restoration habitats, i.e., sites on which it naturally occurs, but also on sites in which it does not naturally



Figure 1. Distribution of tetraploid races of fourwing saltbush. Occidentalis and Aptera overlap extensively in Wyoming.





Table 1. Distinguishing characteristics of races of fourwing saltbush.

Race	Bush height	Leaves	Four-winged fruits	Habitat ^a
Angustifolia (2x)	3 to 6 ft	Narrow to very narrow, often long	Unusually large	Sand (U.S.), other soils (Mexico)
Aptera (4x)	1 to 4 ft	Moderately wide, with curving margins	Wings often small and irregular	Badlands and grasslands
Brevis (2x)	1 to 2 ft, a few 3 ft	Narrow to very narrow, shorter	Size normal to somewhat small	Grasslands on old lava flows
Diegensis (6x)	3 ft to as much as 6 ft	Moderately wide, often green colored	Normal size and shape	Sea coast, ravines, canyons
Garrettii (2x)	Usually 1 ft or less	Leaves roundish, often yellow-green	Normal	Shale slopes, often southwest facing
Gigantea (2x)	4 to 8 ft, often partly buried	Not very narrow, but often long	Unusually large	Active dunes
Grandidentatum (20x)	3 to 6 ft	Bluish colored, small to very large, ends often rounded	Strongly toothed, can be large near the beach	Sandy coastal areas
Laciniata (8x)	6 to 8 ft	Small to long and wide	Wings strongly toothed	Bottomlands
Laciniata (12x)	4 to 6 ft	Extremely small to large	Wings with many teeth and cuts	Sandy desert soils
Laciniata (14x)	4 to 6 ft	Short, or large and long. Bluish	Deeply toothed	Coastal dunes
Linearis (2x)	Usually 1 to 2 ft	Usually very narrow, bluish, the sides straight	Wings very small (1/8 in), the body may be swollen	Bottomlands, may be seasonally flooded
Navajoensis (4x)	One to 2 ft	Shortened and exceptionally wide	Normal	Red or blue shale
Nevadensis (6x)	1 to 4 ft, rarely more	Wider than 4x at the same season	Usually more toothed than 4x	Desert sands and slopes
Obtusifolia (8x)	1 to 2 ft	Egg-shaped with a blunt end, greenish gray	Wings small, toothed	Alkaline bottomlands
Occidentalis (4x)	2 to 6 ft	Small to large, narrower than 6x	May be toothed	Valleys and hills, sometimes on sand
Piochensis (2x)	1 to 6 ft	Narrow to very narrow	Normal in size to rather large	Hills, commonly with juniper
Prosopidium (8x)	1 to 4 ft	Short and wide, bluish	Wings small, toothed	Valleys
Prosopidium (10x)	1 to 4 ft	Short and wide, bluish	Wings small, toothed	Valleys
Prostrata (6x)	1 to several inches, flat topped	Moderately long	Normal	Alkaline valley bottoms
Rachelensis (6x)	1 to 5 ft high	Wider than 4x	Smallish, toothed	Valley
Rachelensis (8x)	1 to 2 ft high	Very gray	Smallish, toothed	Alkaline valley bottoms
Vallis (6x)	Usually 1 to 3 ft high, larger in West	Wider than 4x	Normal to smallish	Bottomlands

^a These specific type sites need to be combined with the general race distribution as shown in figures 1-5.

occur, e.g., Wyoming big sagebrush habitats and pinyon-juniper habitats (Monsen and Shaw 1995; Ott and others 2003). It has also been used with success in the Mediterranean Basin and western and southern Asia (Belal and others 1993; Hyder 1981; Khalii and others 1986; Le Houérou 1992, 2000; Sankary and Goodin 1986; Thompson and others 1998).

Fourwing saltbush includes populations with different chromosome numbers that are distributed in distinct geographical patterns. Sanderson and Stutz (2001) described and characterized this variation as chromosome number or chromosomal races based on previously described formal taxonomic designations and geographical, morphological, chemical, and cytogenetic information that they presented (table 1). The racial names in fourwing saltbush are derived, for the most part, from previously applied formal specific or varietal names but some racial names come from geographical or morphological traits of races that had not been formally characterized prior to the description of chromosomal races by Sanderson and Stutz (2001). This chromosomal race knowledge is useful in clarifying the limits of the species and its subspecific and population variation. Chromosome number races range from 2x to 20x-2x, 4x, 6x, 8x, 10x, 12x, 14x, 20x (Dunford 1984, 1985; Sanderson and Stutz 1994, 2001; Stutz and others 1975; Stutz and Sanderson 1979)-and apparently account for much of the existing genetic differentiation. By far the most widespread of the races is the tetraploid Occidentalis (figures 1-5). Fourwing saltbush's dioecious or trioecious outbreeding fosters genetic variation and contributes to the species? successful occupation of diverse habitats (Freeman and others 1993; McArthur and others 1992). The reason the widespread race is called Occidentalis rather than "Canescens" is that the variety name canescens has to be given to plants from the type locality (site from which the species was first collected and described). The type locality for fourwing saltbush is a badland area along the banks of the Missouri River Big Bend in South Dakota, where the plant was collected by the explorers Lewis and Clark and described by Frederick Pursh. The type specimen, and plants growing there now, have relatively wide leaves and belong to race Aptera (4x), or perhaps to their own separate race because they show abundant sprouting from the roots (in figure 1 they are shown in the distribution map with Aptera).

Fourwing saltbush plants from different areas or belonging to different races have been found to vary in tolerance to environmental factors (Glenn and others 1994, 1996; McArthur and others 1983; Mikhiel and others 1992; Plummer and others 1968; Van Epps 1975). Especially important from the point of view of reclamation in cold desert settings is variation for tolerance to low temperatures (Van Epps 1975; Aberdeen Plant Materials Center staff 2002; USDA-NRCS and University of Idaho staff 2001).

Materials and Methods

Chromosome counts were accumulated over a period of 30 years in the laboratory of Howard C. Stutz, now professor emeritus, Brigham Young University in cooperation with the U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Shrub Sciences Laboratory. Recently, a ploidy analyzer instrument (Partec GmbH, Münster, Germany) has been obtained by the Shrub Sciences Laboratory, Provo, Utah, that facilitates the chromosome count work.

Meiotic chromosome counts were made from male flower buds preserved in the cold (-20° F) in vinegar or 5% acetic acid. Pollen mother cells were squashed on a microscope slide and stained with saturated acetocarmine dye (Sanderson and Stutz 1994). More recently, ploidy levels were also determined on the ploidy analyzer using a DAPI stain for DNA and plants of known ploidy for comparison (Koontz 2001).

Flavonoid chemistry was found to be useful in differentiating some races, due to presence or absence of 6-methoxy flavonols, 3-O methylated flavonols, and the flavone tricin. Flavonoid aglycones were obtained from glycosides by acid hydrolysis with 1 N HCl, separated by paper chromatography, and examined over a black light (longwave ultraviolet) both before and after treatment with 5% aqueous aluminum chloride solution (Sanderson and others 1999). We have also examined isozymes to characterize the genetic variation of fourwing saltbush populations (McArthur and others 1986).

We call on our experience and reports in the literature in planting, transplanting, and survival of fourwing saltbush seed source populations taken together with the genetic and chemical variation data cited in the previous paragraphs to make recommendations for seed transfer for revegetation plantings for this important reclamation species.

Application of Results

Knowledge of chromosome numbers and flavonoid chemistry proved invaluable in understanding the variation observed in fourwing saltbush. Previously described varieties were validated and new races were discovered.

Fourwing saltbush is adaptable to seasonal drought and the leaves vary in size from large to small depending on growth status and moisture availability. This is particularly true of the high polyploids. Plants of Grandidentatum (20x) normally have small leaves of a bluish color, but when we grew plants from seed in a barrel of sand, the juvenile plants had leaves about 3 inches long and 1 inch wide that looked somewhat like peach leaves. In spite of strong foliage variability, differences in leaf characters must often be relied upon in distinguishing the different races (table 1). There is a tendency for leaf width to increase with ploidy, at least at the lower ploidy levels. This is the most useful characteristic in distinguishing diploids from tetraploids. With the notable exception of race Garrettii (A. garrettii), diploids have very narrow leaves that are 10 times or more longer than broad, although they vary seasonally to some degree. With care, tetraploids can also be distinguished from hexaploids by leaf width alone, but plants have to be compared in the field and at the same time of year. Leaf color is also useful in some cases. The diploids, except for race Linearis with rather bluish gray-green leaves (sometimes blackish in herbarium specimens), all tend to have more yellowish foliage during dry seasons and winter due to chlorophyll loss, which has been found to prevent damage from excess light energy during times when the plant is not growing (Munné-Bosch and Alegre 2000). Yellowing may be seen slightly in tetraploid plants also, but not usually in higher polyploids. Like Linearis, several Mexican races of higher ploidy also have a bluish leaf color.

A problem with the identification of the diploids Angustifolia, Brevis, and Piochensis is the occurrence of hybrid intermediates between these and local plants of the tetraploid race Occidentalis. Hybrid intermediates are usually also tetraploid, and therefore interbreed with Occidentalis but not with their parental diploid. As a result the diploids remain unchanged and pure, but tetraploids can be somewhat mongrelized by the hybridization and begin to look more like diploids. Tetraploid plants in an area of past or present diploid occurrence are often sufficiently intermediate in form that it is difficult to tell from appearance alone where the diploid leaves off and the tetraploid begins. However, there are seasons when it is easier to distinguish them, and the tetraploids seldom become exactly like diploids in any characteristic. For instance, flavonoids, a kind of plant chemical, are different between diploids and the tetraploids of race Occidentalis (the latter produce 6methoxy flavonols and the former do not). In many cases that have been examined, hybrid derivatives always produce these compounds too.

Planting Recommendations

Many of the races are relatively localized (figures 1 to 5). Many are also found in Mexico and have little ability to survive the winters of the temperate zone in the U.S. Therefore, they are not likely candidates for use in reclamation in this country. In localities where any of the distinctive less common races of fourwing saltbush are encountered, they alone should be used in reseeding. Ordinarily, however, it is the tetraploid Occidentalis, or sometimes Aptera, that are ordinarily used in reclamation or restoration plantings. When an appropriate seed source is chosen, the primary concerns are to seed each race in its own area of natural distribution and to be mindful of ecotypic variability of Occidentalis (see further discussion, below).

As might be expected, resistance to winter injury increases in a northward direction. Of the various Atriplex species and races from warm desert areas or Mexico (and some from Australia) that were planted in gardens at Provo, Utah, over the years, none survived for long and most died during their first winter (Stutz and Sanderson, personal observations). Van Epps (1975) reported on winter survival of fourwing saltbush plants from a number of sites in the Intermountain area. There was variability, but generally plants from southern Utah and areas farther south suffered high mortality when grown in central Utah, and the best survival for this planting area was in plants obtained from Bridger in southern Montana. The Aberdeen, Idaho, Plant Materials Center has recently announced the release of a cold-adapted strain of fourwing saltbush hardier than strains presently in use (Aberdeen Plant Materials Center staff 2002; USDA-NRCS and University of Idaho staff 2001). It should be especially useful in colder areas, such as northern Nevada, northern Utah, and southern Idaho.

The race Aptera, mainly found in Montana (figure 1), is also very cold hardy. Its native habitat is saline soils such as are found in badlands, in areas of generally greater annual precipitation (≥15 inches), especially summer rainfall, than is found over the general fourwing saltbush distributional range. Plants of Aptera from various locations were grown successfully for many years in gardens at Provo, Utah and at the Gordon Creek Wildlife Management Area near Price, Utah, where the annual rainfall is about 15 inches and 11.3 inches, respectively (Ashcroft and others 1992; McArthur and Welch 1982).

Plants from more northern areas generally perform well in areas to the south of their origin. On the other hand, plants from the south typically grow in northern areas with much vigor during summer but are eliminated during the cold season. Cold resistance probably



Figure 6. Recommended seed transfer zones for race Occidentalis: (1) Northern Intermountain, (2) Western Great Plains, (3) Colorado Plateau/Great Basin/Columbia Basin, and (4) Southwestern.

comes at a certain cost, and that is a reduced growth rate. Usually this is not an adaptive disadvantage. Long-term survival of seeded populations depends on successful recruitment by seeded plants. This may or may not happen depending of the site adaptation of the source population—it usually does not happen in off-site plantings, e.g., Wyoming big sagebrush sites (McArthur and Sanderson, personal observation). Seeded plants do produce viable seed in most locations (Noller and others 1984; Monsen and McArthur 1985; Sanderson and McArthur personal observations).

Our general recommendations for planting race Occidentalis are summarized in figure 6. Common garden plantings and revegetation plantings (Fitzsimmons and others 1998; Geist and Edgerton 1984; McArthur and others 1983; Pendleton and others 1992; Peterson and others 1987; Plummer and others 1966; Van Epps 1975) point to a consensus that source populations should not be moved too far north or up in elevation. In general, source populations from near planting sites do well. So, we recommend that source populations not be obtained out of the general zones (figure 6) and not moved up in elevation more than 2,000 feet within their respective zones. The four zones are Northern Intermountain; Western Great Plains; Colorado Plateau/Great Basin/ Columbia Basin ; and Southwestern (figure 6). Stutz and colleagues (Stutz 1982; Stutz and Estrada 1995) recommend that revegetation be by natural evolutionary processes by seeding mixes of compatible seed sources (sources of the same ploidy level), thus stimulating an evolutionary trajectory for the specific seeded site. Should this be the management choice for revegetation, we recommend that the seed mix chosen include source populations from the zone to be rehabilitated.

Selected Germplasms

Five germplasms of fourwing saltbush have been released for reclamation plantings (Davis and others 2002). Germplasm releases are plant materials that have been identified and depending on class (source identified, selected, tested, source identified, cultivar—see Young 1995 and Young and others 1995 for criteria), evaluated for performance and made available for seed increase. Whereas there is a substantive continuing demand for fourwing saltbush seed for rehabilitation plantings (McArthur and Young 1999), most of this need is met by wildland collected seed. McArthur and others (1978) made an economic analysis on the potential profitability for growing fourwing saltbush in seed orchards. However, the ready availability

of wildland-collected seed has dampened the demand for more expensive field-grown seed. Some commercial production, however, is available. The best way to determine if seed is available is to contact state Crop Improvement Associations or the plant materials specialists of the U. S. Department of Agriculture, Natural Resources Conservation Service.

The released germplasms are 'Marana,' 'Rincon,' 'Santa Rita,' "Snake River Plains," and 'Wytana' (Davis and others 2002). All but "Snake River Plains" were released as cultivars, implying that the indigenous population sources underwent selection for specific trait(s) and were tested in comparisons with other fourwing saltbush source populations. "Snake River Plains" is a tested plant material comprising a synthetic composite of four populations. The released plant materials are all of race Occidentalis except 'Wytana' which is of race Aptera. Both 'Marana' (originally collected from El Cajon, San Diego County, California at an elevation of 600 feet) and 'Santa Rita' (originally from south of Tucson, Pima County, Arizona at an elevation of 2400 feet) are from southern locations at relatively low altitudes and are best adapted to the Mojave and Sonoran Deserts, respectively (Davis and others 2002; Cable 1972; United States Department of Agriculture Soil Conservation Service and others 1980). The source population for 'Rincon' is at Rincon Blanco, Rio Arriba County, New Mexico, at 7.800 feet. This release was a result of selection and testing over a wide geographical area (McArthur and others 1984). The source high elevation site (ponderosa pine, pinyon-juniper, and big sagebrush community) preadapted this release to more northern habitats (as far north as central Utah and Nevada) than might be expected. The "Snake River Plains" tested germplasm was selected for its tolerance to the colder, windier northern conditions of Snake River Plain and Northern Great Basin areas (USDA-NRCS and University of Idaho staff, 2001). 'Wytana' is adapted not only to the Aptera geographical area (Fig. 1) but also to somewhat more southern areas (Sanderson and McArthur, personal observations).

We believe that each of the released plant materials should be considered for use in revegetation areas to which they are adapted but that wildland collected materials should be used prudently within their zones of adaptation (our recommendations are in Fig. 6). As a general rule, close habitat and geographical proximity are preferred. And, when greater movement from source sites occurs, down slope and south is better than upslope and north.

References

- Aldon, Earl F. 1981. Long-term plant survival and density data from reclaimed Southwestern coal mine spoils. Great Basin Naturalist. 41: 271-273.
- Aberdeen Plant Materials Center Staff. 2002. Fact sheet: Snake River Plains fourwing saltbush, selected class germplasm. Aberdeen, ID: USDA-NRCS, Aberdeen PMC, P.O. Box 296, Aberdeen, ID 83210. 1 p.
- Ashcroft, Gaylen L.; Jensen, Donald T.; Brown, Jeffrey L. 1992. Utah climate. Logan, UT: Utah Climate Center, Utah State University. 127 p.
- Belal, A. H.; Rammah, A. M.; Hopkins, M. S.; Hansen, L. D.; McArthur, E. D. 1993. Studies of salt-tolerance and chemical analysis of fodder shrubs in Egypt and Utah (USA). In: Leith, H.; Al Masoom, A., eds. Towards the rational use of high salinity tolerant plants, Vol 2. Dordrecht, Netherlands: Kluwer Academic Publishers: 213-220.
- Blauer, A. Clyde; Plummer, A. Perry; McArthur, E. Durant; Stevens, Richard; Giunta, Bruce C. 1976. Characteristics and hybridization of important Intermountain shrubs. II. Chenopod family. Res. Pap. INT-177. Ogden, UT: U.S. Department Agriculture, Forest Service, Intermountain Forest and Range Experiment Station. 42 p.
- Cable, Dwight R. 1972. Fourwing saltbush revegetation trials in southern Arizona. Journal of Range Management. 25: 150-153.
- Cibils, Andrés F.; Swift, David M.; McArthur, E. Durant. 1998. Plant-herbivore interactions in *Atriplex*: current state of knowledge. Gen. Tech. Rep. RMRS-GTR-14. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 31 p.
- Davis, Kathleen, M.; Englert, John M.; Kujawski, Jennifer L. 2002. Improved conservation plant materials released by NRCS and cooperators through September 2002. Beltsville, MD: U. S. Department of Agriculture, Natural Resources Conservation Service. 57 p.
- Dunford, Max. P. 1984. Cytotype distribution of *Atriplex canescens* (Chenopodiaceae) of southern New Mexico and adjacent Texas. Southwestern Naturalist. 29: 223-228.
- Dunford, Max P. 1985. A statistical analysis of morphological variation in cytotypes of *Atriplex canescens* (Chenopodiaceae). Southwestern Naturalist. 30: 377-381.
- Fitzsimmons, Kevin; Lovely, Cynthia; Glenn, Edward. 1998. Growth differences among widely separated geographic accessions of fourwing saltbush (*Atriplex canescens*) in the Great Basin Desert, New Mexico, USA. Arid Soil Research and Rehabilitation. 12: 87-94.
- Freeman, D. Carl; McArthur, E. Durant; Sanderson, Stewart C.; Tiedemann, Arthur R. 1993. The influence of topography on male and female fitness components of *Atriplex canescens*. Oecologia. 93: 538-547.
- Geist, J. Michael; Edgerton, Paul J. 1984. Fourwing saltbush establishment in the Keating Uniform Garden—first year results. Res. Note PNW-416. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station. 8 p.
- Glenn, Edward P.; Moore, David; Sanderson, Stewart; Brown, J. Jed; Lash, Don; Nelson, Mansel; Waugh, Jody. 1998. Comparison of growth and morphology of

Atriplex canescens varieties *occidentalis* and *angustifolia*. Southwestern Naturalist. 43: 176-182.

- Glenn, Edward P.; Olsen, M. W.; Frye, R. J.; Moore, D. W. 1994. How much sodium accumulation is necessary for salt tolerance in subspecies of the halophyte *Atriplex canescens*? Plant Cell and Environment. 17: 711-719.
- Glenn, Edward; Pfister, Rachel; Brown, J. Jed; Thompson, T. Lewis; O'Leary, James. 1996. Na and K accumulation and salt tolerance of *Atriplex canescens* (Chenopodiaceae) genotypes. American Journal of Botany. 83: 997-1005.
- Hyder, S. Z. 1981. Preliminary observations on the performance of some exotic species of *Atriplex* in Saudi Arabia. Journal of Range Management. 34: 208-210.
- Khalii, Jehangir K.; Sawaya, Wajin N.; Hyder, Syed Z. 1986. Nutrient composition of *Atriplex* leaves grown in Saudi Arabia. Journal of Range Management. 39: 104-107.
- Koontz, Jason A.; Soltis, Pamela S. 2001. Polyploidy and segregation analyses in *Delphinium gypsophilum* (Ranunculaceae). Madroño. 48: 90-97.
- Le Houérou, H. N. 1992. The role of saltbushes (*Atriplex* spp.) in arid land rehabilitation in the Mediterranean Basin: a review. Agroforesty Systems. 18: 107-148.
- Le Houérou, Henry N. 2000. Utilization of fodder trees and shrubs in the arid and semiarid zones of West Africa and North Africa. Arid Soil Research and Rehabilitation. 14: 101-135.
- Maywald, Dione; McArthur, E. Durant; Jorgensen, Gary L.; Stevens, Richard; Walker, Scott C. 1998. Experimental evidence for sex-based palatability variation in fourwing saltbush. Journal of Range Management. 51: 650-654.
- McArthur, E. Durant; Freeman, D. Carl; Luckinbill, Leo S.; Sanderson, Stewart C.; Noller, Gary L. 1992. Are trioecy and sexual liability in *Atriplex canescens* genetically based?: evidence from clonal studies. Evolution. 46: 1708-1721.
- McArthur, E. D.; Plummer, A. P.; Van Epps, G. A.; Freeman, D. C.; Jorgensen, K. R. 1978. Producing fourwing saltbush in seed orchards. In: Hyder, Donald N., ed. Proceedings of the First International Rangeland Congress; 1978 August 14-18; Denver, CO. Denver, CO: Society for Range Management: 406-410.
- McArthur, E. Durant; Sanderson, Stewart C.; Freeman, D. Carl. 1986. Isozymes of an autopolyploid shrub, *Atriplex canescens* (Chenopodiaceae). Great Basin Naturalist. 46: 157-160.
- McArthur, E. Durant; Stranathan, Sam E.; Noller, Gary L. 1984. 'Rincon' fourwing saltbush—proven for better forage and reclamation. Rangelands. 6: 62-64.
- McArthur, E. Durant; Stevens, Richard; Blauer, A. Clyde. 1983. Growth performance comparisons among 18 accessions of fourwing saltbush (*Atriplex canescens*) at two sites in central Utah. Journal of Range Management. 36: 78-81.
- McArthur, E. Durant; Welch, Bruce L. 1982. Growth rate differences among big sagebrush (*Artemisia tridentata*) accessions and subspecies. Journal of Range Management. 35: 396-401.
- McArthur, E. Durant; Young, Stanford A. 1999. Development of native seed supplies to support restoration of pinyonjuniper sites. In: Monsen, Stephen B.; Stevens, Richard, comps. Proceedings: ecology and management of pinyonjuniper communities within the Interior West; 1997 September 15-18; Provo, UT. Proc. RMRS-P-9. Ogden,

UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station: 327-330.

- Mikhiel, Gamal S.; Pendleton, Rosemary L.; Meyer, Susan E. 1992. Variation in germination response to temperature and salinity in shrubby *Atriplex* species. Journal of Arid Environments. 22: 39-49.
- Monsen, Stephen G.; McArthur, E. Durant. 1985. Factors influencing establishment of seeded broadleaf herbs and shrubs following fire. In: Sanders, Ken; Durham, Jack, eds. Rangeland fire effects, a symposium; 1984 November 27-29; Boise, ID. Boise, ID: U.S. Department of the Interior, Bureau of Land Management, Idaho State Office: 112-124.
- Monsen, Stephen B.; Shaw, Nancy L. 1995. Interseeding fourwing saltbush (*Atriplex canescens*) with crested wheatgrass (*Agropyron cristatum*) for winter livestock grazing. In: West, Neil E., ed., Rangelands in a sustainable biosphere, proceedings of the Fifth International Rangeland Congress, Volume 1; 1995 July 23-25; Salt Lake City, UT. Denver, CO: Society for Range Management: 381-382.
- Munné-Bosch, S.; Alegre, L. 2000. Changes in carotenoids, tocopherols and diterpenes during drought and recovery, and the biological significance of chlorophyll loss in *Rosmarinus officinalis* plants. Planta. 210: 925-931.
- Noller, Gary L.; Stranathan, Sam E.; McArthur, E. Durant. 1984. Establishment and initial results from a 'Rincon' fourwing saltbush (*Atriplex canescens* [Pursh] Nutt.) seed orchard. In: Tiedemann, Arthur R.; McArthur, E. Durant; Stutz, Howard C.; Stevens, Richard; Johnson, Kendall R., comps. Proceedings—symposium on the biology of *Atriplex* and related chenopods; 1983 May 2-6; Provo, UT. Gen. Tech. Rep. INT-172. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station: 196-204.
- Ott, Jeffrey E.; McArthur, E. Durant; Roundy, Bruce A. 2003. Vegetation of chained and non-chained seedings after wildfire in Utah. Journal of Range Management. 56: 81-91.
- Pendery, Bruce M.; Provenza, Fredrick D. 1987. Interplanting crested wheatgrass with shrubs and alfalfa; effects of competition and preferential clipping. Journal of Range Management. 40: 514-520.
- Pendleton, Burton, K.; Freeman, D. Carl; McArthur, E. Durant; Pendleton, Rosemary L. 1992. Life history features of three sexual morphs of *Atriplex canescens* (Chenopodiaceae) clones grown in a common garden. American Journal of Botany. 79: 376-382.
- Petersen, Joseph L.; Ueckert, Darrell N.; Potter, Robert L.; Juston, James E. 1987. Ecotypic variation in western Texas fourwing saltbush. Journal of Range Management. 40: 361-366.
- Plummer, A. Perry; Christensen, Donald R.; Monsen, Stephen B. 1968. Salt Lake City, UT: restoring big game range in Utah. Publ. 68-3, Salt Lake City, UT: Utah Division of Fish and Game. 183 p.
- Plummer, A. Perry.; Monsen, Stephen, B.; Christensen, Donald R. 1966. Fourwing saltbush, a shrub for future game ranges. Publ. 66-4. Salt Lake City, UT: Utah State Department of Fish and Game. 12 p.
- Plummer, Mark. 1984. Considerations in selecting chenopod species for range seedings. In: Tiedemann, Arthur R.; McArthur, E. Durant; Stutz, Howard C.; Stevens, R.; Johnson, Kendell L., comps. Proceedings—symposium

on the biology of *Atriplex* and related chenopods; 1983 May 4-6, Provo, UT. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station: 183-186.

- Rumbaugh, Melvin D.; Johnson, Douglas A.; Van Epps, Gordan A. 1982. Forage yield and quality in a Great Basin shrub, grass, and legume pasture experiment. Journal of Range Management. 35: 604-609.
- Sanderson, Stewart C.; Stutz, Howard C. 1994. High chromosome numbers in Mojavean and Sonoran Desert *Atriplex canescens* (Chenopodiaceae). American Journal of Botany. 81: 1045-1053.
- Sanderson, Stewart C.; Stutz, Howard C. 2001. Chromosome races of fourwing saltbush (*Atriplex canescens*), Chenopodiaceae. In: McArthur, E. D.; Fairbanks, D. J., comps. Shrubland ecosystem genetics and biodiversity: proceedings; 2000 June 13-15; Provo, UT. Proc. RMRS-P-21. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station: 75-88.
- Sanderson, Stewart C.; Stutz, Howard C.; Stutz, Mildred R.; Roos, Richard C. 1999. Chromosome races in *Sarcobatus* (Sarcobataceae, Caryophyllales). Great Basin Naturalist. 59: 301-314.
- Sankary, Mohamed Nazir; Goodin, J. R. 1986. The American 'Aptera' complex of *Atriplex canescens* (Pursh) Nuttall in Syria: chromosome number, morphology, germination and field performance in arid environments. Research Journal of Aleppo University Agricultural Sciences Series. 8: 17-50.
- Stutz. Howard C. 1982. Broad gene pools required for disturbed lands. In: Aldon, Earl F.; Oaks, Wendall R., eds. Reclamation of mined lands in the Southwest; 1982 October 20-22; Albuquerque, NM. Albuquerque, NM: Soil Conservation Society of America, New Mexico Chapter: 113-118.
- Stutz, Howard C.; Estrada, Orlando J. 1995. Development of genetic diversity in *Atriplex*. In: Schuman, Gerald E.; Vance, George F., eds. Decades later: a time for reassessment, proceedings of the 12th annual meeting of the American Society for Surface Mining and Reclamation; 1995 June 3-8, Gillette, WY. Princeton, WV: American Society for Surface Mining and Reclamation: 762-765.
- Stutz, Howard C.; Melby, James M.; Livingston, Gordon K. 1975. Evolutionary studies of Atriplex: a relic gigas diploid population of *Atriplex canescens*. American Journal of Botany. 62: 236-245.
- Stutz, Howard C.; Sanderson, Stewart C. 1979. The role of polyploidy in the evolution of *Atriplex canescens*. In: Goodin, J. R.; Northington, David K., eds. Arid land plant resources: proceedings of the International Arid Lands

Conference on Plant Resources. Lubbock, TX: Texas Tech University: 615-621.

- Taylor, Dean; Wilken, Dieter H. 1993. Atriplex. In: Hickman, James C., ed. The Jepson manual: higher plants of California. Berkeley: CA. University of California Press: 501-505.
- Thompson, Euan F.; Mirza, Sarwat N.; Afzal, Javed. 1998. Technical note: predicting the components of aerial biomass of fourwing saltbush from height and volume. Journal of Range Management. 51: 323-325.
- U.S. Department of Agriculture, Soil Conservation Service; California Agriculture Experiment Station; California Department of Fish and Game. 1980. Notice of release of 'Marana' fourwing saltbush. In: Upland game report 1964-1980. Sacramento, CA: California Department of Fish and Game and Soil Conservation Service, U.S. Department of Agriculture: 147-150.
- USDA-NRCS and University of Idaho Staff, 2001. Notice of release of Snake River Plains germplasm fourwing saltbush selected class germplasm. Aberdeen, ID: U.S. Department of Agriculture, Natural Resources Conservation Service, and Moscow, ID: University of Idaho, Agricultural Experiment Station. 8 p.
- Van Epps, Gordon A. 1975. Winter injury to fourwing saltbush. Journal of Range Management. 28: 157-159.
- Welch, Bruce L.; Monsen, Stephen B. 1981. Winter crude protein among accessions of fourwing saltbush grown in a uniform garden. Great Basin Naturalist. 41: 343-346.
- Welch, Bruce L.; Monsen, Stephen B. 1984. Winter nutritive value of accessions of fourwing saltbush (*Atriplex canescens*) grown in a uniform garden. In: Tiedemann, Arthur R.; McArthur, E. Durant; Stutz, Howard C.; Stevens, Richard; Johnson, Kendall R., comps. Proceedings—symposium on the biology of *Atriplex* and related chenopods; 1983 May 2-6; Provo, UT. Gen. Tech. Rep. INT-172. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station: 138-144.
- Young, Stanford A. 1995. Verification of germplasm origin and genetic status by seed certification agencies. In: Roundy, Bruce A.; McArthur, E. Durant; Haley, Jennifer S.; Mann, David K., comps. Proceedings: wildland shrub and arid land restoration symposium; 1993 October 19-21; Las Vegas, NV. Gen. Tech. Rep. INT-GTR-315. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 293-295.
- Young, Stanford, A.; Kitchen, Stanley G.; Armstrong, J.; Watson, V. H. 1995. A.O.S.C.A. approves certification guidelines for wildland collected seed. Seed World. 133: 20-21.



The Rocky Mountain Research Station develops scientific information and technology to improve management, protection, and use of the forests and rangelands. Research is designed to meet the needs of the National Forest managers, Federal and State agencies, public and private organizations, academic institutions, industry, and individuals.

Studies accelerate solutions to problems involving ecosystems, range, forests, water, recreation, fire, resource inventory, land reclamation, community sustainability, forest engineering technology, multiple use economics, wildlife and fish habitat, and forest insects and diseases. Studies are conducted cooperatively, and applications may be found worldwide.

Research Locations

Flagstaff, Arizona Fort Collins, Colorado* Boise, Idaho Moscow, Idaho Bozeman, Montana Missoula, Montana Lincoln, Nebraska Reno, Nevada Albuquerque, New Mexico Rapid City, South Dakota Logan, Utah Ogden, Utah Provo, Utah Laramie, Wyoming

*Station Headquarters, Natural Resources Research Center, 2150 Centre Avenue, Building A, Fort Collins, CO 80526.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326 W, Whitten Building, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

