ACGIH. 1986. Documentation of the Threshold Limit Values and Biological Exposure Indices. 5th ed. American Conference of Governmental Industrial Hygienists, Cincinnati, OH.

*ACGIH. 1991. Threshold limit values for chemical substances and physical agents and biological exposure indices for 1991 to 1992. American Conference of Governmental Industrial Hygienists, Cincinnati, OH.

ACS. 1990. Chemcyclopedia 90. J.H. Kuney, ed. The American Chemical Society, Washington, DC:94.

Anders MW, Jakobson I. 1985. Biotransformation of halogenated solvents. Stand J Work Environ Health 11:23-32.

*APHA. 1992a. M-6410, extractable base/neutrals and acids. In: Greenburg AE, Clesceri LS, Eaton AD, eds. Standard methods for the examination of water and wastewater. 18th ed. Washington, DC: American Public Health Association, 6-77-6-89.

*APHA. 1992b. M-6230, volatile halocarbons. In: Greenburg AE, Clesceri LS, Eaton AD, eds. Standard methods for the examination of water and wastewater. 18th ed. Washington, DC: American Public Health Association, 6-55-6-57.

APHA. 1992c. M-6210, volatile organics. In: Greenburg AE, Clesceri LS, Eaton AD, eds. Standard methods for the examination of water and wastewater. 18th ed. Washington, DC: American Public Health Association, 6-26-6-46.

*Atkinson R. 1987. A structure-activity relationship for the estimation of rate constants for the gasphase reactions of OH radicals with organic compounds. Int J Chem Kinet 19:799-828.

*Cited in text

*Atkinson R, Carter WPL. 1984. Kinetics and mechanisms of the gas-phase reaction of ozone with organic compounds under atmospheric conditions. Chem Rev 84:437-70.

*ATSDR. 1989. Decision guide for identifying substance-specific data needs related to toxicological profiles. Agency for Toxic Substances and Disease Registry. Atlanta, GA.

Badaeva LN, Belousva AA, Ba:ida NA, et al. 1990. [Characteristics of the neuro- and immunotoxic effects of transplacental hexachlorobutadiene]. Gig Sanit Sept. 90:75-78. (Russian).

Badaeva LN, Ovsiannikova LM, Kiseleva NI, et al. 1985. [manifestation of neurotoxic effect of chlororganic pesticide hexachlorobutadiene postnatal period of ontogenesis in rats.] Arkh Anat Gist01 Em Briol Aug. 89:44-49. (Russian).

Baggett JM, Berndt WO. 1984. Renal and hepatic glutathione concentrations in rats after treatment with hexachloro-1,3-butadiene and citrinin. Arch Toxicol 56:46-49.

*Bai CL, Canfield PJ, Stacey NH. 1992. Effects of hexachloro-1,3-butadiene and 1,1,2,2tetrachloroethylene on individual serum bile acids. Toxicology and Industrial Health 8: 19 1-203.

Ban M, de Ceaurriz J. 1988. Probenecid-induced protection against acute hexachloro-1,3-butadiene and methyl mercury toxicity to the mouse kidney. Toxicol Lett 40:71-76.

Barkley J, Bunch J, Bursey JT, et al. 1980. Gas chromatography mass spectrometry computer analysis of volatile halogenated hydrocarbons in man and his environment - A multimedia environmental study. Biomed Mass Spectrometry 7: 139-147.

*Barnes DG, Dourson M. 1988. Reference dose (RfD): Description and use in health risk assessments. Regul Toxicol Pharmacol 8:471-486.

Bauer S, Wolff I, Werner N, et al. 1992. Toxicological investigations in the semiconductor industry: I. Studies on the acute oral toxicity of a complex mixture of waste products from the aluminium plasma etching process. Toxicology and Industrial Health 8: 141-156.

Bernard A, Lauwerys RR. 1991. Proteinuria: changes and mechanisms in toxic nephropathies. Toxiocology 21: 373-405.

Berndt WO, Mehendale HM. 1978. Effect of hexachlorobutadiene on renal function in the rat. Toxicol Appl Pharmacol 45 :296-297.

Berndt WO, Mehendale HM. 1979. Effects of hexachlorobutadiene (HCBD) on renal function and renal organic ion transport in the rat. Toxicology 14:55-65.

*Bristol DW, Crist HL, Lewis RG, et al. 1982. Chemical analysis of human blood for assessment of environmental exposure to semivolatile organochlorine chemical contaminants. J Anal Toxicol 6:269-275.

*Brown KW, Donnelly KC. 1988. An estimation of the risk associated with the organic constituents of hazardous and municipal waste landfill leachates. Hazardous Waste and Hazardous Materials 5: 1-30.

*Camanzo J, Rice CP, Jude DJ. 1987. Organic priority pollutants in nearshore fish from 14 Lake Michigan tributaries and embayments, 1983. J Great Lakes Res 13 :296-309.

*CDC/ATSDR. 1990. Biomarkers of organ damage or dysfunction for the renal, hepatobiliary and immune systems. Atlanta, GA: CDC/ATSDR Subcommittee on Biomarkers of Organ Damage and Dysfunction, Centers for Disease Control, Agency for Toxic Substances and Disease Registery. Summary report, August 27, 1990.

*Class T, Ballschmiter K. 1987. Global baseline pollution studies. Freznius Z. Anal Chem 327: 198-204.

Class T, Ballschmiter K. 1986. Chemistry of organic traces in air VI: Distribution of chlorinated Cl - C4 hydrocarbons in air over the northern and southern Atlantic ocean. Chemosphere 15:413-427.

*Cole RH, Frederick RE, Healy RE, et al. 1984. Preliminary findings of the priority pollutants monitoring project of the nationwide urban runoff program. J WPCF 56:898-908.

*CRISP. 1993. Computer Retrieval of Information on Scientific Projects. National Institutes of Health, Division of Research Grants, Bethesda, MD. May 13, 1993.

*Davis ME. 1984. Changes of hexachlorobutadiene nephrotoxicity after piperonyl butoxide treatment. Toxicology 30:217-225.

Davis ME. 1988. Effects of AT-125 on the nephrotoxicity of hexachloro-1,3-butadiene in rats. Toxicol Appl Pharmacol 95 :44-52.

*Davis ME, Berndt WO, Mehendale HM. 1980. Disposition and nephrotoxicity of hexachloro-1,3butadiene. Toxicology 16: 179-191.

de Ceaurriz J, Ban M. 1990. Role of γ -glutamyltranspeptidase and β -lyase in the neptrhotoxicity of hexachloro-1,3-butadiene and methyl mercury in mice. Toxicol Lett 50:249-256.

*de Ceaurriz J, Gagnaire F, Ban M, et al. 1988. Assessment of the relative hazard involved with airborne irritants with additional hepatotoxic or nephrotoxic properties in mice. J Appl Toxicol 8:417-422.

*De Meester C, Mercier M, Poncelet F. 1980. Mutagenic activity of butadiene, hexachlorobutadiene, and isoprene. Ind Environ Xenobio Pot Int Conf 195-203.

*Dekant W, Schrenk D, Vamvakas S, et al. 1988a. Metabolism of hexachloro-1,3-butadiene in mice: *in vivo* and *in vitro* evidence for activation by glutathione conjugation. Xenobiotica 18:803-816.

*Dekant W, Urban G, Gorsman C, et al. 1991. Thioketene formation from γ-haloalkenyl 2nitrophenyl disulfides: models for biological reactive intermediates of cytotoxic S-conjugates. J Am Chem Soc 113:5120-5122.

*Dekant W, Vamvakas S, Anders MW. 1990a. Review section: Bioactivation of hexachlorobutadiene by glutathione conjugation. Food Chem Toxicol 28:285-293.

*Dekant W, Vamvakas S, Berthold K, et al. 1986. Bacterial β-lyase mediated cleavage and mutagenicity of cysteine conjugates derived from the nephrocarcinogenic alkenes trichloroethylene, tetrachloroethylene and hexachlorobutadiene. Chem Biol Interactions 60:31-45.

*Dekant W, Vamvakas S, Henschler D, et al. 1988b. Enzymatic conjugation of hexachloro-1,3butadiene with gultathione. Formation of l-(Glutathion-S-yl)-1,2,3,4,4-pentachlorobuta-1,3-diene and 1,4-bis(glutathion-S-yl)-1,2,3,4-tetrachlorobuta-1,3-diene. Drug Metabolism and Disposition 16:701-706.

*Dekant W, Vamvakas S, Koob M, et al. 1990b. A mechanism of haloalkene-induced renal carcinogenesis. Environ Health Perspect 88: 107-110.

*DeVault DS. 1985. Contaminants in fish from Great Lakes harbors and tributary mouths. Arch Environ Contam Toxicol 14:587-594.

*Driscoll TR, Hamdan HH, Wang G, et al. 1992. Concentrations of individual serum or plasma bile acids in workers exposed to chlorinated aliphatic hydrocarbons. Br J Ind Med 49:700-705.

*Duprat P, Gradiski D. 1978. Percutaneous toxicity of hexachlorobutadiene. Acta Pharmacol Toxicol 43:346-353.

*Eichelberger JW, Bellar TA, Donnelly JP, et al. 1990. Determination of volatile organics in drinking water with USEPA method 524.2 and the ion trap detector. J Chromatogr Sci 28:460-467.

*Elder V, Proctor B, Hites R. 1981. Organic compounds found near dump sites in Niagara Falls, New York. Environ Sci Technol 15: 1237-1243.

*EPA. 1976. An ecological study of hexachlorobutadiene (HCBD). Washington, DC: U.S. Environmental Protection Agency, Office of Toxic Substances. EPA/560/6-76-010.

*EPA. 1980. Ambient water quality criteria document for hexachlorobutadiene. Washington, DC: U.S. Environmental Protection Agency, Office of Water Regulations and Standards. EPA 440/5-80-053.

*EPA. 1982a. Chlorinated hydrocarbons-method 612. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory.

*EPA. 1982b. U.S. Environmental Protection Agency. Federal Register 47:26992-27008.

*EPA. 1982c. Test method. Base/neutrals and acids methods 625. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory.

*EPA. 1982d. U.S. Environmental Protection Agency. Federal Register 47:58029-58030.

*EPA. 1984. Health effects assessment for hexachlorobutadiene. Washington, DC: U.S. Environmental Protection Agency, Office of Emergency and Remedial Response, Office of Solid Waste and Emergency Response. EPA/540/1-86-053.

*EPA. 1986. Gas chromatography/mass spectrometry for semivolatile organics: capillary column technique-method 8270. In: Test methods for evaluating solid waste. SW-846. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response.

*EPA. 1987a. U.S. Environmental Protection Agency. Part II. Federal Register. 52:25690-25717.

*EPA, 1987b. U.S. Environmental Protection Agency. Part II. Federal Register. 52:25942-25953.

EPA. 1987c. Measurement of hydrolysis rate constants for evaluation of hazardous waste land disposal: Vol. 1. Data on 32 chemicals. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development. EPA/600/S3-86/043.

EPA. 1987d. Measurement of hydrolysis rate constants for evaluation of hazardous waste land disposal: Vol. 2. Data on 54 chemicals. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development. EPA/600/S3-87/019.

*EPA. 1988a. U.S. Environmental Protection Agency. Part II. Federal Register. 53:4500-4539.

*EPA. 1988b. U.S. Environmental Protection Agency. Part V. Federal Register 53:38642-38654.

*EPA. 1988c. U.S. Environmental Protection Agency. Part II. Federal Register 53:4500-4532.

EPA. 1988d. Measurement of hydrolysis rate constants for evaluation of hazardous waste land disposal: Vol. 3. Data on 70 chemicals. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development. EPA/600/S3-88/028.

EPA. 1988e. Interim protocol for measuring hydrolysis rate constants in aqueous solutions. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development. EPA/600/S3-88/014.

*EPA. 1989a. Ambient water quality criteria document addendum for hexachlorobutadiene. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Water Regulations and Standards. Final draft ECAO-CIN-652.

*EPA. 1989b. Health advisory for hexachlorobutadiene. Washington, DC: U.S. Environmental Protection Agency, Office of Water. PB91-160663.

*EPA. 1989c. Measurement of volatile organic compounds in water by purge and trap capillary column gas chromatography with photoionization and electrolytic conductivity detectors in series - method 502.2. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory.

*EPA. 1989d. Measurement of volatile aromatic and unsaturated organic compounds in water by purge and trap gas chromatography - method 503.1. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory.

*EPA. 1989e. Measurement of purgeable organic compounds in water by capillary column gas chromatography/mass spectrometry - method 524.2. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory.

*EPA. 1990a. Application of open-tubular columns to SW-846 GC methods. In: Test methods for evaluating solid waste. SW-846. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. EPA/600/S4-90/021.

*EPA. 1990b. Method IP-1A (In: Compendium of methods for the determination of air pollutants in indoor air. Washington, DC: U.S. Environmental Protection Agency, Atmospheric Research and Exposure Assessment Laboratory, EPA/600/S4-90/1010.

*EPA. 1990c. Method 8120A. Chlorinated hydrocarbons by gas chromatography. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory.

*EPA. 1990d. U.S. Environmental Protection Agency. Part II. Federal Register 55:22520-22720.

*EPA. 1990e. Volatile organic compounds by gas chromatography/mass spectrometry (GC/MS): capillary column technique-method 8260. In: Test methods for evaluating solid waste. SW-846. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response.

*EPA. 1990f. Pathway analysis of chemical hydrolysis for 14 RCRA chemicals. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development. EPA/600/M-89/009.

*EPA. 199Og. U.S. Environmental Protection Agency. Part III. Federal Register 503:47229.

*EPA. 1990h. Interim methods for the development of inhalation reference doses. Washington, DC: U.S. Environmental Protection Agency, Office of Heath and Environmental Assessment. EPA/600/8-90/066A.

*EPA. 1991. U.S. Environmental Protection Agency. Part III. Federal Register 56:7134-7240.

*EPA. 1993. U.S. Environmental Protection Agency. Health effects summary tables. Supplement No. 1 to the March 1993 annual update. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development. EPA 540-R-93-058.

*Ewing BB, Chian ESK. 1977. Monitoring to detect previously unrecognized pollutants in surface waters. Appendix: organic analysis data. National Technical Information Service. EPA-560/6-77-015A.

Flynn LR, Bass SW, Meierer RE. 1991. Headspace screening/capillary column GC/MS analysis for volatile organics: validation studies and applications. Waste Testing and Quality Assurance 106-114.

*FSTRAC. 1990. Summary of state and federal drinking water standards and guidelines. Washington, DC: Federal-State Toxicology and Regulatory Alliance Committee, Chemical Communication Subcommittee. 7: 17.

Ford JH, Evans J. 1985. Distribution of 5'-nucleotidase in the tissues of sheep and the effect of kidney and liver lesions on the activity of enzyme in plasma and urine. Res Vet Sci 39: 103-109.

*Fox ME, Carey JH, Oliver BG. 1983. Compartmental distribution of organochlorine contaminants in the Niagara River and the western basin of Lake Ontario. J Great Lakes Res 9:287-294.

*Gage JC. 1970. The subacute inhalation toxicity of 109 individual chemicals. Brit J Industr Med 27: 1-8.

*Garle MJ, Fry J. 1989. Detection of reactive metabolites in vitro. Toxicology 54:101-110.

Gartland KP, Beddell CR, Lindon JC, et al. 1991. Computer patter recognition analysis of PMR spectroscopic data from urine obtained from rats in various nephrotoxicity states. Proceedings, Fourth International Symposium on Nephrotoxicity, Guilford, England, UK 1989. Marcel Dekker, Inc., New York, NY 519-524.

Gartland KP, Bonner FW, Nicholson JK. 1989. Investigations into the biochemical effects of region-specific nephrotoxins. Mol Pharmacol 35:242-250.

Gartland KP, Sweatman BC, Beddell CR, et al. 1991. Proton nuclear magnetic resonance urinalysis studies on the biochemical effects of a nephrotoxic dose of uranyl nitrate in the Fischer 344 rat. Proceedings, Fourth International Symposium on Nephrotoxicity, Guilford, England, UK 1989. Marcel Dekker, Inc., New York, NY 525530.

*Gietl YS, Anders MW. 1991. Biosynthesis and biliary excretion of S-conjugates of hexachlorobuta-1,3-diene in the perfused rat liver. Drug Metabolism and Disposition 19:274-277.

*Gietl Y, Vamvakas S, Anders MW. 1991. Intestinal absorption of S-(pentachlorobutadienyl)glutathione and S-(pentachlorobutadienyl)-L-cysteine, the glutathione and cysteine S-conjugates of hexachlorobuta-1,3-diene. Drug Metabolism and Disposition 19:703-707.

*Govind R, Flaherty PA, Dobbs RA. 1991. Fate and effects of semivolatile organic pollutants during anaerobic digestion of sludge. Water Res 25:547-556.

Gradiski D, Duprat P, Magadur JL, et al. 1975. [Etude toxicologique experimentale de l'hexachlorobutadiene]. Eur J Toxicol 8: 180-187. (French).

Green T, Odum J. 1985. Structure/activity studies of the nephrotoxic and mutagenic action of cysteine conjugates of chloro- and fluoroalkenes. Chem Biol Interactions 54: 15-3 1.

*Hardin BD, Bond GP, Sikov MR, et al. 1981. Testing of selected workplace chemicals for teratogenic potential. Stand J Work Environ Health 7:66-75.

*Harleman JH, Seinen W. 1979. Short-term toxicity and reproduction studies in rats with hexachloro-(1,3)-butadiene . Toxicol Appl Pharmacol 47: 1-14.

*Hauser TR, Bromberg SM. 1982. EPA's monitoring program at Love Canal 1980. Environmental Monitoring and Assessment 2:249-271.

*Haworth S, Lawlor T, Mortelmans K, et al. 1983. Salmonella mutagenicity test results for 250 chemicals. Environ Mutagen Suppl 1: 3-142.

*HazDat. 1993. Agency for Toxic Substances and Disease Registery (ATSDR), Atlanta, GA. October, 1993.

*Henschler D, Dekant W. 1990. Nephrocarcinogenic xenobiotics. Toxicol Lett 53 : 105-110.

*Hewitt WR, Brown EM. 1984. Nephrotoxic interactions between ketonic solvents and halogenated aliphatic chemicals. Fund Appl Toxicol 4:902-908.

*Hook JB, Ishmael J, Lock EA. 1983. Nephrotoxicity of Hexachloro-1:3-butadiene in the rat: The effect of age, sex, and strain. Toxicol Appl Pharmacol 67:122-131.

*Hook JB, Rose MS, Lock EA. 1982. The nephrotoxicity of hexachloro-1:3-butadiene in the rat: Studies of organic anion and cation transport in renal slices and the effect of monooxygenase inducers. Toxicol Appl Pharmacol 65: 373-382.

*HSDB. 1992. Hazardous Substances Data Bank. National Library of Medicine, National Toxicology Information Program, Bethesda, MD. January 20, 1992.

*HSDB. 1993. Hazardous Substances Data Bank. National Library of Medicine, National Toxicology Information Program, Bethesda, MD. May 13, 1993.

*IARC. 1979. IARC monographs on the evaluation of the carcinogenic risk of chemicals to humans. Vol. 20: Hexachlorobutadiene. World Health Organization, Lyon, France.

*IARC. 1987. IARC monographs on the evaluation of carcinogenic risk of chemicals to humans. Suppl 7. Overall evaluations of carcinogenicity: Updating of IARC monographs volumes 1-42. World Health Organization, International Agency for Research on Cancer, Lyon, France, 64.

*IRIS. 1992. Integrated Risk Information System. U.S. Environmental Protection Agency, Washington, DC. March 26, 1992.

*IRIS. 1993. Integrated Risk Information System. U.S. Environmental Protection Agency, Washington, DC. May 14, 1993.

*Ishmael J, Lock EA. 1986. Nephrotoxicity of hexachlorobutadiene and its glutathione-derived conjugates. In: Presented at the Fourth International Symposium of the Society of Toxicologic Pathologists, June 5-7, 1985. Washington, DC. Toxicol Pathol 14:258-262.

*Ishmael J, Pratt I, Lock EA. 1982. Necrosis of the pars recta (S₃ segment) of the rat kidney produced by hexachloro-1,3-butadiene. J Pathol 138:99-1 13.

*Ishmael J, Pratt I, Lock EA. 1984. Hexachloro-1,3-butadiene-induced renal tubular necrosis in the mouse. J Pathol 142:195-203.

*Jaffe DR, Hassall CD, Brendel K. 1983. *In vivo* and *in vitro* nephrotoxicity of the cysteine conjugate of hexachlorobutadiene. J Toxicol Environ Health 11:857-867.

*Johnson LD, Young JC. 1983. Inhibition of anaerobic digestion by organic priority pollutants. J Water Pollut Control Fed 55: 1141-1 149.

*Jones TW, Gerdes RG, Ormstad K, et al. 1985. The formation of both a mono- and bis-substituted glutathione conjugate of hexachlorobutadiene by isolated hepatocytes and following *in vivo* administration to the rat. Chem Biol Interactions 56:251-267.

*Jones TW, Chen Q, Schaeffer V, et al. 1988. Immunohistochemical localization of glutamine transaminase K, a rat kidney cysteine conjugate β -lyase, and the relationship to the segment specificity of cysteine conjugate nephrotoxicity. Ma1 Pharm 341621-627.

*Jonker D, Jones MA, van Bladeren PJ, et al. 1993a. Acute (24 hr) toxicity of a combination of four nephrotoxicants in rats compared with the toxicity of the individual compounds. Food Chem Toxicol 3 1:45-52.

*Jonker D, Woutersen RA, van Bladeren PJ, et al. 1993b. Subacute (4-wk) oral toxicity of a combination of four nephrotoxins in rats: comparison with the toxicity of the individual compounds. Food Chem Toxicol 3 1: 125-136.

*Junk GA, Ford CS. 1980. A review of organic emissions from selected combustion processes. Chemosphere 9: 187-230.

Kaminksy R, Kaiser KL, Hites RA. 1983. Fates of organic compounds from Niagara Falls dumpsites in Lake Ontario. J Great Lakes Res 9:183-189.

*Kastl PE, Hermann EA. 1983. Quantitative gas chromatographic determination of hexachloro-1,3butadiene in whole rat blood at part per trillion levels. Journal of Chromatography 280:390-393.

Keith LH, Garrison AW, Allen FR, et al. 1976. Identification of organic compounds in drinking water from thirteen U.S. cities. In: Identification & analysis of organic pollutants in water. Ann Arbor, MI: Ann Arbor Science Publishers, Inc., 329-373.

Kennedy GL, Graepel GJ. 1991. Acute toxicity in the rat following either oral or inhalation exposure. Toxicol Lett 56:317-326.

Khalturin GU, Chudin VA, Andriushkeeva NI. 1984. [Hexachlorobutadiene distribution in rats during chronic intragastric adminstration.] Gig Sanit Oct. 10:91. (Russian).

King JW. 1989. Fundamentals and applications of supercritical fluid extraction in chromatographic science. J Chromatog Sci 27:355-364.

Kluwe WM, McNish R, Smithson K, et al. 1981. Depletion by l,2-dibromoethane, 1,2-dibromo-3chloropropane, tris(2,3-dibromopropyl)phosphate, and hexachloro-1,3-butadiene of reduced nonprotein sulfhydryl groups in target and non-target organs. Biochem Pharmacol 30:2265-2271.

Knocke WR, Conley L, van Benschoten JE. 1992. Impact of dissolved organic carbon on the removal of iron during water treatment. Water Res 26:1515-1522.

*Kociba RJ Gehring PJ, Humiston CG, et al. 1971. Toxicologic study of female rats administered hexachlorobutadiene or hexachlorobenzene for thirty days. The Dow Chemical Company, Midland, Michigan.

*Kociba RJ, Keyes DG, Jersey GC. 1977a. Results of a two year chronic toxicity study with hexachlorobutadiene in rats. Am Ind Hyg Assoc J 38589-602.

Kociba RJ, Schwetz BA, Keyes DG. 1977b. Chronic toxicity and reproduction studies of hexachlorobutadiene in rats. Environ Health Perspect 21:49-53,

*Koob M, Dekant W. 1992. Biotransformation of the hexachlorobutadiene metabolites 1-(glutathion-S-yl)-pentachlorobutadiene and l-(cystein-*S*-yl)-pentachlorobutadiene in the isolated perfused rat liver. Xenobiotica 22:125-138.

*Kotzias D, Klein W, Korte F. 1975. [Beitrage Zur Gkologischen Chemie CVI Vorkommen Von Xenobiotika Im Sickerwasser Von Mtilldeponien.] Chemosphere 5:301-306. (German)

Kuehl DW, Leonard EN, Welch KJ, et al. 1980. Identification of hazardous organic chemicals in fish from the Ashtabula River, Ohio, and Wabash River, Indiana. J Assoc Off Anal Chem 63: 1238-1244.

Kuo CH, Hook JB. 1983. Effects of age and sex on hexachloro-1,3-butadiene toxicity in the Fischer 344 rat. Life Sci 33:517-523.

*Laska AL, Bartell CK, Laseter JL. 1976. Distribution of hexachlorobenzene and hexachlorobutadiene in water, soil, and selected aquatic organisms along the lower Mississippi River, Louisiana. Bull Environ Contam Toxicol 15:535-542.

Lau YL, Oliver BG, Krishnappan BG. 1989. Transport of some chlorinated contaminants by the water, suspended sediments, and bed sediments in the St. Clair and Detroit Rivers. Environ Toxicol Chem 8:293-301.

*LeBel GK, Williams DT. 1986. Pesticide and industrial chemical residues: Determination of helogenated contaminants in human adipose tissue. J Assoc Off Anal Chem 69:451-458.

*Levins P, Adams J, Brenner P, et al. 1979. Sources of toxic pollutants found in influents to sewage treatment plants VI. Integrated interpresentation. NTIS PB81-219685.

Ligocki M, Levenberger C, Pankow J. 1985. Trace organic compounds in rain-II. Gas scavenging of neutral organic compounds. Atmos Environ 19: 1609-1617.

*Li RT, Going JE, Spigarelli JL. 1976. Sampling and analysis of selected toxic substances. Task 1B. Hexachlorobutadiene. NTIS PB-253.941.

Lock EA. 1989. Mechanism of nephrotoxic action due to organohalogenated compounds. Toxicol Lett 46:93-106.

*Lock EA, Ishmael J. 1979. Original investigations: The acute toxic effects of hexachloro-1:3butadiene on the rat kidney. Arch Toxicol 43:47-57.

*Lock EA, Ishmael J. 1981. Hepatic and renal nonprotein sulfhydryl concentration following toxic doses of hexachloro-1,3-butadiene in the rat: The effect of Aroclor 1254, Phenobarbitone, or SKF 525A treatment. Toxicol Appl Pharmacol 57:79-87.

*Lock EA, Ishmael J. 1985. Effect of the organic acid transport inhibitor probenecid on renal cortical uptake and proximal tubular toxicity of hexachloro-1,3-butadiene and its conjugates. Toxicol Appl Pharmacol 81: 32-42.

Lock EA, Ishmael J, Hook JB. 1984. Nephrotoxicity of hexachloro-1,3-butadiene in the mouse: The effect of age, sex, strain, monooxygenase modifiers, and the role of glutathione. Toxicol Appl Pharmacol 72:484-494.

*Lock EA, Ishmael J, Pratt I. 1982. Hydropic change in rat liver induced by hexachloro-1 : 3butadiene. J Appl Toxicol 6:315-320.

*Lock EA, Pratt IS, Ishmael J. 1985. Hexachloro-1,3-butadiene-induced hydropic change in mouse liver. J Appl Toxicol 5:74-79.

Lopez-Avila V, Beckert WF, Billets S. 1991. Supercritical fluid extraction and its application to environmental analysis. Waste Testing and Quality Assurance 141-153.

*Lucas SV. 1984. GCYMS (gas chromatography-mass spectrometry) analysis of organics in drinking water concentrates and advanced waste treatment concentrates, Volume I. NTIS PB85-128221.

*Mabey WR, Smith JH, Pod011 RT, et al. 1982. Aquatic fate process data for organic priority pollutants. Report to U.S. Environmental Protection Agency, Office of Water Regulations and Standards, Washington, DC, by SRI International, Menlo Park, CA. EPA 440/4-81-014.

*MacFarlane M, Foster JR, Gibson GG, et al. 1989. Cysteine conjugate β-lyase of rat kidney cytosol: Characterization, immunocytochemical localization, and correlation with hexachlorobutadiene nephrotoxicity. Toxicol Appl Pharmacol 98: 185-197.

Malins DC, McCain BB, Brown DW, et al. 1984. Chemical pollutants in sediments and diseases of bottom-dwelling fish in Puget Sound, Washington. Environ Sci Technol 18:705-713.

Markwell RD, Comrel DW, Gabric AJ. 1989. Bioaccumulation of lipophilic compounds from sediments by oligochaetes. Wat Res 22: 1443-1450.

*McConnell G, Ferguson DM, Pearson CR. 1975. Chlorinated hydrocarbons and the environment. Endeavor 34: 13-18.

Mes J. - 1981. Experiences in human milk analysis for halogenated hydrocarbon residues. Inter J Environ Anal Chem 9:283-299.

*Mes J, Davies DJ, Turton D. 1982. Polychlorinated biphenyl and other chlorinated hydrocarbon residues in adipose tissue of Canadians. Bull Environ Contam Toxicol 28:97-104.

*Mes J, Davies DJ, Turton D. 1985. Environmental contaminants in human fat: A comparison between accidental and nonaccidental causes of death. Ecotoxicol Environ Safety 10:70-74.

*Montgomery JH, Welkom LM. 1990. Groundwater chemicals desk reference. Chelsea, MI: Lewis Publications, Inc. 334-336.

*NAS/NRC. 1989. Bilogic markers in reproductive toxicology. National Academy of Sciencs of National Research Council. Washington, DC: National Academy Press, 15-35.

*Nash JA, King LJ, Lock EA, et al. 1984. The metabolism and disposition of hexachloro-1:3butadiene in the rat and its relevance to nephrotoxicity. Toxicol Appl Pharmacol 73: 124-137.

*NATICH. 1991. National Air Toxics Information Clearing House: NATICH database report on state, local and EPA air toxics activities. Report to U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards. Research Triangle Park, NC, by Radian Corporation, Austin, TX. EPA-450/3-91-018.

*NIOSH. 1981. Tier II mutagenic screening of 13 NIOSH priority compounds: Individual compound report hexachloro- 1,3-butadiene . Cincinnati, OH: National Institute for Occupational Safety and Health, PB83-152397.

*NIOSH. 1990. Hexachlorobutadiene - method P&CAM 307-update. In: NIOSH Manual of Analytical Methods. 3rd Ed. 4th Supplement. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Services, Centers for Disease Control, National Institute for Occupational Safety and Health.

*NIOSH. 1992. Recommendations for occupational safety health, compendium of policy documents and statements. U.S. Department of Health and Human Services, Centers for Disease Control, NIOSH, Cincinnati, OH.

*NSF. 1975. Research program on hazard priority ranking of manufactured chemicals (Chemicals 1-20) 9-A-I. NTIS PB-263161.

*NTP. 1991. National Toxicology Program. Toxicity studies of hexachloro-1,3-butadiene in B6C3F₁ mice (feed studies). Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health. NIH publication no. 91-3120.

Oliver, BG. 1985. Desorption of chlorinated hydrocarbons from spiked and anthropogenically contaminated sediments. Chemosphere 14: 1087-1106.

*Oliver BG, Bourbonniere RA. 1985. Chlorinated contaminants in surficial sediments of Lakes Huron, St. Clair and Erie: implications regarding sources along the St. Clair and Detroit rivers. J Great Lakes Res 11:366-372.

*Oliver BG, Charlton MN. 1984. Chlorinated organic contaminants on settling particulates in the Niagara River vicinity of Lake Ontario. Environ Sci Technol 18:903-908.

*Oliver BG, Niimi AJ. 1983. Bioconcentrations of chlorobenzenes from water by rainbow trout: correlations with partition coefficients and environmental residues. Environ Sci Technol 10: 148-152.

Oliver BG, Nicol KD. 1985. Field testing of a large volume liquid-liquid extraction device for halogenated organics in natural waters. Intern J Environ Anal Chem 25:275-285.

Oliver BG, Niimi AJ. 1988. Trophodynamic analysis of polychlorinated biphenyl congeners and other chlorinated hydrocarbons in the Lake Ontario Ecosystem. Environ Sci Technol 22:388-397.

Oliver BG, Charlton MN, Durham RW. 1989. Distribution, redistribution, and geochronology of polychlorinated biphenyl congeners and other chlorinated hydrocarbons in Lake Ontario Sediments. Environ Sci Technol 23 :200-208.

*OSHA. 1989. Occupational Safety and Health Administration: Part III. Federal Register 54:2332-2959. January 19.

*OTA. 1990. Neurotoxicity: Identifying and controlling poisons of the nervous system. Washington, DC: Office of Technology Assessment, U.S. Congress. OTA-BA-436.

Pankow JF, Rosen ME. 1988. Determination of volatile compounds in water by purging directly to a capillary column with whole column cryotrapping. Environ Sci Technol 22:398-405.

*Pankow JF, Isabelle LM, Asher WE. 1984. Trace organic compounds in rain. 1. Sampler design and analysis by adsorption/thermal desorption (ATD). Environ Sci Technol 18:3 10-3 18.

Patterson JW, Kodukala PS. 1981. Emission and effluent control: biodegradation. Chem Eng Prog 77:48-55.

Payan JP, Beydon D, Fabry JP, et al. 1993. Partial contribution of biliary metabolites to nephrotoxicity, renal content and excretion of [¹⁴C]hexachloro-l,3-butadiene in rats. J Appl Toxicol 13:19-24.

*Payan JP, Fabry JP, Beydon D, et al. 1991. Biliary excretion of hexachloro-1,3-butadiene and its relevance to tissue uptake and renal excretion in male rats. J Appl Toxicol 11:437-442.

*Pearson CR, McConnell G. 1975. Chlorinated C₁ and C₃ hydrocarbons in the marine environment. Proc R Sot Lond Biol 189:305-332.

*Pellizzari ED. 1978. Quantification of chlorinated hydrocarbons in previously collected air samples. EPA-450/3-78-112.

*Pellizzari ED. 1982. Analysis for organic vapor emissions near industrial and chemical waste disposal sites. Environ Sci Technol 16:781-785.

Pereira WE, Rostad CE, Chlou CT, et al. 1988. Contamination of estuarine water, biota, and sediment by halogenated organic compounds: A field study. Environ Sci Technol 22:772-778.

*Piet GJ, Zoeteman BC. 1980. Organic water quality changes during sand bank and dune filtration of surface waters in the Netherlands. J Amer Water Works Assoc 72:400-404.

*Plumb RH. 1991. The occurrence of appendix IX organic constituents in disposal site ground water. Ground Water Monitoring Review 157-164.

Poncelet F, Duverger-van Bogaert M, Lambotte-Vandepaer M, et al. 1984. Mutagenicity, carcinogenicity, and teratogenicity of industrially important monomers. Kirch-Volders M, ed. New York, Plenum Press, p. 205-279.

*Pospisil PA, Marcus MF, Kobus MA. 1991. The application of supercritical fluid capillary chromatography to the analysis of appendix-VIII and IX compounds. Waste Testing and Quality Assurance 154-169.

Poteryaeva G. 1966. [The effect of hexachlorobutadiene on off-springs of albino rats.] Gig Sanit 31:33. (Russian)

Pratt IS, Lock EA. 1988. Deacetylation and further metabolism of the mercapturic acid of hexachloro-1,3-butadiene by rat kidney cytosol *in vitro*. Arch Toxicol 62:341-345.

Public Law 101-541. 1990. Clean Air Act Amendments of 1990. Environmental Reporter 352-353.

Reichert D. 1983. Metabolism and disposition of hexachloro(1,3)butadiene in rats. In: Hayes AW, Schnell RC, Miya TS, eds. Developments in the Science and Practice of Toxicology, Elsevier Science Publishers, 411-414.

*Reichert D, Schutz S. 1986. Mercapturic acid formation is an activation and intermediary step in the metabolism of hexachlorobutadiene. Biochem Pharmacol 35 : 1271-1275.

*Reichert D, Neudecker T, Schutz S. 1984. Mutageneicity of hexachlorobutadiene, perchlorobutenoic acid and perchlorobutenoic acid chloride. Mutat Res 137: 89-93.

*Reichert D, Neudecker T, Spengler U. 1983. Mutagenicity of dichloroacetylene and its degradation products trichloroacetyl chloride, trichloroacryloyl chloride and hexachlorobutadiene. Mutat Res 117:21-29.

*Reichert D, Schutz S, Metzler M. 1985. Excretion pattern and metabolism of hexachlorobutadiene in the rats: Evidence for metabolic activation by conjugation reactions. Biochem Pharmacol 34:499-505.

Rodriguez OM, Desideri PG, Lepri L, et al. 1991. Simultaneous separation and determination of hydrocarbons and organochlorine compounds by using a two-step microcolumn. J Chromatogr 555:221-228.

*Roldan-Arjona T, Garcia-Pedrajas MD, Luque-Romero FL, et al. 1991. An association between mutagenicity of the Ara test of *salmonella typhimurium* and carcinogenicity in rodents for 16 halogenated aliphatic hydrocarbons. Mutagenesis 6: 199-205.

Rosenfeld JK, Plumb RH. 1991. Ground water contamination at wood treatment facilities. Ground Water Monitoring Review XI: 133-140.

*Ruth JH. 1986. Odor thresholds and irritation levels of several chemical substances: A review. J Am Ind Hyg Assoc 47:A-142-A151.

*Saillenfait AM, Bonnet P, Guenier JP, et al. 1989. Inhalation teratology study on hexachloro-1,3butadiene in rats. Toxicol Lett 47:235-240.

*Sauer TC. 1981. Volatile organic compounds in open ocean and coastal surface waters. Organic Geochemistry 3:91-101.

*Sax NI, Lewis RJ. 1989. Dangerous properties of industrial materials. 7th ed. New York, NY: Van Nostrand Reinhold, p. 2687.

"Schiffmann D, Reichert D, Henschler D. 1984. Induction of morphological transformation and unscheduled DNA synthesis in Syrian hamster embryo fibroblasts by hexachlorobutadiene and its putative metabolite pentachlorobutenoic acid. Cancer Lett 23 :297-305.

*Schnellmann RG, Lock EA, Mandel LJ. 1987. A mechanism of S-(1,2,3,4,4-pentachloro-1,3butadienyl)-L-cysteine toxicity to rabbit renal proximal tubules. Toxicol Appl Pharmacol 90:513-521.

Schrenk D, Dekant W. 1989. Short communication. Covalent binding of hexachlorobutadiene metabolites to renal and hepatic mitochondrial DNA. Carcinogenesis 10: 1139-1 141.

Schwartz BS, Ford DP, Bolla KI, et al. 1990. Solvent-associated decrements in olfactory function in paint manufacturing workers. Am J Ind Med 18:697-706.

Schwetz BA, Norris JM, Kociba RJ. 1974. Reproduction study in Japanese quail fed hexachlorobutadiene for 90 days. Toxicol Appl Pharmacol 30:255-265.

*Schwetz BA, Smith FA, Humiston CG. 1977. Results of a reproduction study in rats fed diets containing hexachlorobutadiene. Toxicol Appl Pharmacol 42: 387-398.

*Shah JJ, Heyerdahl EK. 1988. National ambient volatile organic compounds (VOCs) data base update. EPA/600/3-38/010a.

*Shah JJ, Singh HB. 1988. Distribution of volatile organic chemicals in outdoor and indoor air. Environ Sci Technol 22: 1381-1388.

Shen TT. 1982. Air quality assessment for land disposal of industrial wastes. Environ Management 6:297-305.

*Singh HB, Sales LJ, Smith A, et al. 1980. Atmospheric measurements of selected hazardous organic chemicals. Menlo Park, CA: SRI International. Project No. 7774:6.

*Singh HB, Sales LJ, Stiles RE. 1982. Distribution of selected gaseous organic mutagens and suspect carcinogens in ambient air. Environ Sci Technol 16:872-880.

*Stanker LH, Watkins B, Vanderlaan M, et al. 1990. Analysis of heptachlor and related cyclodiene insecticides in food products. In: ACS Symposium Series 451. Immunoassays for trace chemical analysis. Washington, DC; American Chemical Society, 108-123.

*Staples CA, Werner AF, Hoogheem TJ. 1985. Assessment of priority pollutant concentrations in the United States using Storet database. Environ Toxicol Chem 4: 131-142.

Stein VB, Narang RS. 1984. Chlorinated compounds and phenols in tissue, fat, and blood from rats fed industrial waste site soil extract: methods and analysis. J Assoc Off Anal Chem 67:111-116.

Stonard MD, Gore CW, Oliver JA, et al. 1987. Urinary enzymes and protein patterns as indicators of injury to different regions of the kidney. Fund Appl Toxicol 9:339-351.

*Stott WT, Quast JF, Watanabe PG. 1981. Differentiation of the mechanisms of oncogenicity of 1,4-dioxane and 1,3-hexachlorobutadiene in the rat. Toxicol Appl Pharmacol 60:287-300.

*Tabak HH, Quave SA, Mashni CI, et al. 1981. Biodegradability studies with organic priority pollutant compounds. J Water Pollut Control Fed 53: 1503-1518.

Theiss JC, Stoner GD, Shimkin MB, et al. 1977. Test for carcinogenicity of United States drinking waters by pulmonary tumor response in strain A mice. Cancer Res 37:2717-2720.

Thomann RV. 1989. Bioaccumulation model of organic chemical distribution in aquatic food chains. Environ Sci Technol23:699-707.

*TRI90. 1992. Toxic Chemical Release Inventory. National Library of Medicine, National Toxicology Information Program, Bethesda, MD.

U.S. Air Force. 1988. Interactions amond drinking and ground water contaminants on renal and hepatic function. Washington, DC. AFOSR-TR-88-0717.

*Vamvakas S, Kordowich FJ, Dekant W, et al. 1988a. Mutagenicity of hexachloro-1,3-butadiene and its S-conjugates in the Ames test-role of activation by the mercapturic acid pathway in its nephrocarcinogenicity. Carcinogenesis 9:907-910.

*Vamvakas S, Muller DA, Dekant W, et al. 1988b. DNA-binding of sulfur-containing metabolites from ³⁵S-(pentachlorobutadienyl)-L-cysteine in bacteria and isolated renal tubular cells. Drug Metabolic Interact 6: 349-358.

Van de Meent D, den Hollander HA, Pool WG, et al. 1986. Organic micropollutants in Dutch coastal waters. Wat Sci Tech 18:73-81.

*Van Duuren BL, Goldschmidt BM, Loewengart G, et al. 1979. Carcinogenicity of halogenated olefinic and aliphatic hydrocarbons in mice. JNCI 63: 1433-1439.

*Verschueren K. 1983. Handbook of environmental data on organic chemicals. Van Nostrand Reinhold Company, New York, NY: pp. 717-718.

*Walters SM. 1990. Clean-up techniques for pesticides in fatty foods. Anal Chim Acta 236:77-82.

*Wang F, Ji Z, Wang D. 1991. A rapid pretreatment procedure for determining trace organochlorine compounds in biological samples by capillary gas chromatography. Microchemical Journal 44 : 67-7 1.

*Wild D, Schutz S, Reichert D. 1986. Mutagenicity of the mercapturic acid and other S-containing derivatives of hexachloro-1,3-butadiene. Carcinogenesis 7:431-434.

*Wolf CR, Berry PN, Nash JA, et al. 1984. Role of microsomal and cytosolic glutathione Stransferases in the conjugation of hexachloro-1:3-butadiene and its possible relevance to toxicity. J Pharmacol Exp Ther 228 : 202-208.

*Yang RS. 1988. Hexachloro-1,3-butadiene: toxicology, metabolism, and mechanisms of toxicity. Rev Environ Contam Toxicol 101: 121-137.

*Yang RS, Abdo KM, Elwell MR. 1989. Subchronic toxicology studies of hexachloro-1,3-butadiene (HCBD) in B6C3F₁ mice by dietary incorporation. JEPTO 9:323-332.

*Yip G. 1976. Survey of hexachloro-1,3-butadiene in fish, eggs, milk, and vegetables. J Assoc Off Anal Chem 59:559-561.

*Yurawecz MP, Dreifuss PA, Kamps LR. 1976. Determination of hexachloro-1,3-butadiene in spinach, eggs, fish, and milk by electron capture gas-liquid chromatography. J Assoc Off Anal Chem 59:552-558.

*Zoeteman BC, Harmsen K, Linders JB, et al. 1980. Persistent organic pollutants in river water and groundwater in the Netherlands. Chemosphere 9:231-249.