

Table 36. (Corresponds to MMWR Table 1.) Estimated average annual number of persons with self-reported current asthma by age, sex, race, ethnicity, region, and poverty level: National Health Interview Survey, United States, 2001 - 2004

	Number of Persons with Asthma										
	All ages	Children	Adults	Age							
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	15 - 19	20 - 24	25 - 34	35 - 64	65 +
Total	20,172,000	6,196,000	13,976,000	1,151,000	3,834,000	5,778,000	1,904,000	1,498,000	2,377,000	7,336,000	2,073,000
Sex											
Male											
Female	8,548,000	3,634,000	4,913,000	680,000	2,342,000	2,380,000	940,000	586,000	855,000	2,447,000	699,000
Race	11,624,000	2,561,000	9,063,000	472,000	1,492,000	3,398,000	964,000	912,000	1,522,000	4,889,000	1,374,000
White											
Male	15,371,000	4,144,000	11,227,000	659,000	2,608,000	4,331,000	1,391,000	1,139,000	1,801,000	5,966,000	1,807,000
Female	6,362,000	2,390,000	3,972,000	390,000	1,567,000	1,773,000	664,000	462,000	647,000	2,015,000	617,000
Black	9,009,000	1,754,000	7,255,000	269,000	1,041,000	2,558,000	727,000	677,000	1,154,000	3,951,000	1,190,000
Male	3,149,000	1,346,000	1,803,000	331,000	802,000	935,000	342,000	257,000	336,000	905,000	176,000
Female	1,362,000	794,000	568,000	180,000	506,000	373,000	178,000	84,000	111,000	256,000	48,000
Other races NTA	1,787,000	552,000	1,235,000	151,000	296,000	562,000	164,000	174,000	225,000	649,000	128,000
Male	1,652,000	706,000	946,000	162,000	424,000	512,000	171,000	101,000	240,000	464,000	91,000
Female	824,000	450,000	374,000	110,000	270,000	234,000	97,000	40,000	97,000	176,000	34,000
Ethnicity	828,000	255,000	572,000	52,000	154,000	278,000	73,000	61,000	143,000	288,000	56,000
Hispanic or Latino											
Male	2,014,000	904,000	1,110,000	205,000	554,000	605,000	221,000	131,000	253,000	538,000	112,000
Female	930,000	523,000	407,000	123,000	323,000	259,000	118,000	51,000	90,000	184,000	41,000
Puerto Rican	1,084,000	381,000	703,000	82,000	231,000	346,000	103,000	81,000	163,000	354,000	71,000
Male	526,000	228,000	298,000	45,000	142,000	158,000	63,000	37,000	58,000	146,000	34,000
Female	234,000	130,000	104,000	30,000	84,000	56,000	28,000*	15,000*	13,000	54,000	11,000*
Mexican	292,000	98,000	194,000	15,000*	59,000	103,000	35,000	22,000	45,000	93,000	23,000
Male	959,000	444,000	515,000	104,000	274,000	299,000	101,000	67,000	130,000	241,000	42,000
Female	458,000	267,000	191,000	62,000	162,000	130,000	54,000	29,000	48,000	89,000	15,000*
Not Hispanic or Latino	501,000	177,000	324,000	41,000	112,000	168,000	48,000	38,000	82,000	152,000	28,000
Male	18,158,000	5,292,000	12,866,000	946,000	3,280,000	5,173,000	1,683,000	1,366,000	2,124,000	6,798,000	1,961,000
Female	7,618,000	3,112,000	4,506,000	556,000	2,020,000	2,121,000	821,000	535,000	765,000	2,263,000	658,000
Region	10,540,000	2,180,000	8,360,000	390,000	1,260,000	3,052,000	861,000	831,000	1,359,000	4,535,000	1,303,000
Northeast											
Midwest	4,187,000	1,299,000	2,888,000	228,000	821,000	1,110,000	385,000	295,000	429,000	1,614,000	415,000
South	5,095,000	1,536,000	3,559,000	306,000	894,000	1,593,000	527,000	442,000	624,000	1,760,000	542,000
West	6,893,000	2,221,000	4,672,000	442,000	1,379,000	1,897,000	621,000	453,000	823,000	2,466,000	709,000
Ratio of Family Income to Poverty Threshold	3,997,000	1,140,000	2,857,000	174,000	741,000	1,179,000	370,000	307,000	502,000	1,497,000	407,000
0 - .99											
1.00 - 2.49	2,686,000	988,000	1,698,000	272,000	566,000	837,000	292,000	300,000	245,000	792,000	219,000
2.50 - 4.49	4,678,000	1,552,000	3,126,000	346,000	938,000	1,369,000	380,000	401,000	589,000	1,395,000	630,000
4.50 and above	4,237,000	1,282,000	2,955,000	175,000	841,000	1,300,000	402,000	295,000	602,000	1,539,000	383,000

*The estimate is unreliable because the relative standard error of the estimate is 30%–50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 37. (Corresponds to MMWR Table 2.) Estimated average annual prevalence percents for self-reported current asthma by age, sex, race, ethnicity, region and poverty level: National Health Interview Survey, United States, 2001-2004

	Persons with Asthma per 100 Population										
	All ages	Children	Adults	Age							
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	15 - 19	20 - 24	25 - 34	35 - 64	65 +
Total	7.20	8.50	6.70	5.80	9.40	7.50	9.50	7.70	6.30	6.70	6.20
Sex											
Male	6.20	9.80	4.90	6.70	11.20	6.20	9.10	6.10	4.60	4.60	4.90
Female	8.10	7.20	8.30	4.90	7.50	8.70	9.80	9.20	7.90	8.70	7.10
Race											
White	6.90	7.70	6.70	4.60	8.70	7.40	9.20	7.70	6.30	6.70	6.10
Male	5.90	8.70	4.90	5.30	10.20	6.00	8.60	6.20	4.50	4.60	4.90
Female	8.00	6.70	8.30	3.80	7.10	8.80	9.80	9.30	8.00	8.70	7.10
Black	9.20	12.50	7.60	11.50	13.20	9.00	11.50	9.60	7.00	7.40	6.40
Male	8.50	14.50	5.40	12.20	16.30	7.70	11.70	7.10	5.10	4.70	4.50
Female	9.70	10.40	9.50	10.80	9.90	10.10	11.40	11.60	8.50	9.60	7.70
Other races NTA	6.50	8.40	5.60	6.40	9.20	6.00	8.40	4.70	5.50	5.60	6.50
Male	6.40	10.50	4.40	8.50	11.40	5.40	9.10	3.80	4.40	4.20	5.60
Female	6.60	6.20	6.70	4.20	6.80	6.60	7.70	5.60	6.60	7.00	7.20
Ethnicity											
Hispanic or Latino	5.40	6.90	4.60	5.10	7.70	4.70	7.10	4.00	3.90	4.80	5.40
Male	4.90	7.90	3.30	6.00	8.80	3.90	7.30	3.00	2.70	3.30	4.70
Female	5.90	6.00	5.80	4.20	6.60	5.60	6.90	5.10	5.30	6.20	6.00
Puerto Rican	14.40	18.60	12.30	13.40	20.30	13.60	20.80	12.70	10.20	12.20	13.10
Male	12.90	20.00	9.00	16.40	22.50	9.60	18.30	9.8*	4.70	9.50	9.5*
Female	15.80	17.10	15.30	9.8*	17.90	17.50	23.40	15.70	15.40	14.60	15.80
Mexican	4.00	5.00	3.40	3.70	5.60	3.40	4.80	2.90	3.00	3.60	4.10
Male	3.70	5.80	2.40	4.40	6.50	2.80	4.80	2.50	2.10	2.60	3.10
Female	4.30	4.10	4.40	3.00	4.70	4.10	4.80	3.40	4.10	4.70	4.80
Not Hispanic or Latino	7.40	8.90	7.00	6.00	9.80	8.00	9.90	8.40	6.80	6.90	6.20
Male	6.40	10.20	5.10	6.90	11.80	6.60	9.50	6.70	5.00	4.70	4.90
Female	8.40	7.50	8.70	5.10	7.70	9.30	10.30	10.00	8.40	8.90	7.20
Region											
Northeast	7.90	9.90	7.20	6.80	11.00	8.20	10.50	8.70	6.70	7.50	5.80
Midwest	7.50	9.00	7.00	6.50	9.50	8.30	10.50	8.80	6.80	6.60	6.70
South	6.70	8.40	6.10	6.00	9.30	6.70	8.60	6.40	5.80	6.20	5.80
West	6.90	7.10	6.80	4.00	8.10	7.20	8.90	7.60	6.10	6.90	6.60
Ratio of Family Income to Poverty Threshold											
0 - .99	10.20	10.70	9.90	9.60	11.20	9.20	10.80	9.50	7.70	11.30	9.00
1.00 - 2.49	7.80	8.60	7.50	6.90	9.10	7.70	8.70	7.80	7.00	7.90	7.10
2.50 - 4.49	6.80	7.70	6.40	4.00	9.00	7.20	8.90	7.40	6.30	6.20	6.30
4.50 and above	6.50	8.00	6.10	3.80	9.10	7.10	9.60	7.60	5.90	6.20	4.20

*The estimate is unreliable because the relative standard error of the estimate is 30%-50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 38. (Corresponds to MMWR Table 5.) Estimated average annual number of persons reporting an asthma attack in the past year by age, sex, race, ethnicity, region, and poverty level: National Health Interview Survey, United States, 2001-2004

	Number of Persons with an Asthma Attack										
	All ages	Children	Adults	Age							
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	15 - 19	20 - 24	25 - 34	35 - 64	65 +
Total	11,656,000	4,076,000	7,580,000	876,000	2,526,000	3,208,000	1,041,000	796,000	1,370,000	4,125,000	922,000
Sex											
Male	4,839,000	2,433,000	2,406,000	512,000	1,578,000	1,257,000	511,000	315,000	432,000	1,219,000	273,000
Female	6,817,000	1,643,000	5,175,000	364,000	947,000	1,950,000	531,000	481,000	938,000	2,907,000	649,000
Race											
White	8,790,000	2,734,000	6,057,000	509,000	1,730,000	2,413,000	776,000	585,000	1,051,000	3,349,000	790,000
Male	3,547,000	1,610,000	1,937,000	298,000	1,064,000	951,000	382,000	237,000	333,000	996,000	238,000
Female	5,243,000	1,124,000	4,120,000	211,000	666,000	1,461,000	394,000	348,000	718,000	2,353,000	552,000
Black	1,857,000	868,000	988,000	243,000	522,000	503,000	156,000	158,000	188,000	498,000	91,000
Male	803,000	517,000	286,000	123,000	341,000	190,000	73,000	59,000	58,000	129,000	20,000*
Female	1,054,000	352,000	703,000	120,000	181,000	313,000	83,000	99,000	131,000	369,000	71,000
Other races NTA	1,009,000	474,000	535,000	124,000	274,000	292,000	109,000	53,000	130,000	279,000	41,000
Male	489,000	306,000	183,000	91,000	173,000	116,000	56,000	19,000*	41,000	94,000	15,000*
Female	520,000	167,000	353,000	33,000	100,000	176,000	53,000	33,000	89,000	185,000	26,000
Ethnicity											
Hispanic or Latino	1,203,000	551,000	652,000	136,000	326,000	340,000	128,000	66,000	146,000	333,000	68,000
Male	543,000	326,000	217,000	77,000	200,000	137,000	69,000	22,000*	46,000	107,000	22,000
Female	660,000	225,000	435,000	59,000	126,000	204,000	59,000	45,000	100,000	226,000	46,000
Puerto Rican	356,000	149,000	207,000	39,000	87,000	101,000	38,000	22,000*	42,000	101,000	27,000
Male	156,000	84,000	72,000	25,000	52,000	38,000*	**	**	9,000*	30,000	11,000*
Female	201,000	65,000	135,000	14,000*	35,000	64,000	22,000	9,000*	32,000	71,000	16,000*
Mexican	525,000	256,000	269,000	61,000	153,000	154,000	60,000	27,000	67,000	136,000	21,000
Male	254,000	162,000	92,000	34,000	98,000	68,000	37,000	7,000*	23,000*	50,000	5,000*
Female	271,000	94,000	177,000	27,000	55,000	87,000	24,000	20,000	43,000	86,000	16,000
Not Hispanic or Latino	10,453,000	3,525,000	6,929,000	740,000	2,199,000	2,867,000	913,000	730,000	1,224,000	3,793,000	854,000
Male	4,296,000	2,107,000	2,189,000	435,000	1,378,000	1,121,000	442,000	293,000	386,000	1,112,000	251,000
Female	6,157,000	1,417,000	4,739,000	305,000	821,000	1,747,000	472,000	436,000	838,000	2,681,000	603,000
Region											
Northeast	2,390,000	855,000	1,534,000	171,000	552,000	600,000	203,000	155,000	241,000	900,000	166,000
Midwest	2,951,000	1,012,000	1,939,000	241,000	595,000	869,000	277,000	234,000	358,000	1,013,000	233,000
South	3,986,000	1,457,000	2,529,000	329,000	907,000	1,044,000	328,000	249,000	467,000	1,369,000	338,000
West	2,329,000	752,000	1,578,000	135,000	472,000	695,000	233,000	158,000	303,000	843,000	185,000
Ratio of Family Income to Poverty Threshold											
0 - .99	1,657,000	654,000	1,003,000	208,000	357,000	479,000	157,000	163,000	159,000	501,000	111,000
1.00 - 2.49	2,736,000	1,014,000	1,722,000	264,000	611,000	741,000	194,000	205,000	342,000	844,000	276,000
2.50 - 4.49	2,539,000	908,000	1,631,000	144,000	605,000	754,000	246,000	176,000	332,000	860,000	175,000
4.50 and above	2,364,000	693,000	1,671,000	100,000	439,000	605,000	213,000	69,000	323,000	1,136,000	84,000

*The estimate is unreliable because the relative standard error of the estimate is 30%–50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 39. (Corresponds to MMWR Table 6.) Estimated average annual prevalence percents for self-reported asthma attacks by age, sex, race, ethnicity, region, and poverty level: National Health Interview Survey, United States, 2001-2004

	Persons with an Asthma Attack per 100 population										
	All ages	Children	Adults	Age							
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	15 - 19	20 - 24	25 - 34	35 - 64	65 +
Total	4.1	5.6	3.6	4.4	6.2	4.1	5.2	4.1	3.6	3.7	2.7
Sex											
Male	3.5	6.5	2.4	5.0	7.6	3.2	4.9	3.2	2.3	2.3	1.9
Female	4.7	4.6	4.8	3.7	4.7	5.0	5.4	4.9	4.9	5.1	3.4
Race											
White	4.0	5.1	3.6	3.5	5.7	4.1	5.1	4.0	3.7	3.7	2.7
Male	3.3	5.9	2.4	4.0	6.9	3.2	4.9	3.2	2.3	2.3	1.9
Female	4.6	4.3	4.7	3.0	4.5	5.0	5.3	4.8	5.0	5.2	3.3
Black	5.4	8.0	4.2	8.5	8.5	4.8	5.3	5.9	3.9	4.1	3.3
Male	5.0	9.4	2.7	8.4	11.0	3.9	4.8	5.0	2.7	2.3	1.8*
Female	5.7	6.6	5.4	8.6	6.0	5.6	5.8	6.6	4.9	5.5	4.3
Other races NTA	4.0	5.6	3.1	4.9	5.9	3.4	5.4	2.5	3.0	3.3	2.9
Male	3.8	7.1	2.1	7.0	7.4	2.7	5.2	1.8*	1.8	2.2	2.5*
Female	4.1	4.1	4.1	2.7	4.4	4.2	5.6	3.1	4.1	4.5	3.3
Ethnicity											
Hispanic or Latino	3.2	4.2	2.7	3.4	4.6	2.6	4.1	2.0	2.3	2.9	3.3
Male	2.9	4.9	1.8	3.8	5.5	2.0	4.2	1.3*	1.4	1.9	2.5
Female	3.6	3.5	3.6	3.0	3.6	3.3	4.0	2.8	3.2	4.0	3.9
Puerto Rican	9.7	12.2	8.5	11.7	12.5	8.7	12.4	7.6*	7.2	8.5	10.6
Male	8.6	12.9	6.2	13.4	14.0	6.5	**	8.7*	3.3*	5.4	9.9*
Female	10.9	11.4	10.6	9.5*	10.8	10.8	14.8	6.5*	10.9	11.2	11.1
Mexican	2.2	2.9	1.8	2.2	3.1	1.8	2.9	1.2	1.5	2.0	2.0
Male	2.0	3.5	1.2	2.4	3.9	1.5	3.3	0.6*	1.0*	1.4	1.1*
Female	2.3	2.1	2.4	2.0	2.3	2.1	2.4	1.8	2.2	2.6	2.8
Not Hispanic or Latino	4.3	5.9	3.7	4.7	6.5	4.4	5.3	4.5	3.9	3.8	2.7
Male	3.6	6.9	2.5	5.4	8.0	3.5	5.1	3.7	2.5	2.3	1.9
Female	4.9	4.8	4.9	3.9	5.0	5.3	5.6	5.3	5.2	5.3	3.3
Region											
Northeast	4.5	6.5	3.8	5.1	7.4	4.4	5.5	4.6	3.8	4.2	2.3
Midwest	4.3	5.9	3.8	5.1	6.3	4.5	5.5	4.6	3.9	3.8	2.9
South	3.9	5.5	3.3	4.5	6.1	3.7	4.5	3.5	3.3	3.4	2.8
West	4.0	4.7	3.8	3.1	5.1	4.2	5.6	3.9	3.7	3.9	3.0
Ratio of Family Income to Poverty Threshold											
0 - .99	6.3	7.1	5.8	7.3	7.0	5.3	5.8	5.1	5.0	7.2	4.6
1.00 - 2.49	4.6	5.6	4.1	5.2	5.9	4.1	4.4	4.0	4.1	4.8	3.1
2.50 - 4.49	4.1	5.4	3.5	3.3	6.5	4.2	5.5	4.4	3.4	3.5	2.9
4.50 and above	3.6	5.4	3.2	3.0	6.3	3.9	5.9	2.6	3.5	3.3	1.9

*The estimate is unreliable because the relative standard error of the estimate is 30%-50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 40. (Corresponds to MMWR Table 7.) Estimated average annual attack percent for those reporting current asthma by age, sex, race, ethnicity, region and poverty level: National Health Interview Survey, United States, 2001-2004

	Persons with an Asthma Attack per 100 with Current Asthma										
	All ages	Children	Adults	Age							
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	15 - 19	20 - 24	25 - 34	35 - 64	65 +
Total	55.3	62.6	52.0	70.8	62.8	52.7	52.4	50.9	54.0	54.2	43.5
Sex											
Male	54.3	63.9	47.2	69.2	64.8	50.4	52.7	51.0	47.5	48.2	38.6
Female	56.0	60.8	54.6	73.1	59.7	54.2	52.0	50.9	57.7	57.2	46.0
Race											
Single race white	54.6	62.5	51.7	72.5	62.6	52.6	52.9	49.1	54.5	54.1	42.9
Male	53.4	64.0	47.0	71.3	64.7	50.9	55.1	48.4	48.2	48.0	38.2
Female	55.5	60.4	54.3	74.2	59.4	53.8	50.9	49.7	58.0	57.3	45.4
Single race black	57.1	62.2	53.2	69.1	63.7	52.4	44.2	59.8	55.1	53.2	49.0
Male	57.4	63.1	49.5	62.9	66.3	50.5	41.2	69.2	51.3	49.3	39.8
Female	56.8	61.0	55.0	76.6	59.2	53.7	47.5	55.3	56.9	54.7	52.4
Other races NTA	57.8	64.1	53.1	67.3	62.7	54.0	64.0	48.5	49.3	56.7	43.9
Male	56.1	64.7	45.6	71.9	63.1	47.2	57.3	43.2	38.6	49.5	44.2*
Female	59.5	63.0	57.9	57.5	62.1	59.8	73.1	51.8	56.5	61.3	43.7
Ethnicity											
Hispanic or Latino	55.7	57.6	54.1	61.2	55.6	52.3	56.4	43.2	53.6	57.5	55.1
Male	54.4	58.6	49.0	58.8	57.9	50.7	56.6	42.9	47.3	52.1	46.7
Female	56.8	56.3	57.0	64.9	52.4	53.5	56.0	43.3	57.0	60.3	59.9
Puerto Rican	64.2	64.0	64.4	80.2	60.7	60.3	59.2	55.0	64.9	64.7	73.9
Male	62.7	62.9	62.5	78.4	60.9	65.3	55.2	88.8	60.4	49.5	85.8
Female	65.4	65.5	65.4	83.8	60.5	57.6	62.6	32.8*	66.2	73.4	68.4
Mexican	50.3	52.5	48.4	52.3	50.8	47.3	56.7	30.7	48.7	52.9	48.0
Male	50.6	55.3	44.0	48.8	54.5	49.0	65.1	25.8*	44.9*	49.5	36.1*
Female	50.0	48.4	50.9	57.6	45.4	46.1	47.2	34.3*	50.9	54.9	54.4
Not Hispanic or Latino	55.2	63.5	51.8	72.9	64.0	52.7	51.8	51.7	54.1	53.9	42.8
Male	54.3	64.8	47.0	71.5	66.0	50.4	52.2	51.8	47.6	47.9	38.1
Female	55.9	61.6	54.4	74.9	61.0	54.3	51.5	51.6	57.8	56.9	45.2
Region											
Northeast	54.3	62.4	50.6	71.1	62.9	51.1	49.4	51.6	52.4	53.7	38.5
Midwest	55.7	63.5	52.3	74.9	64.4	51.9	50.9	51.6	53.0	55.4	42.4
South	55.5	62.1	52.4	68.0	62.8	52.2	50.5	51.6	53.8	54.2	46.8
West	55.3	62.6	52.4	70.3	60.8	55.9	60.6	48.4	57.1	53.2	44.2
Ratio of Family Income to Poverty Threshold											
0 - .99	58.9	62.6	56.8	68.7	61.4	54.5	49.7	53.0	62.0	61.0	49.9
1.00 - 2.49	56.4	62.9	53.2	72.5	62.6	52.1	50.9	49.1	55.0	58.6	43.0
2.50 - 4.49	57.3	67.5	52.8	77.7	68.1	54.9	59.6	58.2	50.3	53.9	45.5
4.50 and above	53.2	63.0	50.0	71.7	64.1	51.4	56.6	31.8	55.2	50.7	42.7

*The estimate is unreliable because the relative standard error of the estimate is 30%–50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 41. (Corresponds to MMWR Table 11.) Estimated average annual number of physician office visits with asthma as the first listed diagnosis by age, sex, race, and region: National Ambulatory Medical Care Survey, United States, 2001- 2004

	Number of Physician Office Visits for Asthma							
	All ages	Children	Adults	Age				
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	35 - 64	65 +
Total	12,609,000	5,173,000	7,435,000	1,979,000	2,640,000	2,048,000	3,863,000	2,079,000
Sex								
Male	5,474,000	3,001,000	2,473,000	1,279,000	1,538,000	762,000	1,248,000	647,000
Female	7,135,000	2,172,000	4,963,000	701,000	1,101,000	1,286,000	2,615,000	1,432,000
Race								
White	10,566,000	4,295,000	6,271,000	1,671,000	2,168,000	1,680,000	3,197,000	1,850,000
Male	4,544,000	2,525,000	2,020,000	1,087,000	1,307,000	581,000	975,000	594,000
Female	6,021,000	1,771,000	4,251,000	584,000	861,000	1,099,000	2,222,000	1,256,000
Black	1,322,000	509,000	813,000	183,000	280,000	158,000	569,000	132,000*
Male	506,000	258,000	248,000*	91,000*	139,000	**	197,000*	**
Female	817,000	251,000	566,000	93,000*	141,000*	92,000*	372,000	119,000*
Other races NTA	720,000	369,000	351,000*	125,000*	191,000*	210,000*	97,000*	**
Male	424,000	219,000	205,000*	101,000*	92,000*	115,000*	**	**
Female	296,000*	150,000*	**	**	100,000*	**	21,000*	**
Region								
Northeast	2,779,000	1,228,000	1,551,000	502,000	608,000*	388,000*	792,000*	488,000
Midwest	2,601,000	768,000	1,833,000	172,000*	450,000	407,000	1,048,000	525,000*
South	4,453,000	2,103,000	2,350,000	1,072,000	858,000	654,000	1,261,000	608,000*
West	2,775,000	1,075,000	1,700,000	234,000	723,000	599,000	761,000	458,000

*The estimate is unreliable because the relative standard error of the estimate is 30%–50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 42. (Corresponds to MMWR Table 12.) Estimated average annual rate of physician office visits with asthma as the first listed diagnosis, per 100 with current asthma, by age, sex, race, and region: National Ambulatory Medical Care Survey, United States, 2001-2004

	Physician Office Visits for Asthma per 100 with Current Asthma							
	All ages	Children	Adults	Age				
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	35 - 64	65 +
Total	62.5	83.5	53.2	171.9	68.8	35.4	52.7	100.3
Sex								
Male	64.0	82.6	50.3	188.1	65.7	32.0	51.0	92.5
Female	61.4	84.8	54.8	148.6	73.8	37.8	53.5	104.2
Race								
White	68.7	103.6	55.9	253.5	83.1	38.8	53.6	102.4
Male	71.4	105.6	50.9	278.6	83.5	32.8	48.4	96.3
Female	66.8	100.9	58.6	217.1	82.6	42.9	56.2	105.6
Black	42.0	37.8	45.1	55.5*	34.9	16.9*	62.9	74.9*
Male	37.1	32.4	43.6*	50.3*	27.6	**	77.*	**
Female	45.7	45.5	45.8	61.6*	47.5*	16.4*	57.3	92.6*
Other races NTA	43.6	52.3	37.1*	77.4*	45.1*	41.1*	20.8*	**
Male	51.5	48.7*	54.9*	92.4*	33.9*	**	**	**
Female	35.8*	58.7*	**	**	**	**	**	**
Region								
Northeast	66.4	94.5	53.7	220.1	74.1*	34.9*	49.1*	117.7
Midwest	51.1	50.0	51.5	56.*	50.3	25.5	59.6	96.8*
South	64.6	94.7	50.3	242.2	62.2	34.5	51.1	85.8*
West	69.4	94.2	59.5	134.0	97.7	50.8	50.9	112.5

*The estimate is unreliable because the relative standard error of the estimate is 30%–50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 43. (Corresponds to MMWR Table 16.) Estimated average annual number of outpatient department visits with asthma as the first listed diagnosis by age, sex, race, ethnicity and region: National Hospital Ambulatory Medical Care Survey, United States, 2001-2004

	Number of Outpatient Visits for Asthma							
	All ages	Children	Adults	Age				
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	35 - 64	65 +
Total	1,269,500	604,100	665,300	234,400	328,600	258,800	339,500	108,200
Sex								
Male	541,100	370,700	170,400	154,100	195,800	81,000	87,000	23,300*
Female	728,400	233,400	494,900	80,300	132,800	177,900	252,500	84,900
Race								
White	774,600	324,400	450,200	122,000	179,900	159,300	238,100	75,300
Male	318,300	188,400	129,900	80,400	99,500	58,300	62,100	17,900*
Female	456,200	135,900	320,300	41,500	80,400	100,900	176,000	57,300
Black	439,100	249,800	189,300	97,300	136,500	88,100	87,200	30,100*
Male	196,400	163,000	33,400	60,100	92,900	19,800	20,000*	**
Female	242,700	86,700	155,900	37,200	43,500	68,300	67,200	26,500*
Other races NTA	55,800	30,000*	25,800	15,200*	12,200*	11,500*	14,200*	**
Male	26,300*	19,200*	7,100*	**	3,300*	**	**	**
Female	29,500	10,800*	18,700*	**	8,900*	**	9,300*	**
Ethnicity								
Hispanic or Latino	225,400	78,300	147,100*	24,500	47,600	36,400	79,900*	37,100*
Male	77,400	39,500	37,900*	15,600	22,200	6,800*	**	**
Female	148,000	38,800	109,300	8,800	25,400	29,500	59,300*	25,000*
Not Hispanic or Latino	1,044,100	525,900	518,200	209,900	281,000	222,500	259,700	71,000
Male	463,700	331,200	132,500	138,400	173,600	74,100	66,400	11,200*
Female	580,300	194,700	385,700	71,500	107,400	148,300	193,300	59,800
Region								
Northeast	374,300	121,900	252,400	42,600	69,800	74,400	151,600	35,900*
Midwest	313,200	172,000	141,200	65,300	94,600	75,300	51,000	27,100*
South	428,600	235,200	193,400	94,500	129,400	64,100	102,800	37,600*
West	153,400	75,100	78,400	32,000*	34,700	45,000	34,100	**

*The estimate is unreliable because the relative standard error of the estimate is 30%–50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 44. (Corresponds to MMWR Table 17.) Estimated average annual rate of outpatient department visits with asthma as the first listed diagnosis, per 100 with current asthma by age, sex, race, and region: National Hospital Ambulatory Medical Care Survey - United States, 2001- 2004

	Outpatient Visits for Asthma per 100 with Current Asthma							
	All ages	Children	Adults	Age				
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	35 - 64	65 +
Total	6.3	9.8	4.8	20.4	8.6	4.5	4.6	5.2
Sex								
Male	6.3	10.2	3.5	22.7	8.4	3.4	3.6	3.33*
Female	6.3	9.1	5.5	17.0	8.9	5.2	5.2	6.2
Race								
White	5.0	7.8	4.0	18.5	6.9	3.7	4.0	4.2
Male	5.0	7.9	3.3	20.6	6.4	3.3	3.1	2.91*
Female	5.1	7.7	4.4	15.4	7.7	3.9	4.5	4.8
Black	13.9	18.6	10.5	29.4	17.0	9.4	9.6	17.11*
Male	14.4	20.5	5.9	33.4	18.4	5.3	7.81*	**
Female	13.6	15.7	12.6	24.7	14.7	12.1	10.3	20.65*
Other races NTA	3.4	4.25*	2.7	**	2.87*	2.25*	3.06*	**
Male	3.2*	4.27*	1.9*	**	1.23*	**	**	**
Female	3.6	4.22*	3.27*	**	5.75*	**	3.24*	**
Ethnicity								
Hispanic or Latino	**	**	**	**	**	**	**	**
Male	**	**	**	**	**	**	**	**
Female	**	**	**	**	**	**	**	**
Not Hispanic or Latino	**	**	**	**	**	**	**	**
Male	**	**	**	**	**	**	**	**
Female	**	**	**	**	**	**	**	**
Region								
Northeast	8.9	9.4	8.7	18.6	8.5	6.7	9.4	8.66*
Midwest	6.1	11.2	4.0	21.3	10.6	4.7	2.9	4.99*
South	6.2	10.6	4.1	21.4	9.4	3.4	4.2	5.3*
West	3.8	6.6	2.7	18.4*	4.7	3.8	2.3	**

*The estimate is unreliable because the relative standard error of the estimate is 30%–50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 45. (Corresponds to MMWR Table 21.) Estimated average annual number of emergency department visits with asthma as the first listed diagnosis by age, sex, race, and region: National Hospital Ambulatory Medical Care Survey, United States, 2001-2004

	Number of Emergency Department Visits for Asthma										
	All ages	Children	Adults	Age							
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	15 - 19	20 - 24	25 - 34	35 - 64	65 +
Total	1,788,600	711,200	1,077,500	309,700	320,400	547,700	128,100	154,700	264,900	508,200	102,700
Sex											
Male	809,100	427,200	382,000	191,300	196,200	224,700	55,300	63,300	106,200	158,600	38,400
Female	979,500	284,000	695,500	118,400	124,300	323,000	72,800	91,400	158,800	349,600	64,300
Race											
White	1,053,300	378,500	674,900	164,500	167,700	336,600	73,800	105,000	157,800	317,900	66,700
Male	436,700	213,900	222,800	102,700	93,400	120,000	24,100	38,700	57,200	94,100	26,600
Female	616,600	164,600	452,000	61,800	74,300	216,600	49,700	66,300	100,600	223,900	40,100
Black	684,100	306,500	377,600	131,400	143,700	197,600	47,500	48,100	102,000	179,200	32,200
Male	345,500	196,200	149,300	80,400	95,500	98,300	27,500	23,700	47,200	60,700	10,600
Female	338,600	110,400	228,200	50,900	48,200	99,300	20,000	24,500	54,800	118,500	21,600
Other races NTA	51,200	26,100	25,100	13,800*	9,000*	13,500*	6,800*	**	5,100*	11,100	3,800*
Male	26,900	17,100	9,800*	8,100*	7,300*	6,400*	3,700*	**	**	3,800*	**
Female	24,300	9,000*	15,300	5,700*	**	7,100*	**	**	3,300*	7,200	2,500*
Ethnicity											
Hispanic or Latino	252,000	130,100	122,000	62,600	53,600	66,800	16,400	10,700*	39,700	58,200	10,900*
Male	125,600	78,800	46,700	41,500	31,300	30,200	6,300*	2,600*	21,300	18,000	4,500*
Female	126,500	51,200	75,200	21,100	22,200	36,600	10,100*	8,100*	18,400	40,200	6,400*
Not Hispanic or Latino	1,536,600	581,100	955,500	247,100	266,900	480,900	111,700	144,000	225,200	450,000	91,800
Male	683,600	348,300	335,200	149,700	164,800	194,500	49,000	60,700	84,800	140,700	33,900
Female	853,000	232,700	620,300	97,300	102,100	286,400	62,700	83,300	140,400	309,300	57,900
Region											
Northeast	456,200	152,300	303,900	56,800	79,800	132,800	29,700	34,800	68,300	155,100	31,600
Midwest	433,000	183,700	249,300	77,900	77,000	153,000	42,300	42,000	68,700	111,400	13,700*
South	655,200	289,100	366,100	139,600	120,600	178,700	41,800	52,500	84,400	181,200	35,100
West	244,200	86,100	158,100	35,300	43,100	83,100	14,400*	25,400	43,400	60,500	22,200

*The estimate is unreliable because the relative standard error of the estimate is 30%–50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 46. (Corresponds to MMWR Table 22.) Estimated average annual rate of emergency department visits, with asthma as the first listed diagnosis, per 100 with current asthma, by age, sex, race, and region: National Hospital Ambulatory Medical Care Survey, United States, 2001-2004

	Emergency Department Visits for Asthma per 100 with Current Asthma										
	All ages	Children	Adults	Age							
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	15 - 19	20 - 24	25 - 34	35 - 64	65 +
Total	8.9	11.5	7.7	26.9	8.4	9.5	6.7	10.3	11.1	6.9	5.0
Sex											
Male	9.5	11.8	7.8	28.1	8.4	9.4	5.9	10.8	12.4	6.5	5.5
Female	8.4	11.1	7.7	25.1	8.3	9.5	7.6	10.0	10.4	7.2	4.7
Race											
White	6.9	9.1	6.0	25.0	6.4	7.8	5.3	9.2	8.8	5.3	3.7
Male	6.9	8.9	5.6	26.3	6.0	6.8	3.6	8.4	8.8	4.7	4.3
Female	6.8	9.4	6.2	23.0	7.1	8.5	6.8	9.8	8.7	5.7	3.4
Black	21.7	22.8	20.9	39.8	17.9	21.1	13.9	18.7	30.4	19.8	18.3
Male	25.4	24.7	26.3	44.7	18.9	26.4	15.4	28.2	42.6	23.7	22.17*
Female	18.9	20.0	18.5	33.8	16.3	17.7	12.2	14.1	24.4	18.2	16.9
Other races NTA	3.1	3.7	2.6	8.52*	2.13*	2.64*	4.*	**	2.1*	2.4	4.2*
Male	3.3	3.8	2.62*	7.37*	2.69*	2.7*	**	**	**	2.19*	**
Female	2.9	3.53*	2.7	**	**	2.55*	**	**	2.3*	2.5	**
Ethnicity											
Hispanic or Latino	**	**	**	**	**	**	**	**	**	**	**
Male	**	**	**	**	**	**	**	**	**	**	**
Female	**	**	**	**	**	**	**	**	**	**	**
Not Hispanic or Latino	**	**	**	**	**	**	**	**	**	**	**
Male	**	**	**	**	**	**	**	**	**	**	**
Female	**	**	**	**	**	**	**	**	**	**	**
Region											
Northeast	10.9	11.7	10.5	24.9	9.7	12.0	7.7	11.8	15.9	9.6	7.6
Midwest	8.5	12.0	7.0	25.4	8.6	9.6	8.0	9.5	11.0	6.3	2.53*
South	9.5	13.0	7.8	31.6	8.7	9.4	6.7	11.6	10.3	7.3	4.9
West	6.1	7.5	5.5	20.2	5.8	7.1	3.9*	8.3	8.7	4.0	5.5

*The estimate is unreliable because the relative standard error of the estimate is 30%–50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 47. (Corresponds to MMWR Table 26.) Estimated average annual number of hospital discharges with asthma as the first listed diagnosis by age, sex, race, and region: National Hospital Discharge Survey, United States, 2001- 2004

	Number of Hospital Discharges for Asthma										
	All ages	Children	Adults	Age							
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	15 - 19	20 - 24	25 - 34	35 - 64	65 +
Total	502,200	203,000	299,200	117,200	75,500	56,300	14,600	12,700	29,100	161,000	92,300
Sex											
Male	205,300	128,800	76,500	75,900	48,400	19,800	6,400	3,900	9,500	38,800	22,400
Female	297,000	74,300	222,700	41,300	27,100	36,500	8,100	8,800	19,600	122,200	69,900
Race											
White	248,700	93,000	155,800	56,400	31,800	25,600	5,900	5,700	14,100	79,800	55,100
Male	96,300	58,300	38,100	36,000	20,000	8,700	3,000	1,200	4,400	18,300	13,400
Female	152,400	34,700	117,700	20,400	11,800	17,000	2,900	4,400	9,600	61,400	41,800
Black	129,200	64,800	64,400	33,800	27,200	16,500	5,600	2,900	8,000	40,200	11,500
Male	59,600	41,700	17,800	22,200	17,800	6,300	2,300	1,300*	2,800	10,400	2,800
Female	69,600	23,100	46,500	11,600	9,300	10,200	3,300	1,600	5,300	29,800	8,700
Other races NTA	27,000	12,200	14,700	7,700	4,200	3,000	500	1,000*	1,500*	7,600	4,400
Male	12,500	7,700	4,800	4,900	2,700	1,100*	200	**	**	2,500	1,300
Female	14,500	4,500	10,000	2,800	1,500	1,900	300*	700*	800*	5,100	3,200
Unknown Race	97,400	33,000	64,400	19,200	12,400	11,200	2,600	3,100	5,500	33,400	21,200
Male	36,900	21,000	15,800	12,700	7,900	3,700	1,000*	1,200*	1,600	7,600	5,000
Female	60,500	12,000	48,500	6,500	4,500	7,500	1,600	2,000	3,900	25,900	16,200
Region											
Northeast	127,100	47,100	80,000	27,300	16,600	16,600	5,000	3,500	8,100	42,400	24,200
Midwest	112,400	42,400*	70,000	20,800*	18,700*	13,000	3,800*	2,600	6,600	39,100	20,800
South	172,300	69,200	103,100	40,500	26,400	17,600	3,300	4,800	9,600	56,200	31,500
West	90,400	44,400*	46,100	28,500*	13,700	9,100	2,500*	1,800*	4,800	23,300	15,800

*The estimate is unreliable because the relative standard error of the estimate is 30%–50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 48. (Corresponds to MMWR Table 27.) Estimated average annual number of hospital discharges, with asthma as the first listed diagnosis, per 100 with current asthma, by age, sex, race, and region: National Hospital Discharge Survey, United States, 2001- 2004

	Hospital Discharges for Asthma per 100 with Current Asthma										
	All ages	Children	Adults	Age							
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	15 - 19	20 - 24	25 - 34	35 - 64	65 +
Total	2.5	3.3	2.2	10.1	1.9	1.0	0.8	0.8	1.2	2.2	4.7
Sex											
Male	2.4	3.6	1.6	11.1	2.1	0.8	0.7	0.6	1.1	1.6	3.4
Female	2.6	2.8	2.5	8.6	1.7	1.1	0.8	0.9	1.3	2.5	5.4
Race											
White	1.6	2.2	1.4	8.9	1.2	0.6	0.4	0.5	0.8	1.4	3.2
Male	1.5	2.5	1.0	9.4	1.3	0.5	0.5	0.2	0.7	0.9	2.3
Female	1.7	2.0	1.6	8.1	1.1	0.7	0.4	0.7	0.8	1.6	3.7
Black	4.1	4.8	3.6	9.6	3.5	1.8	1.7	1.1	2.3	4.5	7.0
Male	4.6	5.5	3.3	11.9	3.8	1.8	1.4	1.6*	2.5*	4.1	7.0
Female	3.8	4.0	3.8	7.0	3.1	1.8	1.9	0.9	2.3	4.6	7.0
Other races NTA	1.6	1.7	1.6	4.4	1.0	0.6	0.3*	.9*	0.6*	1.8	4.8
Male	1.5	1.7	1.2	4.4	1.0	0.4*	0.2*	**	**	1.5	3.3*
Female	1.8	1.7	1.8	4.5*	1.0	0.7	0.4*	1.1*	0.6*	2.0	5.9*
Unknown Race	na	na	na	na	na	na	na	na	na	na	na
Male	na	na	na	na	na	na	na	na	na	na	na
Female	na	na	na	na	na	na	na	na	na	na	na
Region											
Northeast	3.0	3.5	2.7	11.3	1.9	1.5	1.2	1.1	1.9	2.6	5.8
Midwest	2.2	2.8*	2.0	6.7*	2.1*	0.8	0.8*	0.6	1.0	2.2	4.1
South	2.5	3.1	2.2	9.2	1.9	0.9	0.5	1.0	1.1	2.3	4.8
West	2.4	3.9*	1.7	16.8*	1.8	0.8	0.7*	0.7*	1.0	1.7	4.2

*The estimate is unreliable because the relative standard error of the estimate is 30%–50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.

Table 49. (Corresponds to MMWR Table 31.) Average annual number of deaths with asthma as the underlying cause by age, sex, race, ethnicity, and region: National Vital Statistics System, United States, 2001-2004

	Number of deaths										
	All ages	Children	Adults	Age							
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	15 - 19	20 - 24	25 - 34	35 - 64	65 +
Total	4,015	194	3,821	41	108	372	74	84	215	1,534	1,960
Sex											
Male	1,452	124	1,328	29	65	215	48	48	119	585	558
Female	2,563	70	2,493	12	43	157	26	36	96	949	1,402
Race											
White	2,812	83	2,729	19	46	194	33	45	117	933	1,621
Male	912	51	861	14	26	102	21	25	57	335	435
Female	1,900	32	1,868	5	20	92	12	20	60	598	1,186
Black	1,043	103	940	20	59	168	37	36	95	545	253
Male	461	69	392	14	38	106	25	21	60	220	84
Female	582	34	548	6	21	62	12	15	35	325	169
Other	160	9	151	2	4	12	4	4	4	58	86
Male	80	5	75	1	2	8	3	3	2	31	39
Female	80	4	76	1	2	4	1	1	2	27	47
Ethnicity											
Hispanic or Latino	284	23	261	9	12	35	6	8	23	125	106
Male	114	15	99	6	7	22	4	5	14	49	31
Female	170	8	162	3	5	13	2	3	9	76	75
Not Hispanic or Latino	3,598	166	3,432	30	94	324	69	75	181	1,359	1,791
Male	1,338	108	1,230	23	58	193	45	44	105	537	528
Female	2,260	58	2,202	7	36	131	24	31	76	822	1,263
Region											
Northeast	766	38	728	7	24	70	10	16	44	318	348
Midwest	931	53	878	10	28	103	24	23	57	314	478
South	1,404	69	1,336	16	39	132	26	31	76	569	648
West	913	34	879	8	18	67	15	14	38	318	503

Table 50. (Corresponds to MMWR Table 32.) Average annual number of deaths, with asthma as the underlying cause, per 10,000 with current asthma, by age, sex, race, ethnicity, and region: National Vital Statistics System, United States, 2001-2004

	Number of deaths										
	All ages	Children	Adults	Age							
	Total	Age <18	Age 18+	0 - 4	5 - 14	15 - 34	15 - 19	20 - 24	25 - 34	35 - 64	65 +
Total	2.0	0.3	2.7	0.4	0.3	0.6	0.4	0.6	0.9	2.1	9.5
Sex											
Male	1.7	0.3	2.7	0.4	0.3	0.9	0.5	0.8	1.4	2.4	8.0
Female	2.2	0.3	2.8	0.3	0.3	0.5	0.3	0.4	0.6	1.9	10.2
Race											
White	1.8	0.2	2.4	0.3	0.2	0.4	0.2	0.4	0.6	1.6	9.0
Male	1.4	0.2	2.2	0.4	0.2	0.6	0.3	0.5	0.9	1.7	7.1
Female	2.1	0.2	2.6	0.2	0.2	0.4	0.2	0.3	0.5	1.5	10.0
Black	3.3	0.8	5.2	0.6	0.7	1.8	1.1	1.4	2.8	6.0	14.4
Male	3.4	0.9	6.9	0.8	0.8	2.8	1.4	2.5	5.4	8.6	17.6
Female	3.3	0.6	4.4	0.4	0.7	1.1	0.7	0.9	1.6	5.0	13.2
Other	1.0	0.1	1.6	.1*	0.1	0.2	0.2	0.4	0.2	1.3	9.5
Male	1.0	0.1	2.0	**	.1*	0.3	.3*	.8*	.2*	1.8	11.3
Female	1.0	0.2	1.3	**	.1*	0.1	**	**	.1*	0.9	8.4
Ethnicity											
Hispanic or Latino	1.4	0.3	2.4	0.4	0.2	0.6	0.3	0.6	0.9	2.3	9.5
Male	1.2	0.3	2.4	0.5	0.2	0.8	0.3	1.0	1.6	2.7	7.5
Female	1.6	0.2	2.3	.4*	0.2	0.4	.2*	.4*	0.6	2.1	10.6
Not Hispanic or Latino	2.0	0.3	2.7	0.3	0.3	0.6	0.4	0.5	0.9	2.0	9.1
Male	1.8	0.3	2.7	0.4	0.3	0.9	0.5	0.8	1.4	2.4	8.0
Female	2.1	0.3	2.6	0.2	0.3	0.4	0.3	0.4	0.6	1.8	9.7
Region											
Northeast	1.8	0.3	2.5	0.3	0.3	0.6	0.3	0.5	1.0	2.0	8.4
Midwest	1.8	0.3	2.5	0.3	0.3	0.6	0.5	0.5	0.9	1.8	8.8
South	2.0	0.3	2.9	0.4	0.3	0.7	0.4	0.7	0.9	2.3	9.1
West	2.3	0.3	3.1	0.5	0.2	0.6	0.4	0.5	0.8	2.1	12.4

*The estimate is unreliable because the relative standard error of the estimate is 30%–50%. For missing estimates (**), the relative standard error of the estimate exceeded 50% and the estimate was suppressed. All other relative standard errors are <30%.