

TABLE 1
THE 20 MOST FREQUENTLY REPORTED SALMONELLA SEROTYPES
FROM HUMAN SOURCES REPORTED TO CDC IN 1995 AND FROM
NONHUMAN SOURCES REPORTED TO CDC AND USDA IN 1995

HUMAN 1995				NONHUMAN 1995			
RANK	SEROTYPE	NUMBER	PERCENT	RANK	SEROTYPE	NUMBER	PERCENT
1	ENTERITIS	10201	24.7	1	ENTERITIS	3596	17.1
2	TYPHIMURIUM *	9702	23.5	2	TYPHIMURIUM *	3505	16.7
3	NEWPORT	2566	6.2	3	HEIDELBERG	2299	10.9
4	HEIDELBERG	2095	5.1	4	DERBY	918	4.4
5	HADAR	812	2.0	5	KENTUCKY	832	4.0
6	JAVITANA	758	1.8	6	BRANDENBURG	772	3.7
7	MUENCHEN	754	1.8	7	HADAR	767	3.7
8	MONTEVIDEO	685	1.7	8	MONTEVIDEO	701	3.3
9	AGONA	683	1.7	9	CHOLERAESUIS **	614	2.9
10	THOMPSON	625	1.5	10	MUENSTER	560	2.7
11	ORANIENBURG	595	1.4	11	AGONA	557	2.7
12	BRAENDERUP	588	1.4	12	SENFTENBERG	485	2.3
13	POONA	531	1.3	13	ANATUM	373	1.8
14	INFANTIS	521	1.3	14	BREDENEY	358	1.7
15	STANLEY	481	1.2	15	SCHWARZENGROUND	348	1.7
16	SAINTPAUL	467	1.1	16	MBANDAKA	323	1.5
17	TYPHI	442	1.1	17	NEWPORT	257	1.2
18	BERTA	367	0.9	18	BRAENDERUP	214	1.0
19	BRANDENBURG	284	0.7	19	DUBLIN	197	0.9
20	JAVA	268	0.7	20	ISTANBUL	196	0.9
SUB TOTAL		33425	81.1	17872		85.1	
TOTAL		41222		21008			

* TYPHIMURIUM INCLUDES VAR. COPENHAGEN
** CHOLERAESUIS INCLUDES VAR. KUNZENDORF