USGS
science for a changing world

# Colorado's Ancient Trees

Some of the oldest trees in North America live in rugged and remote mountain areas in Colorado. The oldest known living Rocky Mountain bristlecone pines *(Pinus aristata)*, which are about 2,500 years old, are found in the State. These trees are the oldest known trees in the Rocky Mountains. Five living bristlecone pines in Colorado are estimated to be 2,000 or more years old. Three of those trees are shown on this poster.

Also shown on the poster are an old limber pine *(Pinus flexilis)* and an old ponderosa pine *(Pinus ponderosa)*, both of which are among the oldest known trees of their species in North America. All of these ancient trees are on National Forest land. The exact locations of the trees are not revealed on this poster, in order to protect the trees. If too many people visit such areas, the pristine ecology of the sites will be harmed by trampling of fragile alpine plants and lichen, the loss of which will lead to visual degradation of the sites, soil erosion, and harm to the ancient trees.

*Limber pine (Pinus flexilis), about 1,600 years old, growing on a bedrock outcrop.*



*In the cool and relatively dry environment near upper treeline, the wood of dead bristlecone pines can persist for hundreds to thousands of years. This wood can be used by scientists to extend the tree-ring record far back in time.*



*(ABOVE) Rocky Mountain bristlecone pine (Pinus aristata), about 2,000 years old, growing among large lichen-covered granitic boulders. Several species of lichen are present; their colors are lush shades of green, yellow, gray, and orange. Younger trees are in the background.*

*(LEFT) Exposed dead roots of this ponderosa pine (Pinus ponderosa) about 800 years old, record how much erosion of the granitic soil has taken place during its lifetime.*

Because of the harsh alpine environment in which these trees grow, they are stunted, attaining heights of only 10-25 feet (3-8 meters). Longevity of these trees seems to be related to their ability to adapt to adverse environmental conditions. The ancient trees are partly protected from forest fires because the rocky ground surrounding the trees supports only sparse amounts of low plants and shrubs that could serve as fuel for a fire, and because the trees usually grow far apart from each other.

The trunk of each of these ancient living trees is exposed wood except for thin strips of bark that connect a few living branches to sturdy roots. Through the centuries, wind-driven snow and sand gradually erode and sculpt the exposed wood into the forms we see today. Close inspection of the wood reveals intricate textures, such as furrows and ridges. Wood surfaces are silvery gray and shades of amber to dark brown, which contrast with the trees' green needles.

To determine the age of an ancient tree, without harming it, a tiny core sample is carefully collected from the tree using a special tool called an increment borer. Scientists examine the core sample under a microscope and carefully count the number of rings; one ring forms each year. Scientists use a technique called crossdating to match narrow and wide rings in the core sample with narrow and wide rings in core samples from nearby trees. Crossdating is necessary to make sure that occasional very narrow or missing rings are not overlooked in each core sample. Rings can be very narrow or missing in some trees because of abnormally dry environmental conditions or because of tree injury. For example, trees studied at an arid site in central Colorado have very narrow or missing rings for the year 1963, a year in which the spring and early summer were unusually dry.

*Alpine pennyroyr (Primula angustifolia), only a few inches tall, grows in gravelly granitic soil near upper treeline among bristlecone pines.*

*Fully open cone on branch of Rocky Mountain bristlecone pine has released its crop of seeds. Current year's needles on the tip of the branch have broken through their protective sheath but have not completely spread out.*

*Unopened cones on branch of Rocky Mountain bristlecone pine. These cones will open, turn brown, and release their seeds. Note the bristles on the cones, a feature for which the tree is named ("bristlecone"). The needles have numerous white resin specks, which are characteristic of this tree species. Current year's needles at the tip of the branch have not emerged from their protective sheath.*

*(LEFT) Limber needlepine (Pinus flexilis) grows out of a crack in a vertical face of weathered granitic bedrock. Bristlecone pines grow nearby.*

*(ABOVE) Alpine sunflower (Hymenoxys grandiflora) also known as "Old man of the mountain," growing in a bristlecone pine grove at about 12,500 feet (3,800 meters) altitude.*

*Magnified view of annual rings of a Douglas-fir (Pseudotsuga menziesii), in central Colorado. Note that the 1963 annual ring is much narrower than surrounding rings, probably due to the unusually dry spring and early summer that occurred in 1963 in central Colorado. The ring widths of trees that grow on semi-arid sites are influenced by the amount of precipitation the trees receive. This enables researchers to crossdate precipitation recorded in weather stations with the measured ring widths of trees, and then to reconstruct records of precipitation extending hundreds to thousands of years before the present. The black dots are the A.D. 1960 and 1970 rings on pencil notations made by a researcher to mark these rings for future reference.*

*Each tree ring is made up of earlywood (EW) and latewood (LW), which together constitute one year's growth. Earlywood is composed of relatively light-colored, thin-walled cells that form during early to mid-summer. Latewood is composed of dark, thick-walled cells that form during late summer. As cells are formed in the winter and spring dormant season, trees in Colorado are dormant at this time of the year. Dormancy protects the trees from temperatures that occasionally dip below freezing during the winter and spring in the mountains of Colorado. About 30 times enlarged.*

*Leached cemented outcrop of fractured volcanic rock on a bristlecone pine grove above 11,000 feet (3,350 meters) altitude.*

*A stout, rocky talus slope has been the home of this Rocky Mountain bristlecone pine since about 450 B.C., making this tree about 2,450 years old. The contorted limb at the lower left has living branches; the tree's trunk as well as its top are dead. Younger living trees are in the background.*

*For about 2,200 years, this Rocky Mountain bristlecone pine has lived on this rocky talus slope. The mostly treeless basin in the distance is about 1,500 feet (460 meters) below. The age of this tree and the slow growing lichens that cover the rocks provide evidence that this slope has remained stable for over millennia. Stunted bristlecone pine on the left (its gray trunk is visible at the lower left) is taller than its ancient neighbor, but it is only about 300 years old.*

*(ABOVE) Bristlecone pine grove high up in the Colorado Front Range. The two trees in the foreground are more than 1,400 years old. Kate Ally (alk@bmars.com) grew up on the grassily greener scene.*

*(LEFT) View from a grove of bristlecone pines and Engelmann spruce to distant slope that has scattered dead trees at treeline. Top of distant ridge is about 12,200 feet (3,720 meters) altitude.*

**The World's Oldest Known Trees**

| Common name (scientific name) | Estimated age (years) | Location |
|---|---|---|
| Great Basin bristlecone pine (*Pinus longaeva*) | 4,800-4,900 | California and Nevada, U.S.A. |
| Alerce (*Fitzroya cupressoides*) | 3,620 | Chile |
| Giant Sequoia (*Sequoiadendron giganteum*) | 3,500-3,900 | California, U.S.A. |
| Rocky Mountain bristlecone pine (*Pinus aristata*) | 2,900 | Colorado, U.S.A. |
| Coast Redwood (*Sequoia sempervirens*) | 2,200 | California, U.S.A. |
| Foxtail pine (*Pinus balfouriana*) | 2,300 | California, U.S.A. |
| Western juniper (*Juniperus occidentalis*) | 2,300 | California, U.S.A. |
| Subalpine larch (*Larix lyallii*) | 1,920 | Alberta, Canada |
| Rocky Mountain juniper (*Juniperus scopulorum*) | 1,890 | New Mexico, U.S.A. |
| Limber pine (*Pinus flexilis*) | 1,470-1,600 | Colorado and New Mexico, U.S.A. |
| Baldcypress (*Taxodium distichum*) | 1,400 | North Carolina, U.S.A. |
| Douglas fir (*Pseudotsuga menziesii*) | 1,350 | British Columbia, Canada |
| Huon pine (*Lagarostrobus franklinii*) | 1,000 | Tasmania, Australia |
| Northern white cedar (*Thuja occidentalis*) | 1,040 | Ontario, Canada |
| Pinyon pine (*Pinus edulis*) | 980-1,000 | Utah, U.S.A. |
| Whitebark pine (*Pinus albicaulis*) | 880-890 | Alberta, Canada |
| Ponderosa pine (*Pinus ponderosa*) | 680-850 | Utah, U.S.A. |
| Engelmann spruce (*Picea engelmannii*) | 850-860 | Colorado, U.S.A. |

Text and photographs by Ed. Brunswig
Design and digital layout by Carol Quesenberry

Lorna Carter, Paul Carter, Diane Wirth, Gene Hills, Robert S. Thompson, and Ruth Ann Lucas provided thoughtful reviews of this poster. Alice Hunter to C.W. Ferguson, A.C. LaMarche, and H.E. Wrangham.

Photograph of ponderosa pine taken by UMS photomicrograph taken 1999, all others taken 1988-93.

U.S. Department of the Interior
U.S. Geological Survey
*Printed in cooperation with the*
U.S. Department of Agriculture, Forest Service