

# IMMUNIZATION Follow-up



SAFER • HEALTHIER • PEOPLE™

DATE \_\_\_\_\_ / \_\_\_\_\_ / \_\_\_\_\_

Today, \_\_\_\_\_

received vaccines to prevent:

- Diphtheria, Tetanus and Pertussis
- Diphtheria and Tetanus
- Haemophilus Influenzae* Type B (Hib)
- Hepatitis A
- Hepatitis B
- Influenza
- Measles, Mumps and Rubella
- Pneumococcal Disease
- Meningococcal Disease
- Polio
- Varicella (Chickenpox)
- Other \_\_\_\_\_

## YOUR CHILD'S NEXT IMMUNIZATION IS DUE:

### GENERAL INSTRUCTION/ REACTIONS

It is common for children to experience some discomfort from today's vaccine(s).

- Soreness, redness, swelling, tenderness where shot is given
- Fussiness, tiredness, or poor appetite
- Fever (usually low grade)

For relief, you may

- apply ice for first 24 hours
- administer \_\_\_\_\_ every 4 to 6 hours

Dosage:

DIPHTHERIA ■ TETANUS ■ PERTUSSIS ■ MEASLES ■ MUMPS ■ RUBELLA ■ POLIO ■ HIB ■ HEPATITIS A & B ■ VARICELLA ■ PNEUMOCOCCAL DISEASE ■ INFLUENZA ■ MENINGOCOCCAL DISEASE

In rare cases, moderate or severe reactions to vaccines can occur.  
This will usually be within a few minutes to a few hours after the vaccine.

### SHOULD YOUR CHILD EXPERIENCE

- difficulty breathing
- hoarseness or wheezing
- swelling of the throat
- weakness
- fast heart beat
- dizziness
- hives

CONTACT THIS OFFICE  
IMMEDIATELY AT:

### FOR MORE INFORMATION

[www.cdc.gov/nip](http://www.cdc.gov/nip)  
or call

**800-CDC-INFO (232-4636)**  
In English / En español — 24/7

Reactions to any vaccine should be reported to the  
Vaccine Adverse Event Reporting System (VAERS)  
**800-822-7967**

Get an instant list of recommended  
childhood vaccinations with dates by visiting  
[www.cdc.gov/nip/kidstuff](http://www.cdc.gov/nip/kidstuff)