

2002

Fernald Environmental Management Project Annual Illness and Injury Surveillance Report



Fernald Environmental Management Project 2002 Illness and Injury Surveillance Report

Questions or comments about this report or the Illness and Injury Surveillance Program may be directed to:

E-mail:

Dr. Cliff Strader at cliff.strader@eh.doe.gov
or Dr. Bonnie Richter at bonnie.richter@eh.doe.gov

or direct letters to:

United States Department of Energy
Office of Epidemiology and Health Surveillance
EH-53/270 Corporate Square Building
1000 Independence Avenue, SW
Washington, DC 20585-0270

Additional information about the Department of Energy's Office of Epidemiology and Health Surveillance, the Illness and Injury Surveillance Program, and annual reports for DOE sites participating in this program can be found at:

www.eh.doe.gov/health/epi/surv

ACKNOWLEDGEMENT

LifeART images copyright 2000 Lippincott Williams & Wilkins. All rights reserved.

Fernald Environmental Management Project 2002 Illness and Injury Surveillance Report

At A Glance

The number of absences and lost calendar days has decreased from 2001. A total of 17,005 calendar days of work (9,523 days for men and 7,482 days for women) was lost at Fernald in 2002 due to reported illness or injury, about a 10 percent decrease. Absences dropped from 440 in 2001 to 417 in 2002.

Among women, 52 percent of all reported diagnoses were due to conditions of the muscles and skeleton (18 percent), psychological disorders (12 percent), injuries (11 percent), and unspecified symptoms (11 percent).

Among men, 54 percent of all reported diagnoses were due to muscles and skeleton conditions (23 percent), injuries (21 percent), and digestive conditions (10 percent).

The increase seen in the 2000 age-adjusted rate for all diagnoses combined continued through 2002 among women, but the rate decreased among men.

Service/Security/Craft and Repair workers had the highest rate of OSHA events for both men and women in 2002.

The number of OSHA events more than tripled from 2001 to 2002, with 28 events in 2002 compared with only 8 in 2001. An overall decline in the rate among men in the Craft and Repair group was observed between 1994 and 2001, but the trend was broken by a sharp increase in 2002. In summary, we noted no significant changes in the injury rates from OSHA-recordable events from 1993 to 2002.

Introduction	1	Most Frequently Reported Diagnoses by Job Category and Gender	9
Site Overview	2		
The Fernald Work Force – 2002	3	Rates of Disease Occurrence	9
The Work Force by Gender and Age	3	Rates for All Illnesses and Injuries Combined by Job Category, Gender, and Age.....	10
The Work Force by Job Category and Gender	3	Rates for Selected Diagnostic Categories by Job Category, Gender, and Age	10
Number and Length of Absences	4	Time Trends	12
Absence Rate by Gender and Age	4	Age-Adjusted Rates for All Diagnoses Combined Among Women and Men from 1993 to 2002.....	14
Number of Days Absent by Gender and Age	5	Age-Adjusted Rates for Selected Diagnostic Categories Among Women and Men from 1993 to 2002.....	15
Absence Rate by Job Category and Gender	5	Age-Adjusted Rates for All Diagnoses Combined Among Women and Men by Job Category from 1993 to 2002	16
Average Duration of Absence by Job Category and Gender	6		
Diagnostic Categories	6	Sentinel Health Events for Occupations	17
Number of Diagnoses and Lost Calendar Days by Diagnostic Category (Categorized by ICD-9-CM) and Gender	7	Characteristics of SHEOs by Gender	17
Common Diagnoses Among Female Workers in 2002.....	8		
Common Diagnoses Among Male Workers in 2002.....	8	Disabilities Among Active Workers	17

Deaths Among Active Workers 17

OSHA-Recordable Events..... 18

OSHA-Recordable Events by Gender and Age 18

OSHA-Recordable Events by Job Category and Gender 18

OSHA-Recordable Diagnoses by Diagnostic Category and Gender 19

OSHA-Recordable Accidents by Type and Gender 19

Rates of OSHA-Recordable Events 20

OSHA-Recordable Rates by Age and Job Categories Among Women, All Diagnoses Combined..... 20

OSHA-Recordable Rates by Age and Job Categories Among Men, All Diagnoses Combined 20

Time Trends for OSHA-Recordable Events 20

Age-Adjusted Rates for All OSHA-Recordable Diagnoses Combined Among Women and Men by Job Category from 1993 to 2002..... 21

Glossary 22

Explanation of Diagnostic Categories 23

ICD-9-CM Codes 24

Introduction

The U.S. Department of Energy's (DOE) commitment to assuring the health and safety of its workers includes the conduct of illness and injury surveillance activities that provide an early warning system for health problems among workers. The Illness and Injury Surveillance Program monitors illnesses and health conditions that result in absences, occupational illnesses and injuries, and disabilities and deaths among current workers.

This report provides a summary of illness and injury surveillance data collected from the Fernald Environmental Management Project (FEMP) from January 1, 2002 through December 31, 2002. Illness and injury surveillance has been ongoing at Fernald since 1993. The data were collected by a coordinator at FEMP and submitted to DOE's Illness and Injury Surveillance Data Center at Oak Ridge Institute for Science and Education, where quality control procedures and preliminary data analyses were performed. The analyses were interpreted and the final report prepared by the DOE Office of Epidemiology and Health Surveillance.

The information presented in this report provides highlights of the data analyses conducted. Earlier surveillance reports and additional supporting tables are posted on the Office of Office of Epidemiology and Health Surveillance Web site (www.eh.doe.gov/health/epi/surv) or are available by request. The main sections of the report include: work force characteristics; absences due to illness or injury; and workplace injuries, illnesses, and deaths that were reportable to the Occupational Safety and Health Administration ("OSHA-

recordable" events); and disabilities and deaths among current workers. This report also includes sections on time trends that provide comparative information on the health of the work force from 1993 to 2002.

Note: In the figures and calculations that follow, percentages have been rounded to the nearest whole number.

DOE sites vary by mission, function, job classification, and worker exposures; therefore, comparisons of FEMP with other DOE sites should be made with caution. In addition, many factors can affect the completeness and accuracy of health information reported at the sites, thereby affecting the patterns of illness and injury observed.



Site Overview

The Fernald Environmental Management Project (FEMP), located 18 miles northwest of downtown Cincinnati, Ohio, once produced pure uranium metal products used in various U.S. defense programs. Construction began in 1951 in the midst of the Cold War era. Production operations started in 1953 and were suspended in July 1989. FEMP was originally called the Feed Materials Production Center because it produced “feed” in the form of purified uranium metal for use by other DOE sites that made nuclear weapons. The site was designed as a large-scale, integrated facility capable of converting uranium ore and recycled material into uranium metal through a series of chemical and metallurgical conversions. These activities resulted in contamination with radioactive wastes that included uranium tailings emitting radon gas, thorium, and radium, as well as other hazardous materials such as heavy metals, barium, and asbestos.



In November 1989, the site was added to the Superfund National Priority List, which requires site cleanup and remediation activities. Since December 1992, Fluor Fernald has managed the cleanup of the 1,050-acre FEMP; in November 2000, DOE awarded to Fluor Fernald a 10-year closure contract for the Fernald site that provided the framework for the final cleanup of the site. The final

removal of soil and debris from the Southern Waste Unit at Fernald occurred in June 2001. The cleanup of this 26-acre area was a milestone because much of the contamination lay in direct contact with the Great Miami Aquifer.

Together with the DOE, Fluor Fernald completed construction of 2 new disposal cells in November 2002. The new cells, nearly 40 percent complete, are part of a 7-cell, three-quarter mile long, On-Site Disposal Facility. The disposal facility will be among the last projects finished since it will remain open to receive the last of Fernald’s contaminated material and soils.



In December 2002, the new Radon Control System (RCS) was introduced at Fernald. The RCS provided a major step forward in efforts to safely retrieve and dispose of 8,900 cubic yards of radioactive waste. The system is designed to draw radon gas out of the headspace area in each of Fernald’s K-65 silos and reduce the concentration of the gas by 95 percent.




DOE and Fluor Fernald are on schedule to complete cleanup and safe closure of the site by December 2006.

The Fernald Work Force - 2002

A total of 1,777 Fernald employees were included in illness and injury surveillance in 2002, a decrease of 16 workers from 2001. The age and gender distribution of the 2002 work force is shown in Figure 1. There were 540 (30 percent) women and 1,237 (70 percent) men in the work force. The average age of male and female Fernald workers was 46 years and 43 years, respectively. The majority (87 percent) of the workers were White, 11 percent were African Americans, and the remaining 2 percent were Asians, Hispanics, and others.

Figure 1. The Work Force by Gender and Age



The distribution of workers by gender and job category is shown in Figure 2. As reported by Fernald, individual job titles were grouped into 10 job categories. This is because there were either too few workers or health events within a particular job title, thereby limiting the type of analyses that could be conducted. Men and women were not distributed equally among the various job categories. Half of the women were in the Professional (31 percent) and Clerical (19 percent) groups. Less than 1 percent of men

were in the Clerical job category. The largest percentages of male workers were in the Professional (18 percent) and Craft and Repair (18 percent) categories.

Figure 2. The Work Force by Job Category and Gender

Job Category	Women	Men
Management	9 2%	61 5%
Administration	12 2%	78 6%
Professional	169 31%	227 18%
Engineering, Scientific, & Health Care	52 9%	143 12%
Technical Support	58 11%	176 14%
Clerical	100 19%	3 <1%
Service	86 16%	175 14%
Security	3 1%	22 2%
Craft & Repair	12 2%	227 18%
Nuclear Specialties	39 7%	125 10%

A Note to the Reader:

Prior to the Year 2001 report, illness and injury surveillance at FEMP examined illness and injury absences of 5 or more consecutive workdays (also referred to as “5-day absences”). This approach is based on DOE Order 440.1 that requires contractor management to notify Occupational Medicine when a worker has been absent for 5 or more consecutive workdays. Eligible health events would also include those with an absence on a Friday that continued through Tuesday, with the length of that absence including the weekend.

As indicated in Order 440.1, all illnesses and injuries due to a work-related incident must be reported. Non-occupational illnesses and injuries that

involve absences of fewer than 5 days do not routinely require a medical clearance for return to work and, as noted above, have been excluded from these analyses until now. Beginning with the year 2001, FEMP chose to include absences of shorter duration. Twelve absences of less than 5 days in duration were reported by 12 employees (5 females and 7 males). Rates of OSHA-recordable events, reportable regardless of whether or not an absence is involved, will in general not be affected by this change in reporting.

Number and Length of Absences


Illness and injury surveillance examines absences from work. It is based on DOE Order 440.1 that requires contractor management to notify Occupational Medicine when a worker has been absent for 5 or more consecutive workdays. If an absence on a Friday continues through Tuesday, the length of that absence includes the weekend. Starting with the 2001 data, all reported absences are now included in the data collection and analyses, regardless of the length of absence. Twelve absences of less than 5 days duration were reported by 12 employees in 2001. In 2002, 12 of these absences were reported by 12 employees (5 females and 7 males): 4 Service workers, 3 Technical Support workers, 2 Professional workers, and 1 worker each from the Engineering, Scientific, and Health Care, Craft and Repair, and Nuclear Specialties occupational groups. All illnesses and injuries due to a work-related incident must be reported. Certain types of health events were excluded from the analyses. These include 5 women with 5 reported absences related to normal pregnancy and 1 man and 2 women with 3 reported absences due to elective surgical procedures not related to the treatment of an illness or injury.

Throughout this report, analyses take gender, age, and job category into account because the risk of illness and injury varies by these factors.



As shown in Figure 3, the rate of absence due to illness or injury varied by gender and age. There were 192 absences among 540 women resulting in an absence rate of 36 per 100 workers (192/540). Among the 1,237 men, there were 225 absences resulting in an absence rate of 18 per 100 workers (225/1,237). Among men, the rate decreased with age until age 50 when it increased. Among women, there was not a relationship between age and absence rate.

Figure 3. Absence Rate by Gender and Age



The average length of absence by gender and age is shown in Figure 4. A total of 17,005 calendar days of work (9,523 days for men and 7,482 days for


women) was lost at Fernald in 2002 due to reported illness or injury. The average length of absence was 42 days for men and 39 days for women. The youngest workers had the shortest average length of absence among men and women. Among men, the average length of absence varied little for workers 30 years of age and older. Women aged 30-49 had absences of similar average length. The length of absence among the oldest women was over 3 times as long as the youngest women.

Figure 4. Number of Days Absent by Gender and Age

Gender	Age	Number of Absences	Number of Days Absent	Average Number of Days Absent
Women	16-29	10	147	15
	30-39	76	2,897	38
	40-49	57	1,910	34
	50+	49	2,528	52
	Total	192	7,482	39
Men	16-29	23	791	34
	30-39	62	2,606	42
	40-49	51	2,197	43
	50+	89	3,929	44
	Total	225	9,523	42

As shown in Figure 5, the rate of absence due to illness or injury varied by job category for men and women. Men in the Clerical group did not report any absences in 2002. Women had higher rates of absence compared with men. Nuclear Specialties workers had the highest absence rate among male workers (35 per 100 workers); this same job category has had the highest rate of absence among men since 1995. Among women, Security workers had the highest absence rate (100 per 100 workers), however, there were only 3 workers in that group.


Figure 5. Absence Rate by Job Category and Gender



The average duration of absence by job category and gender is shown in Figure 6. Within a job category, men tended to have a longer average length of absence than women. Women in the Clerical group and men in the Management group had the longest average lengths of absence, 56 days for women and 53 days for men.



Figure 6. Average Duration of Absence by Job Category and Gender



Diagnostic Categories

Illness and injury surveillance monitors *all* illnesses and injuries among active workers because it is not always possible to determine what health effects are due to occupational exposures and what are due to other causes. Most illness and injury diagnoses were reported to the occupational medicine clinic by workers who required return-to-work

clearances. An absence due to illness or injury may involve more than 1 diagnosis, and illness and injury surveillance includes all reported diagnoses. In addition, the OSHA 200 Log provides information on recorded occupational illnesses and injuries whether or not they involve absences.

This report organizes illness and injury categories based on a standard reference, the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*. This reference is used to classify health events for statistical purposes. You can find specific health conditions in the Explanation of Diagnostic Categories.

The number of reported diagnoses categorized according to the ICD-9-CM and number of lost calendar days (may include weekends and holidays) are presented in Figure 7a. Lost calendar days for each absence are counted more than once when multiple diagnoses occur in different diagnostic categories for the same absence. There were 317 diagnoses reported by female and 338 diagnoses reported by male Fernald employees in 2002. Of these diagnoses, 18 were reported for absences less than 5 days in duration (8 reported by women and 10 reported by men). Five (28 percent) of the 18 diagnoses were for unspecified symptoms, while the remaining 13 diagnoses were spread over 7 diagnostic categories.



Figure 7a. Number of Diagnoses and Lost Calendar Days by Diagnostic Category (Categorized by ICD-9-CM) and Gender

Diagnostic Category	Women		Men	
	Number of Diagnoses	Number of Lost Calendar Days	Number of Diagnoses	Number of Lost Calendar Days
Benign Growths	7	284	3	30
Blood	0	0	0	0
Cancer	6	403	2	45
Digestive	18	430	35	970
Endocrine/ Metabolic	6	659	8	273
Existing Birth Condition	3	101	1	40
Genitourinary	28	672	8	183
Heart/ Circulatory	12	240	32	1,205
Infections/ Parasites	4	84	7	168
Injury	35	1,338	71	2,693
Miscarriage	1	16	NA	NA
Muscles & Skeleton	56	2,190	79	3,343
Nervous System	30	1,234	20	646
Psychological	38	1,560	19	449
Respiratory	32	881	26	305
Skin	7	140	2	50
Unspecified Symptoms	34	1,259	25	631

Note: Lost calendar days for each absence are counted more than once when multiple diagnoses occur in different diagnostic categories for the same absence.

The most frequently reported diagnoses varied little by gender. Among women, 41 percent of all reported diagnoses were due to conditions of the muscles and skeleton (18 percent), psychological disorders (12 percent), and injuries (11 percent). Among male workers, 54 percent of all reported diagnoses were due to muscles and skeleton conditions (23 percent), injuries (21 percent), and digestive conditions (10 percent). Major contributors to these diagnostic categories are shown in Figures 7b and 7c for women and men, respectively.

Among men, these diagnoses did not vary much by age. Men aged 30-39 reported more nervous system disorders, while workers 50 years of age and older reported more heart/circulatory diseases. Five of the 8 nervous system diagnoses reported by 6 men aged 30-39 years old were for carpal tunnel syndrome. Sixteen men 50 years old or greater reported 24 heart/circulatory diagnoses: 3 diagnoses for hypertension, 14 for ischemic heart disease (restricted blood flow to an artery), and the remainder for a variety of other heart/circulatory problems.

Among women, the most frequently reported diagnoses were not as consistent among the various age groups. Women less than 40 years old frequently reported genitourinary conditions and respiratory disorders. Among workers 50 years of age and greater, nervous system conditions were commonly reported; 5 of the 9 diagnoses were related to vision.

Figure 8 shows the frequency of reported diagnoses by job category for men and women. The types of diagnoses varied among the job categories, but muscles and skeleton conditions and injuries appeared often among both men and women.



Figure 7b. Common Diagnoses Among Female Workers in 2002




Figure 7c. Common Diagnoses Among Male Workers in 2002




Figure 8. Most Frequently Reported Diagnoses by Job Category and Gender

Job Category	Men	Women
Management	Heart/Circulatory (1) Injury (1)	Muscles & Skeleton (2)
Administration	Cancer (1) Digestive (1) Heart/Circulatory (1) Injury (1)	Digestive (1)
Professional	Muscles & Skeleton (10) Heart/Circulatory (9) Psychological (8)	Muscles & Skeleton (16) Injury (14) Nervous System (11)
Engineering, Scientific, & Health Care	Muscles & Skeleton (6) Nervous System (4) Unspecified Symptoms (3)	Unspecified Symptoms (5) Muscles & Skeleton (4) Psychological (3)
Technical Support	Muscles & Skeleton (7) Heart/Circulatory (4) Endocrine/Metabolic (1) Genitourinary (1) Injury (1) Respiratory (1) Unspecified Symptoms (1)	Unspecified Symptoms (5) Genitourinary (4) Injury (4) Respiratory (4)
Clerical	None	Muscles & Skeleton (9) Injury (6) Unspecified Symptoms (6)
Service	Injury (23) Muscles & Skeleton (17) Digestive (8)	Psychological (14) Muscles & Skeleton (13) Nervous System (11)
Security	Respiratory (2) Digestive (1) Genitourinary (1)	Genitourinary (2) Endocrine/Metabolic (1) Infections/Parasites (1) Injury (1) Muscles & Skeleton (1) Skin (1)
Craft & Repair	Injury (26) Muscles & Skeleton (23) Heart/Circulatory (13)	Muscles & Skeleton (2) Nervous System (2) Heart/Circulatory (1) Infections/Parasites (1) Respiratory (1)
Nuclear Specialties	Muscles & Skeleton (16) Injury (14) Unspecified Symptoms (7)	Muscles & Skeleton (9) Psychological (8) Digestive (7) Respiratory (7)

Note: Numbers in parentheses represent the number of reported diagnoses.

Rates of Disease Occurrence

A Word about Rates: The previous section considered the **number** of absences and health conditions among various worker groups. For example, Figure 7a shows that men reported 71 diagnoses and women reported 35 diagnoses involving injuries during 2002. Men, therefore, reported twice as many injuries as women. As there were more than twice as many men as women at Fernald, it seems reasonable to expect more injuries among men than women. Does this mean that men were at greater risk of injuries compared with women in 2002? To correctly answer the question, the total number of men and women in the work force must be considered. To compare risk among men and women, it is necessary to calculate the injury rate for each gender. Rates are calculated by dividing the number of injury diagnoses in a given gender by the total number of employees of that gender. Multiply this number by 1,000 to get the diagnosis rate per 1,000 workers.

For example:

$$71 \text{ injury diagnoses} \div 1,237 \text{ men} = .057 \times 1,000 = 57 \text{ injury diagnoses per } 1,000 \text{ men}$$

$$35 \text{ injury diagnoses} \div 540 \text{ women} = .065 \times 1,000 = 65 \text{ injury diagnoses per } 1,000 \text{ women}$$

Comparing these rates now correctly suggests that the rate of reported absences due to injuries among women was higher than the rate for men. They are called **crude rates** because they do not account for possible differences between men and women such as age and other factors that might affect the individual's risk of having an injury. Because age is so strongly related to the risk of disease and injury, epidemiologists almost always take age into account when comparing groups. This is done by using age-specific categories or by statistical methods of adjustment.

The diagnosis rate, also called the illness and injury rate, is the number of occurrences of a given disease or health condition observed over the course of a year per 1,000 workers at risk of getting that condition (see shaded box). One health condition, arthritis for example, may result in several absences over a year. Conversely, 1 absence may be associated with multiple diagnoses (e.g., the flu and a sprained wrist) recorded on the return-to-work form.

In the following set of analyses, the 4 age groups were collapsed into 2 groups: workers less than 50 years of age and those 50 or older. In addition, the 10 occupational categories were combined into 5 larger groups. These groups were collapsed to ensure that the number of diagnoses in each group was large enough to analyze. The rates of all illnesses and injuries combined are presented in Figure 9. Four groups of diagnoses of particular interest to workers are presented in Figure 10: cancer, heart/circulatory system, respiratory system, and injury.

Figure 9. Rates for All Illnesses and Injuries Combined by Job Category, Gender, and Age






Diagnostic Category	Rate per 1,000			
	Job Category	Age	Men	Women
	Management/ Administration/ Professional	<50	163	439
		50+	153	549
	Engineering, Scientific, & Health Care/ Technical Support	<50	63	376
		50+	205	412
	Clerical	<50	0	373
		50+	0	485
	Service/Security/ Craft & Repair	<50	380	1,014
		50+	527	704
	Nuclear Specialties	<50	681	1,462
		50+	226	1,000

Figure 10. Rates for Selected Diagnostic Categories by Job Category, Gender, and Age

Diagnostic Category	Rate per 1,000			
	Job Category	Age	Men	Women
	Management/ Administration/ Professional	<50	0	0
		50+	6	78
	Engineering, Scientific, & Health Care/ Technical Support	<50	0	11
		50+	0	0
	Clerical	<50	0	0
		50+	0	30
	Service/Security/ Craft & Repair	<50	4	0
		50+	0	0
	Nuclear Specialties	<50	0	0
		50+	0	0

Diagnostic Category	Rate per 1,000			
	Job Category	Age	Men	Women
	Management/ Administration/ Professional	<50	0	0
		50+	67	0
	Engineering, Scientific, & Health Care/ Technical Support	<50	10	43
		50+	27	59
	Clerical	<50	0	0
		50+	0	61
	Service/Security/ Craft & Repair	<50	14	27
		50+	68	74
	Nuclear Specialties	<50	28	38
		50+	0	0

Diagnostic Category	Rate per 1,000			
	Job Category	Age	Men	Women
	Management/ Administration/ Professional	<50	5	36
		50+	12	39
	Engineering, Scientific, & Health Care/ Technical Support	<50	10	43
		50+	9	0
	Clerical	<50	0	45
		50+	0	0
	Service/Security/ Craft & Repair	<50	43	108
		50+	41	111
	Nuclear Specialties	<50	14	154
		50+	19	231

Diagnostic Category	Rate per 1,000			
	Injury	Job Category	Age	Men
	Management/ Administration/ Professional	<50	25	50
		50+	12	137
	Engineering, Scientific, & Health Care/ Technical Support	<50	0	54
		50+	9	0
	Clerical	<50	0	60
		50+	0	61
	Service/Security/ Craft & Repair	<50	127	81
		50+	95	0
	Nuclear Specialties	<50	153	115
		50+	57	77

Rates of all illnesses and injuries combined tended to be greater for workers 50 years of age and older compared with younger workers among women. Among men, the rates were not related to age. Rates for female employees were higher than those for males in the same job category. The highest illness and injury rates for all employees were among individuals classified as Nuclear Specialties. This occupational group has had the highest rates since 1995.

Cancer rates presented in this report are based on reported absences due to cancer. A worker may experience several periods of absence from 1 cancer diagnosis due to medical complications or treatment regimens. The cancer rates in this report are *not* comparable to the *incidence rates* frequently published in many articles on cancer with which you may be familiar. Cancer *incidence rates* are based on the number of new cancer cases diagnosed within a given time, usually a year.

Eight absences related to cancer were noted with 6 diagnoses reported by 4 women and 2 diagnoses reported by 2 men. Two of the workers reporting cancer in 2002 reported the same cancer during the previous 9 years. The likelihood that an individual in the U.S.

will develop cancer increases with age; our data reflect this observation. Only 2 of the 6 workers reporting cancer were less than 50 years old.

Not unexpectedly, older male and female workers tended to have higher rates of heart/circulatory problems. Twenty-four of the 32 diagnoses reported by men were among workers aged 50 and older; 3 diagnoses were for hypertension and 14 involved ischemic heart disease (restricted blood flow through an artery). Men categorized as Service/Security/Craft and Repair workers had the highest rate of heart/circulatory disorders. Ten of the 14 diagnoses in this group involved hypertension or ischemic heart disease. Among all women, there were 12 heart/circulatory diagnoses; 2 involved hypertension or ischemic heart disease. Craft and Repair workers were twice as likely as other workers to report a heart/circulatory diagnosis.

Women workers aged 50 and older tended to have higher respiratory diagnosis rates compared with younger women. No relationship with age was observed among men. Women had higher rates of respiratory disease than did men with the exception of workers 50 years and older in the Engineering, Scientific, and Health Care/Technical Support group.

Both men and women younger than 50 years tended to have higher rates of injuries than did older workers. The highest injury rates in 2002 among women were in the Nuclear Specialties group. Among men, the highest rate occurred among Service/Security/Craft and Repair workers. Craft and Repair and Nuclear Specialties workers were twice as likely as other groups to report an injury. An increased risk has been seen in the Nuclear Specialties group since 1995. Craft and Repair workers

were almost 9 times more likely to report a dislocation. Nuclear Specialties workers were almost 4 times more likely to report a back sprain or strain. As in 2001, Service workers were 3 times more likely to report a sprain or strain other than to the back.

In other analyses, the risk of illness and injury among workers classified in 1 job category was compared with workers in the other 9 job categories. Service, Craft and Repair, and Nuclear Specialties workers were almost twice as likely to report an illness or injury compared with all other groups. These



same occupational groups have been at a similar increased risk since 2000. They were at increased risk for other illnesses and injuries as well. The risk of psychological disorders was almost 3 times higher and muscles and skeleton disorders almost 2 times higher among Service workers compared with other occupational groups. Craft and Repair workers had twice the risk of muscles and skeleton conditions. Endocrine/metabolic disorders were elevated 7-fold, and psychological disorders and conditions of the digestive system and muscles and skeleton were elevated 2- to 3-fold among workers in the Nuclear Specialties group compared with other workers. Among Security workers, infections/parasites and genitourinary disorders were about 9 times more likely to be reported compared with other workers.

Time Trends

Why Are Rates Age-Adjusted?

The injury and illness rates in this section of the report are **age-adjusted**. Differences in the age composition among groups of workers are taken into consideration in the analyses, and 1 rate is calculated for an entire group. This allows us to make comparisons between groups of different ages. Age-adjusted rates are calculated using the age distribution of the 1970 U.S. population as a reference.

In 1995, Fernald began to report job categories that were not available in 1993 and 1994. In order to examine time trends from 1993 to 2002, some job categories used in 1995 through 2002 were combined to reflect the broader categories used in earlier years. The accompanying table shows how the categories were combined:

1995 - 2002 Job Category	1993 and 1994 Job Category
Management	Office Management and Administration
Administration	Office Management and Administration
Professional	Other Management and Administration
Engineering, Scientific, and Health Care	Engineering, Scientific, and Health Care
Technical Support	Technical Support
Clerical	Office Management and Administration
Service	Service
Security	Service
Craft and Repair	Craft and Repair
Nuclear Specialties	Nuclear Specialties

There are 10 years of illness and injury surveillance data for Fernald workers. It is important to note that the age-adjusted rates for the years 1993 and 1994 presented in this report differ from rates presented in the 1993 and 1994 *Annual Epidemiologic Surveillance Reports* due to the exclusion of absences resulting from maternity leave.

Age-adjusted rates for all diagnoses combined are shown in Figure 11. There was a steady increase in the rates for men and women from 1993 through 1997. This was followed by a decline or leveling off until a resumption in the increase in 2000 for women. The increase seen in the 2000 rates continued through 2002 among women, but among men, the rates



decreased. Figure 12 shows age-adjusted rates for selected diagnostic categories. The most notable trends over the 10-year period have been increases in muscles and skeleton conditions and injuries for men and women. However, since 1999 the injury rates have declined among men and women.

Figure 13 shows the age-adjusted rates for illnesses and injuries by job category. The rates of diagnoses among men and women showed little change from 2001 to 2002 for most job categories. The large changes in the rates for women in the Technical Support, Service, Craft and Repair, and Nuclear Specialties groups are the result of small numbers of female workers in each of these groups. Rates calculated for small groups tend to fluctuate more than those based on larger numbers of workers.



Figure 11. Age-Adjusted Rates for All Diagnoses Combined Among Women and Men from 1993 to 2002




Figure 12. Age-Adjusted Rates for Selected Diagnostic Categories Among Women and Men from 1993 to 2002





Figure 13. Age-Adjusted Rates for All Diagnoses Combined Among Women and Men by Job Category from 1993 to 2002



Note: A job category for “Other” workers appeared in 1993 only; the 36 employees in this job category were excluded from this figure.

Sentinel Health Events for Occupations

A sentinel health event for occupation (SHEO) is a disease, disability, or death that is likely to be occupationally related. Its occurrence may serve as a warning signal that material substitution, engineering control, personal protection, or medical care may be required to reduce the risk of illness or injury among the work force. Sixty-four medical conditions associated with workplace exposures from studies of many different industries have been identified as sentinel health events. Although sentinel health events may indicate an occupational exposure, many may result from non-occupational exposures. Due to this uncertainty, sentinel health events are assessed in 2 categories:

Definite Sentinel Health Events:

Diseases that are unlikely to occur in the absence of an occupational exposure. Asbestosis, a lung disease resulting from exposure to asbestos, is an example.

Possible Sentinel Health Events:

Conditions such as lung cancer or carpal tunnel syndrome may or may not be related to occupation. Detailed occupational and non-occupational information is required to determine the work-relatedness of the illness. For example, lung cancer may result from asbestos exposure or smoking. Carpal tunnel syndrome may result from a job requiring typing or from a hobby such as playing the piano.

Three definite and 14 possible sentinel health event diagnoses were identified among the 655 reported diagnoses (Figure 14). The 3 definite sentinel health diagnoses were reported as 3 events among 3 male workers (1 Management, 1 Craft and Repair, and 1 Nuclear Specialist). These events included 1 fracture, 1 sprained knee, and 1 bruise; they accounted for 108 days absent. The 14 possible sentinel health events, reported by 12 workers (6 women and 6 men) included 1 Raynaud’s syndrome, 1 disorder of the bone and cartilage of the knee, and 12 absences for carpal tunnel syndrome. The carpal tunnel events accounted for 530 days absent.



Figure 14. Characteristics of SHEOs by Gender

	Total Number of SHEO Diagnoses		Total Number of Days Absent	
	Men	Women	Men	Women
Definite	3	0	108	0
Possible	7	7	285	311
Total	10	7	393	311

Disabilities Among Active Workers

No disabilities were reported by FEMP in 2002.

Deaths Among Active Workers

Two deaths (1 female worker and 1 male worker) occurred among FEMP workers in 2002. They were due to brain cancer and a heart attack. Both workers were aged 50 or older.

OSHA-Recordable Events

The Occupational Safety and Health Administration (OSHA) requires employers to maintain a record of occupational illnesses and injuries that have occurred among employees and to make that information available to OSHA upon request. Employers maintain the information from these OSHA-recordable events in the OSHA 200 Log. OSHA-recordable events differ from health events captured through return-to-work clearances in at least 2 important respects: 1) they do not necessarily result in days lost from work, and 2) they are usually accompanied by a specific determination that they are work-related.

The rates of OSHA events by gender and age are shown in Figure 15. Rates were higher among women than among men in most age groups. The rates of



OSHA-recordable events by job category and gender are shown in Figure 16. Seventeen men and 11 women reported 28 OSHA-recordable events, with 15 diagnoses among women and 22

diagnoses among men (Figure 17). All the diagnoses reported by women and 86 percent of the diagnoses among men were injuries. Sprains and strains were the most frequently reported injuries, followed by fractures and unspecified injuries for women and open wounds and bruises for men.

Figure 15. OSHA-Recordable Events by Gender and Age




Figure 16. OSHA-Recordable Events by Job Category and Gender




Figure 17. OSHA-Recordable Diagnoses by Diagnostic Category and Gender

Diagnostic Category	Gender	
	Women	Men
Muscles & Skeleton	0	1
Nervous System	0	1
Skin	0	1
Injury	15	19
Fractures – Upper Limb	2	1
Fractures – Lower Limb	1	1
Back Sprains & Strains	1	4
Other Sprains & Strains	7	2
Internal Injuries – Thorax, Abdomen, Pelvis	0	1
Open Wounds – Head, Neck, Trunk	0	1
Open Wounds – Upper Limb	0	2
Open Wounds – Lower Limb	0	1
Superficial Injuries	0	2
Bruises	1	3
Foreign Bodies Entering Orifice	0	1
Unspecified Injuries	2	0
Adverse Reactions to Non-Medical Substances	1	0

Figure 18 shows that 27 of the 28 events resulted from accidents: 8 falls, 2 motor vehicle accidents, 1 insect sting, and 16 “other accidents” (3 reported by women and 13 reported by men). The 3 other accidents reported by women and 3 (23 percent) of the 13 other accidents reported by men were from overexertion and strenuous movements. The 10 remaining other accidents reported by men included 3 struck by an object, 3 caught between objects, 2 cutting/ piercing instrument /object, and 1 each for hot, corrosive, or caustic material/ steam and repetitive trauma. Overall, this distribution showed no evidence of a pattern involving any particular kind of accident. The accidents resulted in 36 diagnoses (15 reported by women and 21 reported by men): 14 sprains and strains, 5 fractures, 4 bruises, 4 open

wounds, 2 unspecified injuries, and 1 each for eye disorder, abscessed toe, tendonitis of the wrist, infected blister (foot), foreign body in the eye, internal injury to chest, and insect bite. The 1 event not recorded as an accident had a diagnosis of abrasion of the cornea. This distribution of diagnoses was not unusual for Fernald, compared with the site’s experience in previous years.

Figure 18. OSHA-Recordable Accidents by Type and Gender

Accident Category	Gender	
	Women	Men
	Number of Accidents	Number of Accidents
Motor Vehicle Traffic	0	2
Falls	7	1
Natural/Environmental Factors	1	0
Other Accidents	3	13
Struck by an Object	0	3
Caught Between Objects	0	3
Cutting/Piercing Instrument/Object	0	2
Hot, Corrosive, or Caustic Material/Steam	0	1
Overexertion/Strenuous Movements	3	3
Repetitive Trauma	0	1
Total	11	16

A total of 92 days were lost and 687 days were restricted among women. Men had a total of 83 lost days and 120 restricted days due to accidents. The 4 events among women with the largest number of days lost and restricted were the result of sprains and strains to the leg and a fractured arm. Three of these events were caused by falls.

There were no OSHA events reported by women aged 16 to 29 years old. Among both men and women, no OSHA events were reported by Administration or Technical Support workers. These occupational groups also had no OSHA events in 2000 and 2001. In addition,

women in the Management, Professional, and Engineering, Scientific, and Health Care job categories and men in the Clerical and Security groups reported no OSHA events in 2002.

Rates of OSHA-Recordable Events

The rates of all OSHA-recordable events by age and job categories and gender are shown in Figures 19 and 20. Service/Security/Craft and Repair workers had the highest rate of OSHA events for both men and women in 2002.

Figure 19. OSHA-Recordable Rates by Age and Job Categories Among Women, All Diagnoses Combined






Figure 20. OSHA-Recordable Rates by Age and Job Categories Among Men, All Diagnoses Combined



Time Trends for OSHA-Recordable Events

The age-adjusted rates for all OSHA-recordable diagnoses combined from 1993 to 2002 by job category and gender are shown in Figure 21. During the 10-year period, the overall rates for OSHA-recordable events did not change greatly for most of the job categories among men and women. An overall decline in the rate among men in the Craft and Repair group was observed between 1994 and 2001, but the trend was broken by a sharp increase in 2002. In summary, we noted no significant changes in the injury rates from OSHA-recordable events from 1993 to 2002.

Figure 21. Age-Adjusted Rates for All OSHA-Recordable Diagnoses Combined Among Women and Men by Job Category from 1993 to 2002



Note: A job category for “Other” workers appeared in 1993 only; the 36 employees in this job category were excluded from this figure. The 1993 Craft & Repairs rate for women was truncated to 200 (□) for graphical presentation. The actual rate was 273.

Glossary

Adjustment: A mathematical procedure for rates in which the effects of differences of a characteristic (such as age or gender) between groups have been removed. The purpose of adjustment is to allow comparisons between 2 or more groups with the effect of the differences for the characteristic removed.

Age-Adjusted Rate: A rate that has been mathematically adjusted to account for the effects of differences in the age composition between groups.

Age-Specific Rate: A rate that is calculated for a specific age group (e.g., 16 to 29 years old). Only people in the specific age group are included in the calculation of the rate.

Confidence Interval: A range of values determined by the degree of random variability in the data. The width of the confidence interval is affected by the size of the group being studied and how often the event whose true value is sought occurs. Generally, as the size of the group or the frequency of the event increases, the width of the confidence interval decreases. The level of confidence, for example a 95 percent confidence level, indicates the percentage (e.g., 95 percent) of time that the true value is expected to fall within the confidence interval if the mathematical procedure is repeated 100 times.

Demographics: Characteristics of human populations related to their size, density, age distribution, and vital status.

Diagnosis (diagnoses): Identification of a disease or health condition from signs and symptoms.

Diagnosis Rate: The number of occurrences of a given disease or health condition observed during a given time period per the number of workers at risk of getting that disease during that time period. It is usually multiplied by 100 or 1,000 to produce a rate expressed as a convenient number.

Diagnostic Category: A particular type of disease, a group of related health conditions, or diseases that all affect the same organ system.

Epidemiologic Surveillance: The ongoing evaluation of the health of a human population which is based on the collection and interpretation of demographic and health information for that population.

Epidemiology: The study of the distribution and determinants of diseases and health conditions in human populations.

ICD-9-CM Code: An abbreviation for the *International Classification of Diseases, 9th Revision, Clinical Modification*. An internationally accepted standardized system for the classification of disease and health data collected from medical records.

OSHA: An acronym for the Occupational Safety and Health Administration.

OSHA Event: An abbreviation used throughout this report for an OSHA-Recordable Event.

OSHA-Recordable Event: An accident that occurs on the job and involves fatalities (regardless of time between injury and death), time lost from work, transfer of employment, medical treatment other than first aid, loss of consciousness, or restriction of work or motion. Also included is any diagnosed occupational health event reported to the employer that is neither fatal nor results in workdays lost. By law, these events are recordable in the OSHA 200 Log.

Person-Year: A unit of measurement combining the number of people being studied with the time that each was observed equivalent to 1 person followed for 1 year. For example, 5 people followed for 1 year contribute 5 person-years, as do 10 people each followed for half a year.

Relative Risk: The ratio of the occurrence of a disease or health condition in 1 group compared with the rate of occurrence of that same disease or health condition in another group.

Explanation of Diagnostic Categories

Throughout this report, health conditions have been grouped into a number of diagnostic categories which come from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). For the text of this report the categories are abbreviated to make the report easier to read. The following table lists the abbreviated categories used throughout the annual report and the corresponding ICD-9-CM codes found in the supporting tables.

Abbreviated Categories Used in the Annual Report	ICD-9-CM Codes
Benign Growths	210-229 235-239
Blood	280-289
Cancer	140-208 230-234
Digestive	520-579
Endocrine / Metabolic	240-279
Existing Birth Conditions	740-759
Genitourinary	580-629
Heart / Circulatory	390-459
Infections / Parasites	001-139
Injury	800-999
Miscarriage	630-676
Muscles and Skeleton	710-739
Nervous System	320-389
Psychological	290-319
Respiratory	460-519
Skin	680-709
Unspecified Symptoms	780-799

ICD-9-CM Codes

All conditions	001-V82	All reported health events
Infectious and parasitic diseases	001-139	Diseases caused by bacteria, viruses, and parasites
• Intestinal infections	001-009	Infections of the bowel or gut
• Tuberculosis	010-018	TB in the lungs and other organs
• Zoonotic bacterial diseases	020-027	Bacterial diseases that animals transmit to humans
• Other bacterial diseases	030-041	Whooping cough, diphtheria, strep throat, and gangrene
• Human Immunodeficiency Virus (HIV) infection	042	AIDS
• Poliomyelitis and other non-arthropod diseases of the central nervous system	045-049	Viral meningitis (swelling of the layers covering the brain and spinal cord); viral encephalitis (swelling of the brain); and polio
• Viral diseases accompanied by exanthem	050-057	Diseases accompanied by rashes or blisters like chickenpox, measles, shingles, and herpes
• Arthropod-borne viral diseases	060-066	Encephalitis (swelling of the brain) caused by bites from virus-carrying ticks or mosquitoes
• Other diseases caused by viruses and chlamydiae	070-079	Viral hepatitis, mumps, rabies, and mononucleosis
• Rickettsioses and other arthropod-borne diseases	080-088	Rocky Mountain spotted fever, malaria, and lyme disease
• Other spirochetal diseases	100-104	Trench mouth and Weil's disease (jaundice caused by coil-shaped bacteria)
• Mycoses	110-118	Athlete's foot; fungal infections of fingernails and toenails; and thrush
• Helminthiases	120-129	Pinworms, tapeworms, roundworms, and whipworms

• Other infectious and parasitic diseases	130-136	Lice, chiggers, scabies, and mites
• Late effects of infectious or parasitic diseases	137-139	Side effects of TB, chickenpox, or polio even though the disease is no longer active
Malignant neoplasms	140-208, 230-234	All cancers, regardless of the part of the body affected
• Lip, oral cavity, and pharynx	140-149	Lip, mouth, throat, and tongue
• Digestive organs and peritoneum	150-159	Stomach, esophagus (tube that transports food to the stomach), intestines, colon, rectum, anus, liver, pancreas, and gallbladder
• Respiratory system and intrathoracic organs	160-165	Sinuses, throat, voice box, lungs, and heart
• Bone, connective tissue, skin, and breast	170-176	Bone, muscle, ligament, tendon, blood vessels, fat, skin, and breast
• Genitourinary organs	179-189	Kidney, bladder, and cervix, ovary, uterus, and prostate
• Other and unspecified sites	190-199	Eye, brain, and thyroid
• Lymphatic and hematopoietic tissue	200-208	Leukemia, lymphoma, Hodgkin's disease, multiple myeloma, lymphosarcoma, and reticulum cell sarcoma
• Carcinoma in situ	230-234	A cancer that is confined to the site of origin (has not spread to neighboring tissue)
Benign neoplasms and neoplasms of uncertain behavior and unspecified nature	210-229 235-239	Tumors that are not cancerous or do not exhibit cancerous behavior, regardless of the part of the body affected
Endocrine, nutritional, and metabolic diseases and disorders of the immune system	240-279	Diseases affecting the hormone secreting glands and organs. Overactive thyroid; underactive thyroid; vitamin deficiency; diabetes; gout; and problems affecting the antibody producing system

Disorders of the blood and blood forming organs	280-289	Anemia and hemophilia (excludes leukemia)
Mental disorders	290-319	Psychiatric diagnoses - Non-psychotic disorders: depression; anxiety, fear, and stress disorders; alcoholism; drug dependence; and eating disorders, such as anorexia; Psychotic disorders: dementia, schizophrenia, and manic depression
Diseases of the nervous system and sense organs	320-389	Huntington's chorea; Alzheimer's and Parkinson's disease; epilepsy; multiple sclerosis; migraine; diseases of the eye, such as cataract and glaucoma
• Inflammatory diseases of the central nervous system	320-326	Bacterial meningitis (swelling of the layers covering the brain and spine); bacterial encephalitis (swelling of the brain); and brain and spinal abscesses
• Hereditary and degenerative diseases of the central nervous system	330-337	Alzheimer's and Parkinson's disease, tremors, and Huntington's chorea
• Other disorders of the central nervous system	340-349	Multiple sclerosis (MS), cerebral palsy, epilepsy, and migraine
• Disorders of the peripheral nervous system	350-359	Nerve disorders of the face, carpal tunnel syndrome, muscular dystrophy
• Disorders of the eye	360-379	Inflammation and ulcers of the eye and eyelid; detached retina; pink eye; problems with tear ducts; glaucoma; and cataracts
• Diseases of the ear and mastoid process	380-389	Infections of the outer, middle, or inner ear; ringing of the ears; hearing loss

Diseases of the circulatory system	390-459	Rheumatic fever, heart murmurs, heart attacks, angina, hardening of the arteries, varicose veins, hemorrhoids, and phlebitis
• Acute rheumatic fever	390-392	High fever and joint pain with possible heart damage
• Chronic rheumatic heart disease	393-398	Long lasting swelling and damage to the heart which results from rheumatic fever
• Hypertensive disease	401-405	High blood pressure
• Ischemic heart disease (Restricted blood flow to the heart)	410-414	Heart attack and angina
• Diseases of pulmonary circulation	415-417	Blood clots in the lung and pulmonary aneurysm (bulge that develops in the wall of the pulmonary artery, which is the artery that carries blood to the lungs)
• Other forms of heart disease	420-429	Swelling of the inner lining, middle lining, or sac enclosing the heart; heart failure; and irregular heartbeat
• Cerebrovascular disease	430-438	Stroke, bleeding in the brain, and blockage or low blood flow in blood vessels of the brain
• Diseases of the arteries and capillaries	440-448	Hardening of the arteries; aneurysm (bulge that develops in the walls of arteries); and blood clots
• Diseases of the veins, lymphatics, and other circulatory system diseases	451-459	Phlebitis (swelling of a vein), thrombophlebitis (swelling of a vein which has a blood clot), varicose veins, and hemorrhoids

Diseases of the respiratory system	460-519	Colds, sinusitis, laryngitis, pneumonia, influenza, chronic bronchitis, asthma, and emphysema
• Acute respiratory infections	460-466	Colds, sore throat, sinus infections, swollen tonsils, and bronchitis
• Other diseases of the upper respiratory tract	470-478	Allergies, hay fever, sinus infections, bronchitis, and sore throat that continue for a long time
• Pneumonia and influenza	480-487	“The flu” and pneumonia caused by a bacteria or virus
• Chronic obstructive pulmonary diseases and allied conditions	490-496	Emphysema and asthma
• Pneumoconiosis and other lung diseases caused by external agents	500-508	Black lung; miners’ asthma; asbestosis; silicosis; berylliosis; and conditions caused by chemical fumes and vapors
• Other diseases of the respiratory system	510-519	Pleurisy (swelling of the lining of the lungs), collapsed lung, and respiratory failure
Diseases of the digestive system	520-579	Diseases affecting the teeth and mouth, salivary glands, digestive tract, and the abdominal cavity. Examples include dental abscess, ulcers, appendicitis, hepatitis (excluding viral hepatitis), cirrhosis of the liver, gallstones, pancreatitis, abdominal hernia, and intestinal polyps
• Diseases of the oral cavity, salivary glands, and jaw	520-529	Tooth problems (too many, too few, abnormal shape or size, cavities, bleeding gums, toothaches), and infections and swelling of the mouth, jaw, and tongue
• Diseases of the esophagus, stomach, and duodenum	530-537	Ulcers of the esophagus (tube that transports food to the stomach), stomach, and small intestine; indigestion; and uncontrollable vomiting

• Appendicitis	540-543	Swelling of the appendix (rupture, surgery, or both may result)
• Hernia of the abdominal cavity	550-553	Ruptures of the groin and diaphragm (muscle which separates the chest area from the lower part of the trunk)
• Non-infectious enteritis and colitis	555-558	Crohn's disease and swelling of the intestine and colon
• Other diseases of the intestines and peritoneum	560-569	Irritable bowel syndrome, blockage of the intestine, constipation, and diarrhea
• Other diseases of the digestive system	570-579	Diseases of the liver, gallbladder, and pancreas; hepatitis; blood in stool; and bleeding in the stomach and intestine
Diseases of the genitourinary system	580-629	Diseases affecting the kidneys, the prostate, and testes; benign breast diseases; infertility (male and female); diseases of the ovary; pelvic inflammatory disease; and menstrual disorders
• Nephritis, nephrotic syndrome, and nephrosis	580-589	Swelling of the kidney; swelling of the small blood vessels in the kidney; and kidney failure
• Other diseases of the urinary system	590-599	Swelling and infection of the kidney and bladder; kidney stones; and difficulty urinating
• Diseases of the male genital organs	600-608	Enlarged prostate; swelling of the scrotum and prostate; and abscess of the prostate
• Disorders of the breast	610-611	Benign tumors, cysts, and infections of the breast
• Inflammatory disease of the female pelvic organs	614-616	Swelling of the uterus, ovary, fallopian tubes, or cervix
• Other diseases of the female genital tract	617-629	Conditions associated with menopause and postmenopause; PMS; infertility; and cramps

Complications of pregnancy, childbirth, and the puerperium	630-676	Miscarriage; complications of pregnancy, such as hemorrhage; pregnancy-related high blood pressure; preeclampsia; and premature labor or other complications of labor
• Ectopic and molar pregnancy	630-633	Development of fetus outside the uterus and growth of cysts
• Other pregnancy with abortive outcome	634-639	Miscarriage and complications associated with miscarriage
• Complications mainly related to pregnancy	640-648	Abnormal bleeding and possible miscarriage; infections; high blood pressure caused by pregnancy; and premature labor
• Normal delivery, and other indications for care in pregnancy, labor, and delivery	650-659	Delivery requiring little or no assistance; multiple births; breech birth; and problems of the fetus or placenta which affect care of mother
• Complications occurring mainly in the course of labor and delivery	660-669	Long labor; unusually fast delivery; and abnormal bleeding after delivery
• Complications of the puerperium	670-676	Infections of the breast; blood clot in lung; and varicose veins
Diseases of the skin and subcutaneous tissue	680-709	Acne, cellulitis, sunburn, psoriasis, and seborrhea
• Infections of the skin and subcutaneous tissue	680-686	Abscesses, boils, hair-containing cysts, and pus-filled blisters
• Other inflammatory conditions of skin and subcutaneous tissue	690-698	Skin rashes caused by detergents, oils, greases, solvents, sun, food, drugs, or medicine
• Other diseases of the skin and subcutaneous tissue	700-709	Corns, calluses, heat rash, swollen hair follicles, acne, and ingrown fingernails and toenails

Diseases of the musculoskeletal system and connective tissue	710-739	Arthritis, systemic lupus erythematosus, ankylosing spondylitis, herniated intervertebral disk (“slipped disk”), lumbago, sciatica, rheumatism, tendonitis, and osteoporosis
• Arthropathies and related disorders	710-719	Arthritis; joint pain and stiffness; and other diseases of the connective tissue which supports and connects internal organs, forms bones and blood vessel walls, and attaches to bones
• Dorsopathies	720-724	Swelling of the spine; herniated, slipped, and ruptured disk; rheumatoid arthritis of the spine; lumbago; and sciatica
• Rheumatism, excluding the back	725-729	Swelling and degeneration of joints, muscles, tendons; tennis elbow; and bursitis
• Osteopathies, chondropathies, and acquired musculoskeletal deformities	730-739	Fracture caused by bone disease; osteoporosis; curvature of the spine; flat foot; hammer toe; and development of deformities of the nose, toes, feet, legs, arms, and hands
Congenital anomalies	740-759	Spina bifida; cleft palate; harelip; and various chromosomal anomalies, such as Klinefelter’s syndrome
Certain conditions originating in the perinatal period	760-779	Maternal high blood pressure; maternal malnutrition; ectopic pregnancy; breech birth; fetal malnutrition or slow growth; injuries related to birth trauma; and perinatal jaundice
Symptoms, signs, and ill-defined conditions	780-799	Blackout, chills, dizziness, fatigue, pallor, abnormal weight loss, undiagnosed chest pain, and heartburn

- Symptoms 780-789 Hallucinations, fainting, convulsions, dizziness, fatigue, fever, sleep disturbance, rash, headache, sore throat, chest pain, nausea, vomiting, and heartburn
- Non-specific abnormal findings 790-796 Abnormal x-ray, blood, stool, and urine test results
- Ill-defined and unknown causes of morbidity and mortality 797-799 Senility; asphyxia; respiratory arrest; nervousness; and unexplained death within 24 hours of onset of symptoms
- Injury and poisoning** 800-999 Dislocation of joints; sprains and strains of associated muscles; concussions; bruises; cuts; internal injuries from crushing, puncture, tearing, or blunt impact; burns; blisters; poisoning; frostbite; heatstroke; and complications of medical or surgical care
- Fractures, all sites 800-829 Cracks or breaks of any bone
- Dislocations 830-839 Separation of a bone from its normal socket or joint
- Sprains and strains of joints and adjacent muscles 840-848 Strains are injuries to muscle from overuse or stretching the muscle beyond its normal limit; sprains are injuries involving tearing or overextending the ligaments of a joint
- Intracranial injuries excluding those with skull fractures 850-854 Concussions; internal bruises; and bleeding within the head without a fracture of the bones of the skull
- Internal injuries of the thorax, abdomen, and pelvis 860-869 Bruising, crushing, tearing, or rupturing the chest, abdomen, and pelvis and the organs within these areas of the body
- Open wounds 870-897 Animal bites; cuts; lacerations; punctures; and amputations, excluding the arteries and veins

<ul style="list-style-type: none"> • Other injuries and late effects of external causes 	900-999	Miscellaneous injuries, including injuries to the arteries and veins; problems that occur an extended period of time after the injury has taken place ("late effects"); superficial bruises and abrasions; burns; post-injury shock; poisoning; toxic side effects of chemicals; heatstroke; electrocution; and altitude sickness
<p>Supplementary classifications related to personal or family history of disease</p>	V10-V19	Covers situations in which the person is not ill or injured but has a personal or family history of problems, such as cancer, mental illness, allergies, or arthritis that may affect his or her risk of illness
<p>Supplementary classifications related to health care for reproduction and child development</p>	V20-V28	Problems related to pregnancy, postpartum care, contraception, outcome of delivery, and physical development of child
<p>Contact with health services for reasons other than illness or injury</p>	V50-V59	Care for workers who have been treated previously for an illness or injury that is no longer present but who receive care to complete treatment or prevent recurrence

NOTES